

Galois Groups of Schubert Problems


USC Mathematics Colloquium

3 December 2014



Frank Sottile

sottile@math.tamu.edu



Work with: Ravi Vakil, Jan Verschelde, Anton Leykin,
Jacob White, Abraham Martín del Campo, Robert Williams,
Christopher Brooks, Zach Maril, and Aaron Moore.

Galois theory and the Schubert calculus

Galois theory originated from understanding symmetries of roots of polynomials. Later, Galois groups came to be understood as encoding all the symmetries of field extensions. Today, it is a pillar of number theory.

Galois groups also appear in enumerative geometry. This aspect is not well-developed, because of its subtlety and because Galois groups are very hard to determine.

I will describe a project to shed more light on Galois groups in enumerative geometry. It is focussed on Galois groups in the *Schubert calculus*, a well-studied class of geometric problems involving linear subspaces.

It is best to begin with examples.


The Problem of Four Lines

What are the lines m_i meeting four general lines l_1, l_2, l_3 , and l_4 ?

The Problem of Four Lines

What are the lines m_i meeting four general lines $l_1, l_2, l_3,$ and l_4 ?


The lines l_1, l_2, l_3 lie on a unique hyperboloid Q of one sheet, and the lines that meet l_1, l_2, l_3 form one ruling of Q .



The Problem of Four Lines

What are the lines m_i meeting four general lines $l_1, l_2, l_3,$ and l_4 ?

The lines l_1, l_2, l_3 lie on a unique hyperboloid Q of one sheet, and the lines that meet l_1, l_2, l_3 form one ruling of Q . Thus the solutions m_i are the lines in that ruling passing through the points of intersection $l_4 \cap Q$.




The Problem of Four Lines

What are the lines m_i meeting four general lines l_1, l_2, l_3 , and l_4 ?

The lines l_1, l_2, l_3 lie on a unique hyperboloid Q of one sheet, and the lines that meet l_1, l_2, l_3 form one ruling of Q . Thus the solutions m_i are the lines in that ruling passing through the points of intersection $l_4 \cap Q$.

Rotating the line l_4 180° around the point p interchanges the two solution lines m_1, m_2 .




The Problem of Four Lines

What are the lines m_i meeting four general lines l_1, l_2, l_3 , and l_4 ?

The lines l_1, l_2, l_3 lie on a unique hyperboloid Q of one sheet, and the lines that meet l_1, l_2, l_3 form one ruling of Q . Thus the solutions m_i are the lines in that ruling passing through the points of intersection $l_4 \cap Q$.

Rotating the line l_4 180° around the point p interchanges the two solution lines m_1, m_2 .



This shows that

The Galois group of the problem of four lines is the symmetric group \mathcal{S}_2 .

A Problem with Exceptional Geometry

Q: What 4-planes H in \mathbb{C}^8 meet four general 4-planes K_1, K_2, K_3, K_4 in a 2-dimensional subspace of each?

Auxiliary problem: There are four (h_1, h_2, h_3, h_4) 2-planes in \mathbb{C}^8 meeting each of K_1, K_2, K_3, K_4 . Schematically, $\square\square\square\square^4 = 4$.

Fact: All solutions H to our problem have the form $H_{i,j} = \langle h_i, h_j \rangle$ for $1 \leq i < j \leq 4$. Schematically, $\boxplus^4 = 6$.

It follows that the Galois group of $\boxplus^4 = 6$ equals the Galois group of $\square\square\square\square^4 = 4$, which is the symmetric group \mathcal{S}_4 . This action is *not* two-transitive.

This problem $\boxplus^4 = 6$ also has exceptional reality: If K_1, K_2, K_3, K_4 are real, then either two or six of the $H_{i,j}$ are real, and never four or zero.

Galois Groups of Enumerative Problems

In 1870, Jordan explained how *algebraic* Galois groups arise naturally from problems in enumerative geometry; earlier (1851), Hermite showed that such an algebraic Galois group coincides with a geometric monodromy group.

This Galois group of a geometric problem is a subtle invariant. When it is *deficient* (i.e. not the full symmetric group), the geometric problem has some exceptional, intrinsic structure.

Hermite's observation, work of Vakil, and some number theory together with modern computational tools give several methods to determine Galois groups, at least experimentally.

I will describe a project to study Galois groups for problems coming from the Schubert calculus using numerical algebraic geometry, symbolic computation, combinatorics, and more traditional methods.

Some Theory

A degree e surjective map $E \xrightarrow{\pi} B$ of equidimensional irreducible varieties (up to codimension one, $E \rightarrow B$ is a covering space of degree e)

\rightsquigarrow degree e extension of fields of rational functions $\pi^* K(B) \subset K(E)$.

Define the Galois group $\text{Gal}(E/B) \subset \mathcal{S}_e$ to be the Galois group of the Galois closure of this extension.

Hermite's Theorem. (Work over \mathbb{C} .) Restricting $E \rightarrow B$ to open subsets over which π is a covering space, $E' \rightarrow B'$, the Galois group is equal to the monodromy group of deck transformations.

This is the group of permutations of a fixed fiber induced by analytically continuing the fiber over loops in the base.

Point de départ: Such monodromy permutations are readily and reliably computed using methods of numerical algebraic geometry.

Enumerative Geometry

“Enumerative Geometry is the art of determining the number e of geometric figures x having specified positions with respect to other, fixed figures b .”
— Hermann Cäsar Hannibal Schubert, 1879.

B := configuration space of the fixed figures, and X := the space of the figures x we count. Then $E \subset X \times B$ consists of pairs (x, b) where $x \in X$ has given position with respect to $b \in B$.


The projection $E \rightarrow B$ is a degree e cover outside of some discriminant locus, and the *Galois group of the enumerative problem* is $\text{Gal}(E/B)$.

In the problem of four lines, B = four-tuples of lines, X = lines, and E consists of 5-tuples $(m, \ell_1, \ell_2, \ell_3, \ell_4)$ with m meeting each ℓ_i . We showed that this has Galois group the symmetric group \mathcal{S}_2 .

Schubert Problems

The Schubert calculus is an algorithmic method promulgated by Schubert to solve a wide class of problems in enumerative geometry.

Schubert problems are problems from enumerative geometry involving linear subspaces of a vector space incident upon other linear spaces, such as the problem of four lines, and the problems $\square\square\square^4 = 4$ and $\square\square^4 = 6$.



As there are many millions of computable Schubert problems, many with their own unique geometry, they provide a rich and convenient laboratory for studying Galois groups of geometric problems.

Proof-of-concept computation

Leykin and I used off-the-shelf numerical homotopy continuation software to compute Galois groups of some Schubert problems formulated as the intersection of a skew Schubert variety with Schubert hypersurfaces.

In every case, we found monodromy permutations generating the full symmetric group (determined by Gap). This included one Schubert problem with $e = 17,589$ solutions.

We conjectured that problems of this type will always have the full symmetric group as Galois group.

As a first step, White and I showed these Galois groups are 2-transitive as permutation groups.

The bottleneck to studying more general problems numerically is that we need numerical methods to solve *one* instance of the problem.

Numerical Project

Recent work, including certified continuation (Beltrán and Leykin), Littlewood-Richardson homotopies (Vakil, Verschelde, and S.), regeneration (Hauenstein), implementation of Pieri and of Littlewood-Richardson homotopies (Martín del Campo and Leykin) and new algorithms in the works will enable the reliable numerical computation of Galois groups of more general problems.

We plan to use a supercomputer whose day job is calculus instruction to investigate many of the millions of accessible and computable Schubert problems. Our intention is to build a library of Schubert problems (expected to be very few) whose Galois groups are deficient.

These data would be used to generate conjectures, leading to proofs about Galois groups of Schubert problems, as well as showcase the possibilities of numerical computation.

Vakil's Criteria

Vakil introduced two combinatorial criteria which can be used to show that the Galois group of a Schubert problem contains the alternating group. (Is *at least alternating*). The first criterion is simple combinatorics, while the second requires knowledge of 2-transitivity.

Theorem. (Brooks, Martín del Campo, S.) *The Galois group of any Schubert problem involving 2-planes in \mathbb{C}^n is at least alternating.*

By Vakil's second criterion, to show high-transitivity (S_e or A_e), we often only need 2-transitivity. All known Galois groups of Schubert problems are either at least alternating or fail to be 2-transitive.

White and I are studying 2-transitivity.

Theorem. *Every Schubert problem involving 3-planes in \mathbb{C}^n is 2-transitive.
Every special Schubert problem (partition a single row) is 2-transitive.*

↪ The proof suggests that not 2-transitive implies imprimitive.

Vakil's Criteria II

Vakil's geometric Littlewood-Richardson rule, his criteria, and some 2-transitivity give an algorithm that can show a Schubert problem has at least alternating monodromy.

Python code written by Brooks is being modified by Maril and Moore to implement this algorithm. There are serious computer-science challenges to overcome.

Our goal is to use it to test all Schubert problems on all small Grassmannians (many hundreds of million Schubert problems), and get a second library of Schubert problems with deficient Galois groups.

Specialization Lemma


Given $\pi: E \rightarrow B$ with B rational, the fiber $\pi^{-1}(b)$ above a \mathbb{Q} -rational point $b \in B(\mathbb{Q})$ has a minimal polynomial $p_y(t) \in \mathbb{Q}[t]$. In this situation, the algebraic Galois group of $p_y(t)$ is a subgroup of $\text{Gal}(E/B)$.

Working modulo a prime, the minimal polynomial of such fibers are easy to compute when $e \lesssim 500$. The degrees of its irreducible factors give the cycle type of a Frobenius element in the Galois group.

This quickly determines the Galois group when it is the full symmetric group, and allows the estimation of the Galois group when it is not.

Using Vakil's criteria and this method, we have determined the Galois groups of all Schubert problems involving 4-planes in \mathbb{C}^8 and \mathbb{C}^9 . (The first interesting case.) The deficient Schubert problems fall into a few easily-identified families, which suggests the possibility of classifying all deficient Schubert problems and identifying their Galois groups.

Thank You!



A Problem with Exceptional Geometry

Q: What 4-planes H in \mathbb{C}^8 meet four general 4-planes K_1, K_2, K_3, K_4 in a 2-dimensional subspace of each?

Auxiliary problem: There are four (h_1, h_2, h_3, h_4) 2-planes in \mathbb{C}^8 meeting each of K_1, K_2, K_3, K_4 . Schematically, $\square\square\square^4 = 4$.

Fact: All solutions H to our problem have the form $H_{i,j} = \langle h_i, h_j \rangle$ for $1 \leq i < j \leq 4$. Schematically, $\boxplus^4 = 6$.

It follows that the Galois group of $\boxplus^4 = 6$ equals Galois group of $\square\square\square^4 = 4$, which is the symmetric group \mathcal{S}_4 . This action is imprimitive.

This problem $\boxplus^4 = 6$ also has exceptional reality: If K_1, K_2, K_3, K_4 are real, then either two or six of the $H_{i,j}$ are real, and never four or zero.