

1. (15) Define the following:

a. a linear transformation

Let T be a mapping from the vector space V into the vector space W . T is a linear transformation if for every \vec{x} and $\vec{y} \in V$ and $\alpha \in R$ we have

$$T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$$

$$T(\alpha\vec{x}) = \alpha T(\vec{x}).$$

b. matrix representation of a linear transformation

If T is a linear transformation from V into W and B_1 and B_2 are bases of V and W respectively, then A is a matrix representation of T with respect to the two bases if

$$[T\vec{x}]_{B_2} = A[\vec{x}]_{B_1}.$$

c. row space of a matrix

Let $\vec{r}_1, \vec{r}_2, \dots, \vec{r}_m$ be the m rows of an $m \times n$ matrix. That is, each of the \vec{r}_i can be thought of as a vector in R^n . The row space of A is the span of these row vectors.

2. (20) $S = \{1, t, t^2\}$ is the standard basis of P_3 , the vector space of polynomials of degree 2 or less. Let $U = \{t^2 + 2t, 6 - t^2, 1 + t + t^2\}$.

a. Show that U is a basis of P_3 .

Since the dimension of P_3 is 3, and the set U contains 3 vectors, it will suffice to show that U is either linearly independent or spanning. To see that this set is linearly independent suppose there are constants c_i such that

$$c_1(t^2 + 2t) + c_2(6 - t^2) + c_3(1 + t + t^2) = \vec{0}$$

$$(6c_2 + c_3) + (2c_1 + c_3)t + (c_1 - c_2 + c_3)t^2 = \vec{0}.$$

This polynomial can be the zero polynomial if and only if

$$6c_2 + c_3 = 0$$

$$2c_1 + c_3 = 0$$

$$c_1 - c_2 + c_3 = 0.$$

These equations have only one solution: $c_1 = 0$, $c_2 = 0$, $c_3 = 0$. Thus, the polynomials in the set U are linearly independent.

- b. If $[\vec{x}]_U = [3, -2, 5]$, what is \vec{x} ?

$$\begin{aligned}\vec{x} &= 3(t^2 + 2t) - 2(6 - t^2) + 5(1 + t + t^2) \\ &= -7 + 11t + 10t^2.\end{aligned}$$

- c. Find the change of basis matrix P such that $[\vec{x}]_S = P[\vec{x}]_U$.

$$P = \begin{bmatrix} 0 & 6 & 1 \\ 2 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix}.$$

- d. Using the matrix P , calculate $[1 + 2t + 3t^2]_U$.

From $[\vec{x}]_S = P[\vec{x}]_U$, we have $[\vec{x}]_U = P^{-1}[\vec{x}]_S$. So

$$\begin{aligned}[1 + 2t + 3t^2]_U &= \begin{bmatrix} 0 & 6 & 1 \\ 2 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \\ &= \begin{bmatrix} -\frac{1}{8} & \frac{7}{8} & -\frac{3}{4} \\ \frac{1}{8} & \frac{1}{8} & -\frac{1}{4} \\ \frac{1}{4} & -\frac{3}{4} & \frac{3}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \\ &= \begin{bmatrix} -\frac{5}{8} \\ -\frac{3}{8} \\ \frac{13}{4} \end{bmatrix}.\end{aligned}$$

3. (15) Various mappings are defined below, determine which, if any, are linear transformations.

- a. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^1, T(x_1, x_2) = x_1$.

This is a linear transformation, as is easily checked.

- b. $T : \mathbb{R}^2 \rightarrow \mathbb{R}^1, T(x_1, x_2) = x_1x_2$.

This is not a linear transformation. As

$$T(2(x_1, x_2)) = T((2x_1, 2x_2)) = 4x_1x_2 \neq 2T((x_1, x_2)).$$

- c. $T : \mathbb{R}^2 \rightarrow M_{2,3}, T(x_1, x_2) = \begin{bmatrix} x_1 & 0 & 1 \\ 0 & x_2 & 0 \end{bmatrix}$.

This too is not a linear transformation. A quick check shows that $T(2\vec{x}) \neq 2T(\vec{x})$.

4. (25) Let T denote the linear transformation of R^3 into itself that first reflects R^3 through the plane $x_1 + x_2 - 2x_3 = 0$, and then rotates R^3 about the z -axis 60° in a counter clockwise direction when viewed from the positive z -axis towards the origin. Hint: if $T = T_2T_1$, then the matrix representation of T is A_2A_1 where A_i is the matrix representation of T_i .

a. Find the matrix representation of T with respect to the standard basis.

Let's find the matrix representation of T_1 , the transformation that reflects through the plane, with respect to the standard basis first. To do this easily we first find a matrix representation of T_1 with respect to the basis

$B_1 = \{(0, 2, 1), (2, 0, 1), (1, 1, -2)\}$ If \hat{A}_1 denotes this matrix, then

$$\hat{A}_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}.$$

If P denotes the change of basis matrix $[\vec{x}]_S = P[\vec{x}]_{B_1}$, then

$$P = \begin{bmatrix} 0 & 2 & 1 \\ 2 & 0 & 1 \\ 1 & 1 & -2 \end{bmatrix}, \text{ and the matrix we are looking for } A_1, \text{ which represents}$$

T_1 with respect to the standard basis is

$$A_1 = P\hat{A}_1P^{-1}$$

$$\begin{aligned} &= \begin{bmatrix} 0 & 2 & 1 \\ 2 & 0 & 1 \\ 1 & 1 & -2 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \begin{bmatrix} 0 & 2 & 1 \\ 2 & 0 & 1 \\ 1 & 1 & -2 \end{bmatrix}^{-1} \\ &= \begin{bmatrix} 0 & 2 & 1 \\ 2 & 0 & 1 \\ 1 & 1 & -2 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \begin{bmatrix} -\frac{1}{12} & \frac{5}{12} & \frac{1}{6} \\ \frac{5}{12} & -\frac{1}{12} & \frac{1}{6} \\ \frac{1}{6} & \frac{1}{6} & -\frac{1}{3} \end{bmatrix} \\ &= \frac{1}{3} \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & 2 \\ 2 & 2 & -1 \end{bmatrix}. \end{aligned}$$

The matrix A_2 equals

$$\begin{aligned}
 A_2 &= \begin{bmatrix} \cos 60 & -\sin 60 & 0 \\ \sin 60 & \cos 60 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}.
 \end{aligned}$$

We're now able to compute the matrix representation A of T with respect to the standard basis.

$$\begin{aligned}
 A &= A_2 A_1 \\
 &= \begin{bmatrix} \frac{1}{2} & -\frac{1}{2}\sqrt{3} & 0 \\ \frac{1}{2}\sqrt{3} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \frac{1}{3} \begin{bmatrix} 2 & -1 & 2 \\ -1 & 2 & 2 \\ 2 & 2 & -1 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1}{6}\sqrt{3} + \frac{1}{3} & -\frac{1}{3}\sqrt{3} - \frac{1}{6} & -\frac{1}{3}\sqrt{3} + \frac{1}{3} \\ \frac{1}{3}\sqrt{3} - \frac{1}{6} & -\frac{1}{6}\sqrt{3} + \frac{1}{3} & \frac{1}{3}\sqrt{3} + \frac{1}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix}.
 \end{aligned}$$

- b. Do we get the same linear transformation if the rotation is performed first?

The answer is no. If the rotation is performed first, then the matrix representation for this composition of the two transformations is $A_1 A_2$, and this product equals

$$A_1 A_2 = \begin{bmatrix} -\frac{1}{6}\sqrt{3} + \frac{1}{3} & -\frac{1}{3}\sqrt{3} - \frac{1}{6} & \frac{2}{3} \\ \frac{1}{3}\sqrt{3} - \frac{1}{6} & \frac{1}{6}\sqrt{3} + \frac{1}{3} & \frac{2}{3} \\ \frac{1}{3}\sqrt{3} + \frac{1}{3} & -\frac{1}{3}\sqrt{3} + \frac{1}{3} & -\frac{1}{3} \end{bmatrix},$$

and this does not equal the matrix $A_2 A_1$.

5. (25) Let T denote a linear transformation of $P_3 \rightarrow P_2$. Let $A = \begin{bmatrix} 2 & 1 & -1 \\ 3 & 0 & 5 \end{bmatrix}$ be the matrix representation of T with respect to the standard basis $S = \{1, t, t^2\}$ in P_3 and the basis $B = \{2t, 3 - t\}$ in P_2 .

- a. $T(6 - t + t^2)$ equals?

$$\begin{aligned} [T(6 - t + t^2)]_B &= A[6 - t + t^2]_S \\ &= \begin{bmatrix} 2 & 1 & -1 \\ 3 & 0 & 5 \end{bmatrix} \begin{bmatrix} 6 \\ -1 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} 10 \\ 23 \end{bmatrix}. \end{aligned}$$

Thus, $T(6 - t + t^2) = 10(2t) + 23(3 - t) = 69 - 3t$.

- b. Find a basis for the kernel of T .

The matrix A is row equivalent to the matrix $\begin{bmatrix} 1 & 0 & \frac{5}{3} \\ 0 & 1 & -\frac{13}{3} \end{bmatrix}$. The rank of

this matrix is 2, hence its null space has dimension 1 and thus, the kernel of T also has dimension 1. Since the vector $(-5, 13, 3)$ spans the null space of A , we know that the vector

$$-5 + 13t + 3t^2$$

will span the kernel of T .

- c. Find a basis for the image of T .

A basis for the column space of A is $\{(2, 3), (1, 0)\}$. Thus a basis for the image of T is

$$\{2(2t) + 3(3 - t), 2t\} = \{t + 9, 2t\}.$$