Name\_\_\_\_\_ MATH 221 Exam 3, Version B Spring 2024 501 P. Yasskin


Multiple Choice: (9 points each. No part credit.)
Circle you answers here and bubble on the Scantron.
Show your work, in case I give some part credit.

1-6	/54
7	/20
8	/15
9	/15
Total	/104

- 1. Compute  $\int_0^2 \int_p^{2p} pq \, dq \, dp.$ 
  - **a**. 24
  - **b**. 12
  - **c**. 8
  - **d**. 6
  - **e**. 4

- **2**. Approximate the integral  $\int_2^6 \int_1^5 (x^2 + y) \, dy \, dx$  by a Riemann sum using 4 squares and evaluating at the center of each square.
  - **a**. 80
  - **b**. 160
  - **c**.  $\frac{488}{3}$
  - **d**.  $\frac{976}{3}$
  - **e**. 320

3. Find the area of the interior of the upper half of the cardioid  $r = 1 + \cos \theta$ .



- **a**.  $A = \frac{3\pi}{2}$
- **b**.  $A = \frac{3\pi}{4}$
- **c**.  $A = \frac{2}{3\pi}$
- **d**.  $A = 2\pi$
- $\mathbf{e.}\ A=\pi$

- **4**. Find the average value of the function f(x,y) = y on the interior of the upper half of the cardioid  $r = 1 + \cos \theta$  as shown in problem 3.
  - **a**.  $f_{\text{ave}} = \frac{16}{9\pi}$
  - **b**.  $f_{\text{ave}} = \frac{9\pi}{16}$
  - **c**.  $f_{\text{ave}} = \frac{4}{9\pi}$
  - **d**.  $f_{ave} = \frac{4}{3}$
  - **e**.  $f_{ave} = \frac{3}{4}$

5. Find the volume of the apple given in spherical coordinates by  $\rho = 1 - \cos \phi$ .



- **a**.  $2\pi^2$
- b.  $\frac{4\pi}{3}$ c.  $\frac{8\pi}{3}$ d.  $\frac{4\pi}{5}$

- **6**. Find the surface area of the parametric surface parametrized by  $\vec{R}(u,v) = \langle u+v, u-v, 2u+2v \rangle$ for  $0 \le u \le 2$  and  $0 \le v \le 3$ .
  - **a**.  $12\sqrt{5}$
  - **b**.  $6\sqrt{5}$
  - **c**.  $2\sqrt{5}$
  - **d**. 6
  - **e**. 36

Work Out: (Points indicated. Part credit possible. Show all work.)

- 7. (20 points) Consider the region between the y-axis and the parabola  $x = 9 y^2$ .
  - **a**. Find the mass of the region, if the surface density is  $\delta = y^2$ .

**b**. Find the center of mass of the region, if the surface density is  $\delta = y^2$ . Write your answer as an ordered pair.

**8**. (15 points) Compute  $\iiint_P \vec{\nabla} \cdot \vec{F} dV$  for  $\vec{F} = \langle xz, yz, z^2 \rangle$  over the solid above the paraboloid  $z = x^2 + y^2$  below the plane z = 4.



**9**. (15 points) Compute  $\iint_P \vec{\nabla} \times \vec{F} \cdot d\vec{S}$  for  $\vec{F} = \langle -yz, xz, z^2 \rangle$  over the parabolic surface  $z = x^2 + y^2$  below z = 4 oriented down and out. It may be parametrized by  $\vec{R}(r,\theta) = \langle r\cos\theta, r\sin\theta, r^2 \rangle$ 

