

Fall 2004 Math 151
Exam 2B: Solutions
Mon, 01/Nov **©2004, Art Belmonte**

- (d) As $x \rightarrow \infty$, we see that $3^{-x} = \frac{1}{3^x} \rightarrow 0$ since the numerator is fixed and the denominator increases without bound.
- (c) The natural logarithm is defined for all positive real arguments; i.e., for $|x| > 0$ or $x \neq 0$.
- (d) Isolate x step by step.

$$\begin{aligned} \log_3(3x+2) &= 2 \\ 3x+2 &= 3^2 = 9 \\ 3x &= 7 \\ x &= 7/3 \end{aligned}$$

- (c) The range of f^{-1} is the domain of f . Since we are given $f(x) = \sqrt{1-x}$, we require $1-x \geq 0$ or $1 \geq x$. This is the interval $(-\infty, 1]$.
- (b) Since $f(1) = 2$ and $g = f^{-1}$, we have $g(2) = 1$ and thus

$$\begin{aligned} g'(2) &= \frac{1}{f'(g(2))} \\ &= \frac{1}{f'(1)} \\ &= \frac{1}{(5x^4 + 3x^2) \Big|_{x=1}} = \frac{1}{8}. \end{aligned}$$

- (c) Use the product rule to differentiate $f(x) = x^2 \tan x$.

$$f'(x) = 2x \tan x + x^2 \sec^2 x$$

- (b) Now $f(x) = 2\sqrt{e^x} = 2(e^x)^{1/2} = 2e^{x/2}$, whence

$$f'(x) = 2e^{x/2} \left(\frac{1}{2}\right) = e^{x/2} = (e^x)^{1/2} = \sqrt{e^x}.$$

- (d) The derivative of position is velocity. In turn the derivative of velocity is acceleration.

$$\begin{aligned} \mathbf{r}(t) &= [(\cos t)^2, t] \\ \mathbf{v}(t) = \mathbf{r}'(t) &= [2(\cos t)(-\sin t), 1] = [-\sin 2t, 1] \\ \mathbf{a}(t) = \mathbf{v}'(t) = \mathbf{r}''(t) &= [-2\cos 2t, 0] \\ &= \left[-2\cos \frac{2\pi}{3}, 0\right] = [1, 0] \end{aligned}$$

when $t = \frac{\pi}{3}$.

- (b) Repeatedly differentiate until a pattern is evident.

$$\begin{aligned} f(x) &= \cos x \\ f'(x) &= -\sin x \\ f''(x) &= -\cos x \\ f'''(x) &= \sin x \\ f^{(4)}(x) &= \cos x \end{aligned}$$

We see that the derivatives repeat in a cycle of four. Hence $f^{(28)}(x) = \cos x$. More precisely, $f^{(k)} = f^{(k \bmod 4)}$ where $k \bmod 4$ signifies remainder upon division by 4. Because $28 \bmod 4 = 0$, we conclude that

$$f^{(28)}(x) = f^{(0)}(x) = f(x) = \cos x.$$

- (d) At $x = \sqrt{\pi}$, we have

$$\begin{aligned} f(x) &= \sin(x^2) = \sin \pi = 0, \\ f'(x) &= 2x \cos(x^2) = -2\sqrt{\pi}. \end{aligned}$$

Therefore, the desired linear approximation is

$$\begin{aligned} L(x) &= f(\sqrt{\pi}) + f'(\sqrt{\pi})(x - \sqrt{\pi}) \\ &= 0 - 2\sqrt{\pi}(x - \sqrt{\pi}) \\ &= 2\pi - 2\sqrt{\pi}x. \end{aligned}$$

- (d) The distance between $(x, 0)$ and $(0, 1)$ is

$$s = \sqrt{(x-0)^2 + (0-1)^2} = (x^2 + 1)^{1/2}.$$

Differentiation with respect to time t yields


$$\begin{aligned} \frac{ds}{dt} &= \frac{1}{2}(x^2 + 1)^{-1/2} \left(2x \frac{dx}{dt}\right) \\ &= \frac{2x}{\sqrt{x^2 + 1}}, \end{aligned}$$

since $dx/dt = 2$ from the statement of the problem.

- (d) Let $w = 2^{-x}$ (to give it a name). Then

$$2^{-x} + 2^{-x} = w + w = 2w = (2^1)(2^{-x}) = 2^{1-x}.$$

- (c) Geometrically, Newton's method amounts to following tangent lines to where they intersect the x -axis. With the point $P(2, 2)$ on the graph of f and the slope of the tangent line thereat between 3 and 5, we have the following diagram. It is clear from the picture that the next Newton iterate lies in the interval $1 < x < 2$ or $(1, 2)$. No computation is required!



Alternatively, take an analytical approach. Since the graph of $y = f(x)$ passes through $(2, 2)$, we have $f(2) = 2$. Using a slope of $f'(2) = 3$, the next Newton iterate is

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = 2 - \frac{2}{3} = \frac{4}{3} \approx 1.33 \in (1, 2).$$

Similarly, using a slope of $f'(2) = 5$ gives

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)} = 2 - \frac{2}{5} = \frac{8}{5} = 1.6 \in (1, 2).$$

14. Implicitly differentiate the equation

$$(x^2 + y^2)^2 = 2(x^2 - y^2) + 6x + 7$$

with respect to x , then solve for dy/dx .

$$\begin{aligned} 2(x^2 + y^2) \left(2x + 2y \frac{dy}{dx} \right) &= 2 \left(2x - 2y \frac{dy}{dx} \right) + 6 \\ (x^2 + y^2) \left(x + y \frac{dy}{dx} \right) &= \left(x - y \frac{dy}{dx} \right) + \frac{3}{2} \\ (y + (x^2 + y^2)y) \frac{dy}{dx} &= x + \frac{3}{2} - (x^2 + y^2)x \\ \frac{dy}{dx} &= \frac{x + \frac{3}{2} - (x^2 + y^2)x}{y + (x^2 + y^2)y} \end{aligned}$$

Finally, plug in $(x, y) = (2, 1)$.

$$\frac{dy}{dx} = \frac{\frac{7}{2} - 10}{6} = \frac{-\frac{13}{2}}{6} = -\frac{13}{12}$$

15. We are given $x = te^{-t}$ and $y = (t + 1)^{1/3}$. The point $(x, y) = (0, 1)$ corresponds to $t = 0$.

(i) The parametric derivatives are

$$\begin{aligned} \frac{dx}{dt} &= (1)e^{-t} + te^{-t}(-1) = (1-t)e^{-t}, \\ \frac{dy}{dt} &= \frac{1}{3}(t+1)^{-2/3}. \end{aligned}$$

(ii) Therefore,


$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{\frac{1}{3}(t+1)^{-2/3}}{(1-t)e^{-t}}.$$

(iii) The slope m of the desired tangent line is obtained by plugging $t = 0$ into the expression for dy/dx . This yields $m = \frac{1}{3}$. Hence an equation of the tangent line is $y - 1 = \frac{1}{3}(x - 0)$ or $y = \frac{1}{3}x + 1$.

16. Given $f(x) = e^{bx} \sin ax$, we have

$$\begin{aligned} f'(x) &= (be^{bx}) \sin ax + e^{bx} (a \cos ax) = e^{bx} (b \sin ax + a \cos ax) \\ f''(x) &= (be^{bx}) (b \sin ax + a \cos ax) + e^{bx} (ab \cos ax - a^2 \sin ax) \\ &= e^{bx} ((b^2 - a^2) \sin ax + 2ab \cos ax) \end{aligned}$$

17. Let y be the height of the rocket and θ the camera's angle of elevation. Here is a diagram followed by the relevant computations involved in this related rates problem.



$$\begin{aligned} \tan \theta &= \frac{y}{4000} \\ \sec^2 \theta \cdot \frac{d\theta}{dt} &= \frac{dy/dt}{4000} \\ \frac{d\theta}{dt} &= \frac{dy/dt}{4000} (\cos \theta)^2 \\ \frac{d\theta}{dt} &= \frac{600}{4000} \left(\frac{4}{5} \right)^2 = \frac{12}{125} = 0.096 \end{aligned}$$

since for $y = 3000$, we have $\cos \theta = \frac{4000}{\sqrt{3000^2 + 4000^2}} = \frac{4}{5}$. The camera angle is changing at $\frac{12}{125} = 0.096$ rad/s or $5.5^\circ/\text{s}$.

18. The linear approximation to f at $a = 1$ is $L(x) = 3x + 2$. Thus $f(1) = L(1) = 5$ and $f'(1) = L'(1) = 3$. We are given $H = \sqrt{f}$ or $H(x) = (f(x))^{1/2}$. At $x = 1$, $H(1) = \sqrt{5}$ and $H'(x) = \frac{1}{2}(f(x))^{-1/2} f'(x) = \frac{1}{2} \left(\frac{1}{\sqrt{5}} \right) (3) = \frac{3}{2\sqrt{5}}$. Hence the linear approximation to H at $a = 1$ is

$$L_H(x) = H(1) + H'(1)(x - 1) = \sqrt{5} + \frac{3}{2\sqrt{5}}(x - 1).$$