## MATH 151, FALL 2011 COMMON EXAM II - VERSION B

Last Name:	First Name:
G:	Cardina NI.

PART I: Multiple Choice (15 questions, 4 points each. No Calculators) Write your name, section number, and version letter (B) of the exam on the ScanTron form.

1. Find the derivative of  $g(x) = \frac{x^3 + 1}{x^2 + 1}$ .

(a) 
$$g'(x) = \frac{x^4 + 3x^2 - 2x}{(x^2 + 1)^2}$$

(b) 
$$g'(x) = \frac{5x^4 + 3x^2 + 2x}{x^2 + 1}$$

(c) 
$$g'(x) = \frac{x^4 + 3x^2 + 2x}{(x^2 + 1)^2}$$

(d) 
$$g'(x) = \frac{x^4 + 3x^2 - 2x}{x^2 + 1}$$

(e) 
$$g'(x) = \frac{5x^4 + 3x^2 + 2x}{(x^2 + 1)^2}$$

- 2. A ball is thrown vertically upward with a velocity of 80 feet per second and the height, s, of the ball at time t seconds is given by  $s(t) = 80t 16t^2$ . What is the velocity of the ball when it is 96 feet above the ground on its way up?
  - (a) 112 ft/sec
  - (b) 24 ft/sec
  - (c) 16 ft/sec
  - (d) 48 ft/sec
  - (e) 64 ft/sec
- 3. Which of the following vectors is tangent to the curve  $\mathbf{r}(\mathbf{t}) = \langle \sqrt{t^2 + 1}, t \rangle$  at the point  $(2, \sqrt{3})$ ?
  - (a)  $\left\langle \frac{1}{\sqrt{5}}, 1 \right\rangle$
  - (b)  $\left\langle \frac{1}{2}, 1 \right\rangle$
  - (c)  $\left\langle \frac{\sqrt{3}}{4}, 1 \right\rangle$
  - (d)  $\left\langle \frac{2}{\sqrt{5}}, 1 \right\rangle$
  - (e)  $\left\langle \frac{\sqrt{3}}{2}, 1 \right\rangle$

- 4. Find the  $81^{st}$  derivative of  $f(x) = \frac{1}{x}$ .
  - (a)  $f^{(81)}(x) = -\frac{(81)!}{x^{81}}$
  - (b)  $f^{(81)}(x) = \frac{(80)!}{x^{80}}$
  - (c)  $f^{(81)}(x) = -\frac{(81)!}{x^{82}}$
  - (d)  $f^{(81)}(x) = -\frac{(80)!}{x^{80}}$
  - (e)  $f^{(81)}(x) = \frac{(81)!}{x^{81}}$
- 5.  $\lim_{x \to -\infty} (9 7e^{-x}) =$ 
  - (a)  $-\infty$
  - (b) 0
  - (c)  $\infty$
  - (d) 7
  - (e) 9
- 6. At what point on the graph of  $f(x) = \sqrt{x}$  is the tangent line parallel to the line 2x 3y = 4?
  - (a)  $\left(\frac{16}{9}, \frac{4}{3}\right)$
  - (b)  $\left(\frac{9}{16}, 0\right)$
  - (c)  $\left(\frac{4}{3}, \frac{2}{\sqrt{3}}\right)$
  - (d)  $\left(\frac{9}{16}, \frac{3}{4}\right)$
  - (e)  $\left(\frac{1}{16}, \frac{1}{4}\right)$
- 7. Given the equation  $2xy + \pi \sin(y) = 2\pi$ , find  $\frac{dy}{dx}$  when x = 1 and  $y = \frac{\pi}{2}$ .
  - (a)  $-\frac{\pi}{2-\pi}$

  - (b)  $-\frac{\pi}{3}$ (c)  $-\frac{\pi}{2+\pi}$
  - (d) 0
  - (e)  $-\frac{\pi}{2}$

- 8. Find the equation of the tangent line to the graph of  $f(x) = \frac{x}{1+2x}$  at x = 1.
  - (a)  $y \frac{1}{3} = -\frac{1}{9}(x-1)$
  - (b)  $y \frac{1}{3} = -\frac{4}{9}(x 1)$
  - (c)  $y \frac{1}{3} = \frac{x}{1+2x}(x-1)$
  - (d)  $y \frac{1}{3} = \frac{1}{9}(x 1)$
  - (e)  $y \frac{1}{3} = -\frac{x}{1+2x}(x-1)$
- 9. If  $f(x) = \sin(g(x))$ , find f'(2) given that  $g(2) = \frac{\pi}{3}$  and  $g'(2) = \frac{\pi}{4}$ .
  - (a)  $\frac{\pi}{8}$
  - (b)  $\frac{\sqrt{3}\pi}{8}$
  - (c)  $\frac{1}{2}$
  - (d)  $-\frac{\sqrt{3}\pi}{8}$
  - (e)  $-\frac{\pi}{8}$
- 10.  $\lim_{x \to 0} \frac{\sin^3(4x)}{x^3} =$ 
  - (a) ∞
  - (b) 64
  - (c) 1
  - (d) 0
  - (e) 4
- 11. Find the slope of the tangent line to the curve  $x = t^2 + t + 1$ ,  $y = \sqrt{t} + 4$  at t = 9.
  - (a)  $\frac{1}{114}$
  - (b)  $\frac{3}{5}$
  - (c)  $\frac{19}{6}$
  - (d)  $\frac{5}{12}$
  - (e) 114


- 12. Find the derivative of  $h(t) = (t^4 + 7t)^5$ .
  - (a)  $h'(t) = 5(4t^3 + 7)^4$
  - (b)  $h'(t) = 5(t^4 + 7t)^4(4t^3)$
  - (c)  $h'(t) = 5(t^4 + 7t)(4t^3 + 7)$
  - (d)  $h'(t) = 20t^{19} + 7^5(5t^4)$
  - (e)  $h'(t) = 5(t^4 + 7t)^4(4t^3 + 7)$
- 13. Given f(x) is a one-to-one function, find g'(3) where g is the inverse of the function  $f(x) = x^9 + x^3 + x$ .
  - (a)  $g'(3) = \frac{1}{12}$
  - (b) g'(3) = 1
  - (c)  $g'(3) = \frac{1}{13}$
  - (d)  $g'(3) = \frac{1}{9}$
  - (e) g'(3) = 13
- 14. Find the derivative of  $f(x) = x^3 e^{2x}$ .
  - (a)  $f'(x) = 3x^2e^{2x} + 2x^3e^{2x}$
  - (b)  $f'(x) = 6x^2e^{2x}$
  - (c)  $f'(x) = 3x^2e^{2x} + x^3e^{2x}$
  - (d)  $f'(x) = 3x^2e^{2x}$
  - (e)  $f'(x) = 3x^2e^{2x} + 2x^4e^{2x-1}$
- 15. Find the linear approximation, L(x), for  $f(x) = \sqrt[3]{x}$  at x = -8.
  - (a)  $L(x) = -2 + \frac{1}{12}(x+8)$
  - (b)  $L(x) = -2 \frac{1}{12}(x+8)$
  - (c)  $L(x) = -2 + \frac{1}{12}(x-8)$
  - (d)  $L(x) = -2 \frac{1}{12}(x-8)$
  - (e)  $L(x) = -2 + \frac{1}{4}(x+8)$

## PART II WORK OUT

<u>Directions</u>: Present your solutions in the space provided. *Show all your work* neatly and concisely and *Box your final answer*. You will be graded not merely on the final answer, but also on the quality and correctness of the work leading up to it.


16. (8 pts) An observer is standing 8 feet from the base of a balloon launching point. At the instant the balloon has risen vertically 6 feet, the height of the balloon is increasing at a rate of 10 feet per minute. How fast is the distance from the observer to the balloon changing at this same instant? Assume the balloon starts on the ground and rises vertically.

17. (8 pts) Find the second derivative of  $f(x) = \tan(x^3)$ .



19. (8 pts) For the equation  $y = e^{2x} + e^{-3x}$ , show y'' + y' - 6y is a constant. Find the constant.

20. (8 pts) Draw a diagram to show there are two tangent lines to the parabola  $y = 2x^2$  that pass through the point (1, -3) by sketching the parabola and both tangent lines on the grid provided below. Find the x-coordinates where these tangent lines touch the parabola.



Last Name:	First Name:	

## Section No:

Question	Points Awarded	Points
1-15		60
16		8
17		8
18		8
19		8
20		8
		100