

Name _____ Sec _____

MATH 251 Exam 2 Spring 2011
Section 511 Solutions P. Yasskin

Multiple Choice: (5 points each. No part credit.)

1. Compute $\int_0^2 \int_0^y xy \, dx \, dy$.

- a. 1
- b. 2 Correct Choice
- c. 3
- d. 4
- e. y^2

$$\int_0^2 \int_0^y xy \, dx \, dy = \int_0^2 \frac{x^2 y}{2} \Big|_{x=0}^y \, dy = \frac{1}{2} \int_0^2 y^3 \, dy = \frac{1}{2} \frac{y^4}{4} \Big|_{y=0}^2 = 2$$

2. Find the area of one loop of the rose $r = \sin(3\theta)$.

- a. $\frac{\pi}{12}$ Correct Choice
- b. $\frac{\pi}{6}$
- c. $\frac{\pi}{4}$
- d. $\frac{\pi}{3}$
- e. $\frac{\pi}{2}$

$$\sin(3\theta) = 0 \text{ at } \theta = 0 \text{ and } 3\theta = \pi \text{ or } \theta = \pi/3$$

$$\begin{aligned} A &= \int_0^{\pi/3} \int_0^{\sin(3\theta)} r \, dr \, d\theta = \int_0^{\pi/3} \left[\frac{r^2}{2} \right]_{r=0}^{\sin(3\theta)} \, d\theta = \frac{1}{2} \int_0^{\pi/3} \sin^2(3\theta) \, d\theta \\ &= \frac{1}{2} \int_0^{\pi/3} \frac{1 - \cos(6\theta)}{2} \, d\theta = \frac{1}{4} \left[\theta - \frac{\sin(6\theta)}{6} \right]_{\theta=0}^{\pi/3} = \frac{\pi}{12} \end{aligned}$$

3. Compute $\iiint x^2 + y^2 \, dV$ over the region between the cones $z = \sqrt{x^2 + y^2}$ and $z = 4 - \sqrt{x^2 + y^2}$.

- a. $\frac{8\pi}{3}$
- b. $\frac{16\pi}{3}$
- c. $\frac{32\pi}{3}$
- d. $\frac{16\pi}{5}$
- e. $\frac{32\pi}{5}$ Correct Choice

In cylindrical coordinates, the cones are $z = r$ and $z = 4 - r$ which intersect at $r = 2$.

$$\int_0^{2\pi} \int_0^2 \int_r^{4-r} r^2 r \, dz \, dr \, d\theta = 2\pi \int_0^2 \left[r^3 z \right]_{z=r}^{4-r} \, dr = 2\pi \int_0^2 r^3 (4 - 2r) \, dr = 2\pi \left[r^4 - 2 \frac{r^5}{5} \right]_{r=0}^2 = 2\pi \left(16 - \frac{64}{5} \right) = \frac{32\pi}{5}$$

1-12	/60	14	/15
13	/15	15	/15
		Total	/105

4. Find the mass of the hemisphere $x^2 + y^2 + z^2 \leq 4$ with $y \geq 0$ if the density is $\delta = y$.

- a. $\frac{\pi}{2}$
- b. π
- c. 2π
- d. 4π Correct Choice
- e. 8

$$M = \iiint \delta dV = \int_0^\pi \int_0^\pi \int_0^2 \rho \sin \varphi \sin \theta \rho^2 \sin \varphi d\rho d\theta d\varphi = \int_0^2 \rho^3 d\rho \int_0^\pi \sin \theta d\theta \int_0^\pi \sin^2 \varphi d\varphi$$

$$= \left[\frac{\rho^4}{4} \right]_0^2 \left[-\cos \theta \right]_0^\pi \left[\frac{1}{2} \left(\varphi - \frac{\sin 2\varphi}{2} \right) \right]_0^\pi = 4(2) \left(\frac{\pi}{2} \right) = 4\pi$$

5. Find the center of mass of the hemisphere $x^2 + y^2 + z^2 \leq 4$ with $y \geq 0$ if the density is $\delta = y$.

- a. $\left(0, \frac{64\pi}{15}, 0\right)$
- b. $\left(0, \frac{16}{15}, 0\right)$ Correct Choice
- c. $\left(0, \frac{\pi^2}{12}, 0\right)$
- d. $\left(0, \frac{15}{16}, 0\right)$
- e. $\left(0, \frac{12}{\pi^2}, 0\right)$

$$M_{xz} = \iiint y \delta dV = \int_0^\pi \int_0^\pi \int_0^2 \rho^2 \sin^2 \varphi \sin^2 \theta \rho^2 \sin \varphi d\rho d\theta d\varphi = \int_0^2 \rho^4 d\rho \int_0^\pi \sin^2 \theta d\theta \int_0^\pi \sin^3 \varphi d\varphi$$

$$= \left[\frac{\rho^5}{5} \right]_0^2 \left[\frac{1}{2} \left(\theta - \frac{\sin 2\theta}{2} \right) \right]_0^\pi (1 - \cos^2 \varphi) \sin \varphi d\varphi = \frac{2^5}{5} \left(\frac{\pi}{2} \right) \left[-\cos \varphi + \frac{\cos^3 \varphi}{3} \right]_0^\pi$$

$$= \frac{16\pi}{5} \left(1 - \frac{1}{3} + 1 - \frac{1}{3} \right) = \frac{64\pi}{15}$$

$$\bar{y} = \frac{M_{xz}}{M} = \frac{64\pi}{15} \frac{1}{4\pi} = \frac{16}{15} \quad \bar{x} = \bar{z} = 0 \text{ by symmetry.}$$

6. Compute $\oint \vec{F} \cdot d\vec{s}$ for $\vec{F} = (-16x^2y, 9xy^2)$ counterclockwise around the ellipse $\frac{x^2}{9} + \frac{y^2}{16} = 1$.

HINTS: The ellipse may be parametrized by $\vec{r}(\theta) = (3 \cos \theta, 4 \sin \theta)$.

Since $\sin(2\theta) = 2 \sin \theta \cos \theta$, we have $4 \sin^2 \theta \cos^2 \theta = \sin^2(2\theta)$.

- a. -864π
- b. -288π
- c. 144π
- d. 288π
- e. 864π Correct Choice

$$\vec{F}(\vec{r}(\theta)) = (-16 \cdot 9 \cos^2 \theta \cdot 4 \sin \theta, 9 \cdot 3 \cos \theta \cdot 16 \sin^2 \theta) \quad \vec{v} = (-3 \sin \theta, 4 \cos \theta)$$

$$\vec{F} \cdot \vec{v} = 64 \cdot 27 \cos^2 \theta \sin^2 \theta + 27 \cdot 64 \cos^2 \theta \sin^2 \theta = 27 \cdot 64 \cdot 2 \cos^2 \theta \sin^2 \theta = 27 \cdot 32 \sin^2(2\theta)$$

$$= 27 \cdot 32 \frac{1 - \cos(4\theta)}{2} = 27 \cdot 16(1 - \cos(4\theta))$$

$$\oint \vec{F} \cdot d\vec{s} = \oint \vec{F} \cdot \vec{v} d\theta = \int_0^{2\pi} 27 \cdot 16(1 - \cos(4\theta)) d\theta = 27 \cdot 16 \left[\theta - \frac{\sin 4\theta}{4} \right]_0^{2\pi} = 864\pi$$

7. The point $\left(\frac{\pi}{3}, \frac{\pi}{6}\right)$ is a critical point of the function $f(x,y) = \sin(x)\cos(y) - \frac{\sqrt{3}}{4}x + \frac{\sqrt{3}}{4}y$

Use the Second Derivative Test to classify this critical point.

- a. Local Maximum Correct Choice
- b. Local Minimum
- c. Inflection Point
- d. Saddle Point
- e. Test Fails

$$f_x = \cos(x)\cos(y) - \frac{\sqrt{3}}{4}$$

$$f_x\left(\frac{\pi}{3}, \frac{\pi}{6}\right) = \cos\left(\frac{\pi}{3}\right)\cos\left(\frac{\pi}{6}\right) - \frac{\sqrt{3}}{4} = 0$$

$$f_y = -\sin(x)\sin(y) + \frac{\sqrt{3}}{4}$$

$$f_y\left(\frac{\pi}{3}, \frac{\pi}{6}\right) = -\sin\left(\frac{\pi}{3}\right)\sin\left(\frac{\pi}{6}\right) + \frac{\sqrt{3}}{4} = 0$$

$$f_{xx} = -\sin(x)\cos(y)$$

$$f_{xx}\left(\frac{\pi}{3}, \frac{\pi}{6}\right) = -\sin\left(\frac{\pi}{3}\right)\cos\left(\frac{\pi}{6}\right) = -\frac{3}{4} < 0$$

$$f_{yy} = -\sin(x)\cos(y)$$

$$f_{yy}\left(\frac{\pi}{3}, \frac{\pi}{6}\right) = -\sin\left(\frac{\pi}{3}\right)\cos\left(\frac{\pi}{6}\right) = -\frac{3}{4}$$






$$f_{xy} = -\cos(x)\sin(y)$$

$$f_{xy}\left(\frac{\pi}{3}, \frac{\pi}{6}\right) = -\cos\left(\frac{\pi}{3}\right)\sin\left(\frac{\pi}{6}\right) = -\frac{1}{4}$$

$$D = f_{xx}f_{yy} - f_{xy}^2$$

$$D\left(\frac{\pi}{3}, \frac{\pi}{6}\right) = \left(-\frac{3}{4}\right)\left(-\frac{3}{4}\right) - \left(-\frac{1}{4}\right)^2 = \frac{1}{2} > 0 \quad \text{Local Maximum}$$

8. Which of the following is the plot of the vector field $\vec{F}(x,y) = \frac{1}{\sqrt{x^2+y^2}}(|x|, |y|)$?



(d) Correct Choice

All arrows must be up and right. So (c) or (d). On the x -axis, $y = 0$ and so the arrows are horizontal there. So (d).

9. Find the area of the parametric surface $\vec{R}(u, t) = (ue^t, ue^{-t}, \sqrt{2}u)$ for $0 \leq u \leq 2$ and $0 \leq t \leq 1$.

HINT: Look for a perfect square.

a. $2\sqrt{2}\left(e + \frac{1}{e} - 2\right)$

b. $2\sqrt{2}\left(e + \frac{1}{e}\right)$

c. $2\sqrt{2}\left(e - \frac{1}{e} - 2\right)$

d. $2\sqrt{2}\left(e - \frac{1}{e}\right)$ Correct Choice

e. $2\left(e + \frac{1}{e}\right)$

$$\vec{e}_u = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ (e^t) & e^{-t} & \sqrt{2} \\ (ue^t) & -ue^{-t} & 0 \end{vmatrix} \quad \vec{N} = \vec{e}_u \times \vec{e}_t = \hat{i}(\sqrt{2}ue^{-t}) - \hat{j}(-\sqrt{2}ue^t) + \hat{k}(-u - u) = (\sqrt{2}ue^{-t}, \sqrt{2}ue^t, -2u)$$

$$|\vec{N}| = \sqrt{2u^2e^{-2t} + 2u^2e^{2t} + 4u^2} = u\sqrt{2}\sqrt{e^{-2t} + 2 + e^{2t}} = u\sqrt{2}(e^{-t} + e^t)$$

$$A = \iint dS = \iint |\vec{N}| du dt = \int_0^1 \int_0^2 u\sqrt{2}(e^{-t} + e^t) du dt = \sqrt{2} \left[\frac{u^2}{2} \right]_0^2 \left[-e^{-t} + e^t \right]_0^1 = 2\sqrt{2}\left(e - \frac{1}{e}\right)$$

10. Find the equation of the plane tangent to the parametric surface $\vec{R}(u, t) = (ue^t, ue^{-t}, \sqrt{2}u)$ at the point $P = \vec{R}(2, 0)$ where $u = 2$ and $t = 0$.

Hint: Evaluate the normal \vec{N} at $u = 2$ and $t = 0$.

a. $x + y - \sqrt{2}z = -4\sqrt{2}$

b. $x + y - \sqrt{2}z = 0$ Correct Choice

c. $x + y - \sqrt{2}z = 16\sqrt{2}$

d. $\sqrt{2}x - \sqrt{2}y + 2z = -4\sqrt{2}$

e. $\sqrt{2}x - \sqrt{2}y + 2z = 0$

$$P = \vec{R}(2, 0) = (2, 2, 2\sqrt{2}) \quad \vec{N} = (\sqrt{2}ue^{-t}, \sqrt{2}ue^t, -2u) = (2\sqrt{2}, 2\sqrt{2}, -4)$$

$$\vec{N} \cdot X = \vec{N} \cdot P \quad 2\sqrt{2}x + 2\sqrt{2}y - 4z = 2\sqrt{2}(2) + 2\sqrt{2}(2) - 4(2\sqrt{2}) = 0$$

11. If $\vec{F} = (xy \tan z, yz \cos x, xz \sin y)$, then $\vec{\nabla} \cdot \vec{\nabla} \times \vec{F} =$

- a. $-2z \cos y - 2x(\tan^2 z + 1)$
- b. $-2z \cos y + 2x(\tan^2 z + 1)$
- c. $2z \cos y - 2x(\tan^2 z + 1)$
- d. $2z \cos y + 2x(\tan^2 z + 1)$
- e. 0

Correct Choice

$\vec{\nabla} \cdot \vec{\nabla} \times \vec{F} = 0$ for any twice differentiable vector field.

12. Let f be the scalar potential for $\vec{F} = (2xz - 3y, 8yz - 3x, x^2 + 4y^2 + 2z)$ for which $f(0,0,0) = 0$. Then $f(1,1,1) =$

- a. 1
- b. 2
- c. 3 Correct Choice
- d. 4
- e. 5

$$\vec{\nabla}f = \vec{F} \quad \text{or} \quad (1) \partial_x f = 2xz - 3y \quad (2) \partial_y f = 8yz - 3x \quad (3) \partial_z f = x^2 + 4y^2 + 2z$$

$$(1) \Rightarrow f = x^2z - 3xy + g(y, z) \Rightarrow (4) \partial_y f = -3x + \partial_y g$$

$$(2) \text{ and } (4) \Rightarrow \partial_y g = 8yz \Rightarrow g = 4y^2z + h(z) \Rightarrow f = x^2z - 3xy + 4y^2z + h(z)$$

$$\Rightarrow (5) \partial_z f = x^2 + 4y^2 + \frac{dh(z)}{dz}$$

$$(3) \text{ and } (5) \Rightarrow \frac{dh(z)}{dz} = 2z \Rightarrow h = z^2 + C \Rightarrow f = x^2z - 3xy + 4y^2z + z^2 + C$$

To have $f(0,0,0) = 0 \Rightarrow C = 0$. So $f(1,1,1) = 1 - 3 + 4 + 1 = 3$.

Work Out: (Points indicated. Part credit possible. Show all work.)

13. (15 points) The plane $x + 2y + 4z = 8$ intersects the 1st octant ($x > 0, y > 0, z > 0$) in a triangle. Find the point on this triangle at which the function $f = xy^2z^3$ is a maximum.

Method of Eliminating a Variable:

$$x = 8 - 2y - 4z \quad f = (8 - 2y - 4z)y^2z^3 = 8y^2z^3 - 2y^3z^3 - 4y^2z^4$$

$$f_y = 16yz^3 - 6y^2z^3 - 8yz^4 = 0 \quad \Rightarrow \quad 2yz^3(8 - 3y - 4z) = 0$$

$$f_z = 24y^2z^2 - 6y^3z^2 - 16y^2z^3 = 0 \quad \Rightarrow \quad 2y^2z^2(12 - 3y - 8z) = 0$$

Since $f = xy^2z^3$ is positive in the 1st octant, we know the solution cannot have $y = 0$ or $z = 0$.

So we solve $8 - 3y - 4z = 0$ and $12 - 3y - 8z = 0$.

Solving each for $3y$ and equating gives $3y = 8 - 4z = 12 - 8z$, or $4z = 4$. So $z = 1$.

$$\text{Then } 3y = 4. \text{ So } y = \frac{4}{3}, \text{ and } x = 8 - 2\left(\frac{4}{3}\right) - 4(1) = \frac{4}{3}.$$

The point is $\left(\frac{4}{3}, \frac{4}{3}, 1\right)$.

Method of Lagrange Multipliers:

$$f = xy^2z^3 \quad \vec{\nabla}f = (y^2z^3, 2xyz^3, 3xy^2z^2) \quad g = x + 2y + 4z \quad \vec{\nabla}g = (1, 2, 4)$$

$$\text{Lagrange equations: } y^2z^3 = \lambda, \quad 2xyz^3 = 2\lambda, \quad 3xy^2z^2 = 4\lambda$$

$$\text{Substitute the first eq into the other two: } 2xyz^3 = 2y^2z^3, \quad 3xy^2z^2 = 4y^2z^3.$$

Since $f = xy^2z^3$ is positive in the 1st octant, we know the solution cannot have $x = 0$ or $y = 0$ or $z = 0$. So we cancel: $x = y$, $3x = 4z$, So $y = x$ and $z = \frac{3}{4}x$.

$$\text{Substitute into the constraint: } x + 2y + 4z = 8 \quad x + 2x + 3x = 8 \quad 6x = 8 \quad x = \frac{4}{3} \quad y = \frac{4}{3} \quad z = 1.$$

The point is $\left(\frac{4}{3}, \frac{4}{3}, 1\right)$.

14. (15 points) Compute $\iiint \vec{\nabla} \cdot \vec{F} dV$ for $\vec{F} = (xy^2, yz^2, zx^2)$ over the solid above the cone $z = \sqrt{x^2 + y^2}$ below the sphere $x^2 + y^2 + z^2 = 4$.

$$\vec{\nabla} \cdot \vec{F} = y^2 + z^2 + x^2 = \rho^2 \text{ in spherical coordinates and } dV = \rho^2 \sin \varphi d\rho d\theta d\varphi$$

$$\text{The cone is } \rho \cos \varphi = \sqrt{\rho^2 \sin^2 \varphi \cos^2 \theta + \rho^2 \sin^2 \varphi \sin^2 \theta} = \rho \sin \varphi. \text{ So } \tan \varphi = 1 \text{ or } \varphi = \pi/4.$$

$$\iiint \vec{\nabla} \cdot \vec{F} dV = \int_0^{\pi/4} \int_0^{2\pi} \int_0^2 \rho^2 \rho^2 \sin \varphi d\rho d\theta d\varphi = \left[\frac{\rho^5}{5} \right]_0^2 (2\pi) \left[-\cos \varphi \right]_0^{\pi/4} = \frac{64\pi}{5} \left(1 - \frac{1}{\sqrt{2}} \right)$$

15. (15 points) Compute $\iint_E \vec{\nabla} \times \vec{F} \cdot \hat{k} dx dy$ for $\vec{F} = (-16x^2y, 9xy^2, 0)$ over the interior of the ellipse $\frac{x^2}{9} + \frac{y^2}{16} = 1$.

HINTS: First compute $\vec{\nabla} \times \vec{F} \cdot \hat{k}$ in rectangular coordinates.

Then compute the integral in elliptic coordinates $x = 3u \cos \theta$, $y = 4u \sin \theta$.

$$\vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \partial_x & \partial_y & \partial_z \\ -16x^2y & 9xy^2 & 0 \end{vmatrix} = \hat{i}(0) - \hat{j}(0) + \hat{k}(9y^2 - -16x^2)$$

$$\vec{\nabla} \times \vec{F} \cdot \hat{k} = 9y^2 + 16x^2 = 9(4u \sin \theta)^2 + 16(3u \cos \theta)^2 = 9 \cdot 16u^2 = 144u^2$$

$$\frac{\partial(x, y)}{\partial(u, \theta)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial y}{\partial u} \\ \frac{\partial x}{\partial \theta} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} 3 \cos \theta & 4 \sin \theta \\ -3u \sin \theta & 4u \cos \theta \end{vmatrix} = 12u \quad dx dy = J du d\theta = 12u du d\theta$$

$$\iint_E \vec{\nabla} \times \vec{F} \cdot \hat{k} dx dy = \int_0^{2\pi} \int_0^1 144u^2 12u du d\theta = 2\pi 144 \cdot 12 \left[\frac{u^4}{4} \right]_0^1 = 864\pi$$