

Name _____ ID _____

MATH 251 Exam 2 Version A Spring 2013
 Sections 506 Solutions P. Yasskin

1-13	/52
14	/12
15	/28
16	/12
Total	/104

Multiple Choice: (4 points each. No part credit.)

1. Compute $\iint_R xy \, dA$ over the region R between the parabola $y = x^2$ and the line $y = 2x$.

- a. $\frac{4}{3}$
- b. $\frac{8}{3}$ Correct Choice
- c. $\frac{32}{15}$
- d. $\frac{10}{3}$
- e. $\frac{29}{6}$

SOLUTION: $x^2 = 2x$ $x = 0, 2$

$$\begin{aligned} \iint_R xy \, dA &= \int_0^2 \int_{x^2}^{2x} xy \, dy \, dx = \int_0^2 x \left[\frac{y^2}{2} \right]_{x^2}^{2x} dx = \frac{1}{2} \int_0^2 x(4x^2 - x^4) dx = \frac{1}{2} \int_0^2 (4x^3 - x^5) dx \\ &= \frac{1}{2} \left[x^4 - \frac{x^6}{6} \right]_0^2 = \frac{1}{2} \left(16 - \frac{32}{3} \right) = \frac{8}{3} \end{aligned}$$

2. Compute $\int_0^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} x \, dy \, dx$ by converting to polar coordinates.

- a. 3
- b. 9
- c. 18 Correct Choice
- d. 27
- e. 36


SOLUTION: The region is a semicircle in the right half plane.

$$\int_0^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} x \, dy \, dx = \int_{-\pi/2}^{\pi/2} \int_0^3 r \cos \theta \, r \, dr \, d\theta = \left[\sin \theta \right]_{-\pi/2}^{\pi/2} \left[\frac{r^3}{3} \right]_0^3 = (1 - -1)(9 - 0) = 18$$

3. Find the mass of the triangular plate with vertices $(0,0)$, $(2,-6)$ and $(2,6)$ if the surface density is $\rho = x^2$.
- 4
 - 6
 - 12
 - 16
 - 24 **Correct Choice**

SOLUTION: Edges: $y = -3x$ and $y = 3x$

$$M = \iint \rho dA = \int_0^2 \int_{-3x}^{3x} x^2 dy dx = \int_0^2 [x^2 y]_{-3x}^{3x} dx = \int_0^2 6x^3 dx = \left[\frac{3}{2} x^4 \right]_0^2 = 24$$



4. Find the center of mass of the triangular plate with vertices $(0,0)$, $(2,-6)$ and $(2,6)$ if the surface density is $\rho = x^2$.
- $\left(\frac{8}{5}, 0\right)$ **Correct Choice**
 - $\left(\frac{9}{5}, 0\right)$
 - $\left(\frac{5}{8}, 0\right)$
 - $\left(\frac{5}{9}, 0\right)$
 - $\left(\frac{6}{5}, 0\right)$


SOLUTION: $\bar{y} = 0$ by symmetry.

$$M_y = \iint x \rho dA = \int_0^2 \int_{-3x}^{3x} x^3 dy dx = \int_0^2 [x^3 y]_{-3x}^{3x} dx = \int_0^2 6x^4 dx = \left[\frac{6x^5}{5} \right]_0^2 = \frac{192}{5}$$

$$\bar{x} = \frac{M_y}{M} = \frac{192}{5 \cdot 24} = \frac{8}{5}$$

5. Which of the following integrals is NOT equivalent to $\int_0^4 \int_0^{4-z} \int_0^{\sqrt{y}} f(x,y,z) dx dy dz$? The region is shown.

- $\int_0^4 \int_0^{\sqrt{4-z}} \int_{x^2}^{4-z} f(x,y,z) dy dx dz$
- $\int_0^4 \int_0^{4-y} \int_{y^2}^2 f(x,y,z) dx dz dy$ **Correct Choice**
- $\int_0^4 \int_0^{\sqrt{y}} \int_0^{4-y} f(x,y,z) dz dx dy$
- $\int_0^2 \int_0^{4-x^2} \int_{x^2}^{4-z} f(x,y,z) dy dz dx$
- $\int_0^2 \int_{x^2}^4 \int_0^{4-y} f(x,y,z) dz dy dx$



SOLUTION: (b) should be $\int_0^4 \int_0^{4-y} \int_0^{\sqrt{y}} f(x,y,z) dx dz dy$

6. Find the area of one petal of the 8 petaled daisy $r = \sin(4\theta)$.

- a. $\frac{\pi}{2}$
- b. $\frac{\pi}{4}$
- c. $\frac{\pi}{8}$
- d. $\frac{\pi}{16}$ Correct Choice
- e. $\frac{\pi}{32}$

SOLUTION: $r = 0$ when $4\theta = 0, \pi$ or $\theta = 0, \pi/4$

$$A = \iint 1 dA = \int_0^{\pi/4} \int_0^{\sin(4\theta)} r dr d\theta = \int_0^{\pi/4} \left[\frac{r^2}{2} \right]_0^{\sin(4\theta)} d\theta = \frac{1}{2} \int_0^{\pi/4} \sin^2(4\theta) d\theta = \frac{1}{2} \int_0^{\pi/4} \frac{1 - \cos(8\theta)}{2} d\theta$$

$$= \frac{1}{4} \left[\theta - \frac{\sin(8\theta)}{8} \right]_0^{\pi/4} = \frac{\pi}{16}$$

7. Find the mass of the solid between the paraboloids $z = x^2 + y^2$ and $z = 8 - x^2 - y^2$ if the volume density is $\rho = z$.

- a. 64π Correct Choice
- b. 32π
- c. 16π
- d. 8π
- e. 4π

SOLUTION: $z = r^2 = 8 - r^2$ $2r^2 = 8$ $r = 2$

$$M = \iiint \rho dV = \int_0^{2\pi} \int_0^2 \int_{r^2}^{8-r^2} z r dz dr d\theta = 2\pi \int_0^2 \left[r \frac{z^2}{2} \right]_{z=r^2}^{8-r^2} dr = \pi \int_0^2 r [(8 - r^2)^2 - r^4] dr$$

$$= \pi \int_0^2 64r - 16r^3 dr = \pi [32r^2 - 4r^4]_0^2 = 64\pi$$

8. Compute $\iiint (x^2 + y^2) z dV$ over the solid hemisphere $0 \leq \sqrt{x^2 + y^2 + z^2} \leq 2$.

- a. 0
- b. $\frac{8}{3}\pi$
- c. $\frac{16}{3}\pi$ Correct Choice
- d. $\frac{64}{9}\pi$
- e. 2π

SOLUTION: $(x^2 + y^2)z = ((\rho \sin \phi \cos \theta)^2 + (\rho \sin \phi \sin \theta)^2) \rho \cos \phi = \rho^3 \sin^2 \phi \cos \phi$ $dV = \rho^2 \sin \phi$

$$\iiint (x^2 + y^2) z dV = \int_0^{2\pi} \int_0^{\pi/2} \int_0^2 \rho^3 \sin^2 \phi \cos \phi \rho^2 \sin \phi d\rho d\phi d\theta = 2\pi \int_0^{\pi/2} \sin^3 \phi \cos \phi d\phi \int_0^2 \rho^5 d\rho$$

$$= 2\pi \left[\frac{\sin^4 \phi}{4} \right]_0^{\pi/2} \left[\frac{\rho^6}{6} \right]_0^2 = 2\pi \frac{1}{4} \frac{32}{3} = \frac{16}{3}\pi$$

9. Which integral gives the arclength of the ellipse $\vec{r}(\theta) = (6 \cos \theta, 3 \sin \theta, 3 \sin \theta)$?

a. $\int_0^{2\pi} 3\sqrt{2 + 2 \sin^2 \theta} \, d\theta$ Correct Choice

b. $\int_0^{2\pi} 3\sqrt{4 + 2 \cos^2 \theta} \, d\theta$

c. $\int_0^{2\pi} 3\sqrt{2 + 2 \cos^2 \theta} \, d\theta$

d. $\int_0^{2\pi} \sqrt{2} (3 + 3 \sin^2 \theta) \, d\theta$

e. $\int_0^{2\pi} \sqrt{54} \, d\theta$

SOLUTION: $\vec{v} = (-6 \sin \theta, 3 \cos \theta, 3 \cos \theta)$ $|\vec{v}| = \sqrt{36 \sin^2 \theta + 9 \cos^2 \theta + 9 \cos^2 \theta}$

$$L = \oint ds = \int_0^{2\pi} |\vec{v}| \, d\theta = \int_0^{2\pi} 3\sqrt{4 \sin^2 \theta + 2 \cos^2 \theta} \, d\theta = \int_0^{2\pi} 3\sqrt{2 + 2 \sin^2 \theta} \, d\theta$$

10. A helical thermocouple whose shape is the curve $\vec{r}(t) = (3 \cos t, 3 \sin t, 4t)$ for $0 \leq t \leq 4\pi$ is placed in a pot of water where the temperature is $T = (41 + x^2 + y^2 + z)$ °C. Find the average temperature of the water as measured by the thermocouple.

HINT: $f_{\text{ave}} = \frac{\int f \, ds}{\int ds}$

a. $\frac{173}{4} + 8\pi$

b. $50 + 16\pi$

c. $1000\pi + 160\pi^2$

d. $250 + 40\pi$

e. $50 + 8\pi$ Correct Choice

SOLUTION: $\vec{v} = (-3 \sin t, 3 \cos t, 4)$ $|\vec{v}| = \sqrt{9 \sin^2 t + 9 \cos^2 t + 16} = 5$

$$\int ds = \int_0^{4\pi} 5 \, dt = 20\pi \quad T = 41 + x^2 + y^2 + z = 41 + 9 + 4t = 50 + 4t$$

$$\int T \, ds = \int_0^{4\pi} (50 + 4t)5 \, dt = 5[50t + 2t^2]_0^{4\pi} = 1000\pi + 160\pi^2$$

$$T_{\text{ave}} = \frac{1000\pi + 160\pi^2}{20\pi} = 50 + 8\pi$$

11. Find a scalar potential, $f(x, y, z)$, for $\vec{F}(x, y, z) = (2xy^2 + 2x + 2xz, 2x^2y - 3z, x^2 + 3z^2 - 3y)$. Then compute $f(2, 2, 2) - f(1, 1, 1)$.

- a. 0
- b. 1
- c. 7
- d. 23 Correct Choice
- e. 25

SOLUTION:

$$\partial_x f = 2xy^2 + 2x + 2xz \Rightarrow f = x^2y^2 + x^2 + x^2z + g(y, z)$$

$$\partial_y f = 2x^2y - 3z \Rightarrow f = x^2y^2 - 3yz + h(x, z)$$

$$\partial_z f = x^2 + 3z^2 - 3y \Rightarrow f = x^2z + z^3 - 3yz + k(x, y)$$

$$f(x, y, z) = x^2y^2 + x^2 + x^2z - 3yz + z^3 + C$$

$$f(2, 2, 2) - f(1, 1, 1) = (16 + 4 + 8 - 12 + 8) - (1 + 1 + 1 - 3 + 1) = 23$$

12. If $f = x^2 + y^2 - 2z^2$ and $\vec{F} = (xz, yz, -z^2)$, which of the following is false?

- a. $\vec{\nabla} \times \vec{\nabla} f = \vec{0}$
- b. $\vec{\nabla} \cdot \vec{\nabla} f = 0$
- c. $\vec{\nabla} \cdot \vec{\nabla} \times \vec{F} = 0$
- d. $\vec{\nabla}(\vec{\nabla} \cdot \vec{F}) = \vec{0}$
- e. None of the above. They are all true. Correct Choice

SOLUTION: $\vec{\nabla} \times \vec{\nabla} f = \vec{0}$ and $\vec{\nabla} \cdot \vec{\nabla} \times \vec{F} = 0$ are always true.

$$\vec{\nabla} f = (2x, 2y, -4z) \quad \vec{\nabla} \cdot \vec{\nabla} f = 2 + 2 - 4 = 0$$

$$\vec{\nabla} \cdot \vec{F} = z + z - 2z = 0 \quad \vec{\nabla}(\vec{\nabla} \cdot \vec{F}) = \vec{\nabla}(0) = \vec{0}$$

13. Compute $\iiint_C \vec{\nabla} \cdot \vec{G} dV$ for $\vec{G} = (xz, yz, z^2)$ over the solid cylinder $x^2 + y^2 \leq 25$ with $0 \leq z \leq 4$.

- a. 800π Correct Choice
- b. 400π
- c. 200π
- d. 80π
- e. 40π

SOLUTION: $\vec{\nabla} \cdot \vec{G} = z + z + 2z = 4z$

$$\iiint_C \vec{\nabla} \cdot \vec{G} dV = \int_0^4 \int_0^{2\pi} \int_0^5 4zr dr d\theta dz = 2\pi [z^2]_0^4 [r^2]_0^5 = 2\pi 16 \cdot 25 = 800\pi$$

Work Out: (Points indicated. Part credit possible. Show all work.)


14. (12 points) Compute $\iint_D x^2 dA$ over the "diamond"

shaped region bounded by the curves

$$y = 1 + x^3, \quad y = 3 + x^3, \quad y = 4 - x^3, \quad y = 7 - x^3.$$

HINT: Define curvilinear coordinates (u, v) so that

$$y = u + x^3 \quad \text{and} \quad y = v - x^3.$$



a. (2 pts) What are the boundaries in terms of u and v ?

SOLUTION: $u = 1, \quad u = 3, \quad v = 4, \quad v = 7$

b. (3 pts) Find formulas for x and y in terms of u and v .

SOLUTION: Add the formulas: $2y = u + v \quad y = \frac{u+v}{2}$

Subtract the formulas: $0 = u - v + 2x^3 \quad x = \left(\frac{v-u}{2}\right)^{1/3}$

c. (4 pts) Find the Jacobian factor $J = \left| \frac{\partial(x,y)}{\partial(u,v)} \right|$.

SOLUTION:
$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial y}{\partial u} \\ \frac{\partial x}{\partial v} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{1}{3} \left(\frac{v-u}{2}\right)^{-2/3} \left(-\frac{1}{2}\right) & \frac{1}{2} \\ \frac{1}{3} \left(\frac{v-u}{2}\right)^{-2/3} \left(\frac{1}{2}\right) & \frac{1}{2} \end{vmatrix}$$

$$= -\frac{1}{12} \left(\frac{v-u}{2}\right)^{-2/3} - \frac{1}{12} \left(\frac{v-u}{2}\right)^{-2/3} = -\frac{1}{6} \left(\frac{v-u}{2}\right)^{-2/3}$$

$$J = \left| \frac{\partial(x,y)}{\partial(u,v)} \right| = \frac{1}{6} \left(\frac{v-u}{2}\right)^{-2/3}$$

d. (1 pts) Express the integrand in terms of u and v .

SOLUTION: $x^2 = \left(\frac{v-u}{2}\right)^{2/3}$

e. (2 pts) Compute the integral.

SOLUTION:

$$\iint_D x^2 dA = \iint_D x^2 J du dv = \int_4^7 \int_1^3 \left(\frac{v-u}{2}\right)^{2/3} \frac{1}{6} \left(\frac{v-u}{2}\right)^{-2/3} du dv = \frac{1}{6} (2)(3) = 1$$

15. (28 points) Consider the elliptical region, E , in the plane $z = 2 + x + y$ above the circle $x^2 + y^2 \leq 4$ oriented upwards.

HINT: This ellipse may be parametrized by $\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, 2 + r \cos \theta + r \sin \theta)$.

- a. (10 pts) Find the normal vector \vec{N} to the ellipse and its length $|\vec{N}|$.

Note: \vec{N} starts hard but simplifies!

SOLUTION:

$$\vec{e}_r = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \cos \theta & \sin \theta & \cos \theta + \sin \theta \\ -r \sin \theta & r \cos \theta & -r \sin \theta + r \cos \theta \end{vmatrix} \quad \vec{N} = \begin{matrix} \hat{i}[\sin \theta(-r \sin \theta + r \cos \theta) - r \cos \theta(\cos \theta + \sin \theta)] \\ -\hat{j}[\cos \theta(-r \sin \theta + r \cos \theta) + r \sin \theta(\cos \theta + \sin \theta)] \\ +\hat{k}[r \cos^2 \theta + r \sin^2 \theta] \end{matrix}$$

$$\vec{N} = (-r, -r, r) \quad \text{Oriented correctly because } N_3 = r \geq 0. \quad |\vec{N}| = \sqrt{r^2 + r^2 + r^2} = \sqrt{3} r$$

- b. (3 pts) Find the surface area of the ellipse.

SOLUTION: $0 \leq r \leq 2 \quad 0 \leq \theta \leq 2\pi$

$$A = \iint_E dS = \iint_E |\vec{N}| dr d\theta = \int_0^{2\pi} \int_0^2 \sqrt{3} r dr d\theta = 2\pi \sqrt{3} \left[\frac{r^2}{2} \right]_0^2 = 4\pi \sqrt{3}$$

- c. (3 pts) Find the mass of the ellipse if the surface density is $\rho = x^2 + y^2$.

SOLUTION: $\rho = x^2 + y^2 = r^2$

$$M = \iint_E \rho dS = \int_0^{2\pi} \int_0^2 \sqrt{3} r^3 dr d\theta = 2\pi \sqrt{3} \left[\frac{r^4}{4} \right]_0^2 = 8\pi \sqrt{3}$$

- d. (12 pts) If $\vec{F} = (-yz, xz, z^2)$, compute the surface integral $\iint_E \vec{\nabla} \times \vec{F} \cdot d\vec{S}$

$$\text{SOLUTION: } \vec{\nabla} \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \partial_x & \partial_y & \partial_z \\ -yz & xz & z^2 \end{vmatrix} = \hat{i}(0 - x) - \hat{j}(0 - -y) + \hat{k}(z - -z) = (-x, -y, 2z)$$

$$(\vec{\nabla} \times \vec{F})(\vec{R}(r, \theta)) = (-r \cos \theta, -r \sin \theta, 4 + 2r \cos \theta + 2r \sin \theta) \quad \text{Recall: } \vec{N} = (-r, -r, r)$$

$$\vec{\nabla} \times \vec{F} \cdot \vec{N} = r^2 \cos \theta + r^2 \sin \theta + 4r + 2r^2 \cos \theta + 2r^2 \sin \theta = 4r + 3r^2 \cos \theta + 3r^2 \sin \theta$$

$$\iint_E \vec{\nabla} \times \vec{F} \cdot d\vec{S} = \int_0^{2\pi} \int_0^2 \vec{\nabla} \times \vec{F} \cdot \vec{N} dr d\theta = \int_0^2 \int_0^{2\pi} (4r + 3r^2 \cos \theta + 3r^2 \sin \theta) d\theta dr$$

$$= \int_0^2 [4r\theta + 3r^2 \sin \theta - 3r^2 \cos \theta]_{\theta=0}^{2\pi} dr = \int_0^2 4r(2\pi) dr = [4\pi r^2]_0^2 = 16\pi$$

16. (12 points) Compute $\iint_C \vec{G} \cdot d\vec{S}$ for $\vec{G} = (xz, yz, z^2)$ over the cylinder $x^2 + y^2 = 25$ for $0 \leq z \leq 4$ with outward normal.

SOLUTION: $\vec{R}(\theta, z) = (5 \cos \theta, 5 \sin \theta, z)$

$$\begin{array}{l} \vec{e}_\theta = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -5 \sin \theta & 5 \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} \\ \vec{e}_z = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 0 & 1 \end{vmatrix} \end{array} \quad \begin{array}{l} \vec{N} = \hat{i}(5 \cos \theta) - \hat{j}(-5 \sin \theta) + \hat{k}(0) = (5 \cos \theta, 5 \sin \theta, 0) \\ \text{Oriented correctly} \end{array}$$

$$\vec{G}(\vec{R}(\theta, z)) = (5z \cos \theta, 5z \sin \theta, z^2) \quad \vec{G} \cdot \vec{N} = 25z \cos^2 \theta + 25z \sin^2 \theta = 25z$$

$$\iint_C \vec{G} \cdot d\vec{S} = \iint_C \vec{G} \cdot \vec{N} d\theta dz = \int_0^4 \int_0^{2\pi} 25z d\theta dz = 2\pi \cdot 25 \left[\frac{z^2}{2} \right]_0^4 = 400\pi$$