

Name\_\_\_\_\_

MATH 251 Final Exam Version B Fall 2016

Sections 504

P. Yasskin

1-13	/65	16	/10
14	/10	E.C.	/ 5
15	/20	Total	/110

Multiple Choice: (5 points each. No part credit.)

- Find the point where the line  $x = 3 - t$ ,  $y = 1 + 2t$ ,  $z = 1 - 2t$  intersects the plane  $x - 2y + z = -2$ . At this point  $x + y + z =$ 
  - 0
  - 1
  - 2
  - 3
  - 4
  
- Find the plane tangent to the graph of  $z = x^2y^3$  at the point  $(2, 1)$ . The  $z$ -intercept is
  - 24
  - 16
  - 4
  - 0
  - 4
  
- Find the line perpendicular to the cone  $z^2 - x^2 - y^2 = 0$  at the point  $P = (4, 3, 5)$ . This line intersects the  $xy$ -plane at
  - $(3, 4, 0)$
  - $(-4, -3, 0)$
  - $(8, 6, 0)$
  - $(-6, -8, 0)$
  - $(\frac{4}{5}, \frac{3}{5}, 0)$

4. A box currently has length  $L = 20$  cm which is increasing at 4 cm/sec, width  $W = 15$  cm which is increasing at 2 cm/sec, and height  $H = 12$  cm which is decreasing at 1 cm/sec. At what rate is the volume changing?
- $3 \text{ cm}^3/\text{sec}$
  - $7 \text{ cm}^3/\text{sec}$
  - $540 \text{ cm}^3/\text{sec}$
  - $900 \text{ cm}^3/\text{sec}$
  - $3600 \text{ cm}^3/\text{sec}$
5. In order to hide from the Dark Invader, Duke Skywalker flies the Millennium Eagle into a galactic dust storm. Currently, his position is  $P = (30, -20, 10)$  and his velocity is  $\vec{v} = (-4, 3, 12)$ . He measures that currently the dust density is  $\rho = 450$  and its gradient is  $\vec{\nabla}\rho = (2, -2, 1)$ . Find the current rate of change of the dust density as seen by Duke.
- 2
  - 2
  - 110
  - 448
  - 560
6. Under the same conditions as in #5, in what **unit** vector direction should Duke travel to **increase** the dust density as quickly as possible?
- $(-2, 2, -1)$
  - $(2, -2, 1)$
  - $\left(\frac{4}{13}, \frac{-3}{13}, \frac{-12}{13}\right)$
  - $\left(\frac{-4}{13}, \frac{3}{13}, \frac{12}{13}\right)$
  - $\left(\frac{2}{3}, \frac{-2}{3}, \frac{1}{3}\right)$

7. The point  $(-2, 1)$  is a critical point of the function  $f = \frac{2x - 4y - x^2y^2}{xy}$ .

Use the Second Derivative Test to classify the point.

- a. Local Minimum
- b. Local Maximum
- c. Inflection Point
- d. Saddle Point
- e. Test Fails

8. Find the mass of the plate between  $y = 3x$  and  $y = x^2$  if the surface density is  $\delta = xy$ .

- a.  $\frac{729}{16}$
- b.  $\frac{243}{8}$
- c.  $\frac{81}{8}$
- d.  $\frac{81}{4}$
- e.  $\frac{9}{2}$

9. Find the  $x$ -component of the center of mass of the plate between  $y = 3x$  and  $y = x^2$  if the surface density is  $\delta = xy$ .

- a.  $\frac{72}{35}$
- b.  $\frac{35}{72}$
- c.  $\frac{2187}{35}$
- d.  $\frac{35}{2172}$
- e.  $\frac{729}{35}$

10. Compute  $\oint \vec{F} \cdot d\vec{S}$  counterclockwise around the rectangle  $0 \leq x \leq 2\pi$  and  $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$  for  $\vec{F} = (y \cos x - \sin y, x \cos y + \sin x)$ .

HINT: Use the Fundamental Theorem of Calculus for Curves or Green's Theorem.

- a.  $-16\pi$
- b.  $-8\pi$
- c. 0
- d.  $8\pi$
- e.  $16\pi$

11. Compute  $\int_{(0,0,0)}^{(2,0,1)} \vec{F} \cdot d\vec{S}$  for  $\vec{F} = (2x + y, x + 2y, 2z)$  along the curve  $\vec{r}(t) = (t^2 + t^3, t^2 - t^3, \sqrt{t})$ .

HINT: Find a scalar potential.

- a. 1
- b. 3
- c. 5
- d. 7
- e. 9

12. Compute  $\iint \vec{\nabla} \times \vec{F} \cdot d\vec{S}$  over the sphere  $x^2 + y^2 + z^2 = 9$  with outward normal for  $\vec{F} = (xy^2z, yz^2x, zx^2y)$ .

HINT: Use Stokes' Theorem or Gauss' Theorem.

- a.  $4\pi$
- b.  $12\pi$
- c.  $\frac{32}{3}\pi$
- d.  $\frac{64}{3}\pi$
- e. 0

13. Compute  $\iint \vec{F} \cdot d\vec{S}$  outward through the complete surface of the cylinder  $x^2 + y^2 = 4$  for  $1 \leq z \leq 4$  for  $\vec{F} = (xz^2, yz^2, z^3)$ .


HINT: Use a Theorem.

- a.  $840\pi$
- b.  $420\pi$
- c.  $252\pi$
- d.  $210\pi$
- e.  $126\pi$

Work Out: (Points indicated. Part credit possible. Show all work.)

14. (10 points) Compute  $\iint_D x \, dx \, dy$  over the "diamond shaped" region  $D$  in the first quadrant bounded by the parabolas

$$\begin{aligned} y = 16 - x^2 & & y = 8 - x^2 \\ y = 4 + x^2 & \text{ and } & y = x^2 \end{aligned}$$




HINTS: Use the coordinates:  $u = y + x^2$ ,  $v = y - x^2$ . Solve for  $x$  and  $y$ .

15. (20 points) Verify Stokes' Theorem  $\iint_H \nabla \times \vec{F} \cdot d\vec{S} = \oint_{\partial H} \vec{F} \cdot d\vec{s}$

for the vector field  $\vec{F} = (y, -x, xz + yz)$

and the hemisphere  $z = \sqrt{9 - x^2 - y^2}$  oriented up.

Be sure to check the orientation. Use the following steps:



a. RHS: Parametrize the boundary curve and compute the line integral:

$$\vec{r}(\theta) =$$

$$\vec{v}(\theta) =$$

$$\vec{F}(\vec{r}(\theta)) =$$

$$\oint_{\partial H} \vec{F} \cdot d\vec{s} =$$

(continued)

b. LHS: Compute the surface integral using the parametrization:

$$\vec{R}(\varphi, \theta) = ( 3 \sin \varphi \cos \theta, 3 \sin \varphi \sin \theta, 3 \cos \varphi )$$

$$\vec{\nabla} \times \vec{F} =$$

$$[\vec{\nabla} \times \vec{F}]_{\vec{R}(\varphi, \theta)} =$$

$$\vec{e}_\varphi =$$

$$\vec{e}_\theta =$$

$$\vec{N} =$$

$$\iint_H \vec{\nabla} \times \vec{F} \cdot d\vec{S} =$$

16. (10 points) Find 3 positive numbers  $x$ ,  $y$  and  $z$ , whose sum is 120 such that  $f(x,y,z) = xy^2z^3$  is a maximum.

Solve either by Eliminating a Variable or by Lagrange Multipliers.

5 points extra credit for doing both. Clearly separate solutions.

METHOD 1: Lagrange Multipliers:

METHOD 2: Eliminate a Variable: