

Name _____

MATH 253H

Final Exam

Fall 2014

Sections 201-202

P. Yasskin

1-12	/48
10	/20
11	/10
12	/25
Total	/103

Multiple Choice: (4 points each. No part credit.)

1. If $z = f(x, y)$ where $x = r \cos \theta$ and $y = r \sin \theta$ then

a. $\frac{\partial z}{\partial r} = -\frac{\partial f}{\partial x} \sin \theta + \frac{\partial f}{\partial y} \cos \theta$

b. $\frac{\partial z}{\partial r} = \frac{\partial f}{\partial x} r \cos \theta + \frac{\partial f}{\partial y} r \sin \theta$

c. $\frac{\partial z}{\partial \theta} = \frac{\partial f}{\partial x} r \sin \theta - \frac{\partial f}{\partial y} r \cos \theta$

d. $\frac{\partial z}{\partial \theta} = -\frac{\partial f}{\partial x} r \sin \theta + \frac{\partial f}{\partial y} r \cos \theta$

e. $\frac{\partial z}{\partial \theta} = \frac{\partial f}{\partial x} r \cos \theta + \frac{\partial f}{\partial y} r \sin \theta$

2. If a car is currently at the point $P = (3, -4)$ and has velocity $\vec{v} = (-2, 1)$ at what rate is its distance from the origin currently changing?

a. $\frac{dD}{dt} = -2$

b. $\frac{dD}{dt} = -\frac{2}{5}$

c. $\frac{dD}{dt} = \frac{2}{5}$


d. $\frac{dD}{dt} = 2$

e. $\frac{dD}{dt} = \frac{11}{5}$

3. Find the point where the line $\vec{r}(t) = (4t + 1, 2t - 1, 3t)$ intersects the plane $\vec{R}(u, v) = (3 + u, 2 - v, u + v)$. At this point (x, y, z) what is $x + y + z$?
- 9
 - 0
 - 9
 - 18
 - 45
4. Ham Duet is flying the Centennial Eagle above the Death Star along the curve $\vec{r}(t) = (t^2, t^3, 8 - t^2)$. At $t = 2$, Ham releases a space torpedo which then glides with constant velocity equal to the velocity of the Centennial Eagle at the instant of release. Where does the torpedo hit the surface of the Death Star, which is at $z = 0$?
- $(12, 32, -4)$
 - $(8, 20, 0)$
 - $(4, 8, 4)$
 - $(0, -4, 8)$
 - $(8, 8 + 6\sqrt{2}, 0)$
5. Duke Skywalker is flying the Centennial Eagle through a dangerous polaron field whose density is given by $\delta = \frac{1}{\sqrt{x^2 + y^2 + z^2}}$. If Duke is currently at the point $P = (1, -2, 2)$ in what unit vector direction should he fly to **REDUCE** the polaron density as fast as possible?
- $\left(\frac{2}{3}, \frac{-1}{3}, \frac{1}{3}\right)$
 - $\left(\frac{-2}{3}, \frac{1}{3}, \frac{-1}{3}\right)$
 - $\left(\frac{-1}{3}, \frac{-2}{3}, \frac{-2}{3}\right)$
 - $\left(\frac{-1}{3}, \frac{2}{3}, \frac{-2}{3}\right)$
 - $\left(\frac{1}{3}, \frac{-2}{3}, \frac{2}{3}\right)$


6. Queen Leah is flying the Centennial Eagle above Planet Tattooo along the curve $\vec{r}(t) = (t^2, t^3, 8 - t^2)$. What is her tangential acceleration, a_T , at $t = \frac{1}{3}$?
- a. $\frac{10}{3}$
 - b. $\frac{5}{3}$
 - c. $\frac{10}{9}$
 - d. $\sqrt{3}$
 - e. $2\sqrt{3}$
7. Find the tangent plane to the graph of $z = \frac{1}{xy}$ at $(2, 3)$. The z -intercept is
- a. 0
 - b. $\frac{1}{2}$
 - c. 1
 - d. $\frac{1}{6}$
 - e. $\frac{1}{36}$
8. Find the tangent plane to the graph of the ellipsoid $x^2 + 3xy + 2y^2 + z^2 = 21$ at $(2, 1, 3)$. The z -intercept is
- a. 3
 - b. 4
 - c. 6
 - d. 7
 - e. 42

9. A wire has the shape of the spiral given in polar coordinates by $r = \theta$ from $\theta = 0$ to $\theta = \pi$ and has linear density $\rho = \sqrt{x^2 + y^2}$. Find its mass.



Hint: Parametrize the curve.

- a. $\frac{1}{3}(1 + \pi^2)^{3/2} - \frac{1}{3}$
 b. $\frac{2}{3}(1 + \pi^2)^{3/2} - \frac{2}{3}$
 c. $\frac{4}{3}(1 + \pi^2)^{3/2} - \frac{4}{3}$
 d. $\frac{2}{3}(1 + \pi^2)^{3/2}$
 e. $\frac{4}{3}(1 + \pi^2)^{3/2}$
10. Compute $\int \vec{F} \cdot d\vec{s}$ for $\vec{F} = (2x, 2y)$ counterclockwise around the piece of the spiral given in polar coordinates by $r = \theta$ from $\theta = \pi$ to $\theta = 2\pi$.



Hint: Use a theorem.

- a. π^2
 b. $2\pi^2$
 c. $3\pi^2$
 d. $4\pi^2$
 e. $5\pi^2$

11. Compute $\iint_D e^{x-y} dA$ over the region D of all points (x,y) such that $|x| + |y| \leq 1$.
Hints: Draw the region and use a change of coordinates with $u = x + y$ and $v = x - y$.
- a. $2e + \frac{2}{e}$
 - b. $e + \frac{1}{e}$
 - c. $2e - \frac{2}{e}$
 - d. $e - \frac{1}{e}$
 - e. $\frac{1}{e} - e$

12. Compute the integral $\iint \vec{F} \cdot d\vec{S}$ for $\vec{F} = (2xz^2, yz^2, z^3)$ over the complete surface of the solid between the paraboloid $z = x^2 + y^2$ and the plane $z = 4$, with outward normal.
Hint: Use a theorem.
- a. 96π
 - b. 192π
 - c. 384π
 - d. 768π
 - e. 0


Work Out: (Points indicated. Part credit possible. Show all work.)

13. (20 points) For each integral, plot the region of integration and then compute the integral.

a. $I = \int_0^{27} \int_{\sqrt[3]{y}}^3 e^{-x^4} dx dy$

b. $J = \int_{-\sqrt{2}}^{\sqrt{2}} \int_{|y|}^{\sqrt{4-y^2}} e^{-x^2-y^2} dx dy$

14. (10 points) Compute $\oint \vec{F} \cdot d\vec{s}$ for $\vec{F} = (3y - 2x^2, 3y^2 - 2x)$ over the complete boundary of the shape at the right, which is a square of side 2 under an isosceles triangle with height 3. If you use a theorem, name it.



15. (25 points) Verify Stokes' Theorem $\iint_P \vec{\nabla} \times \vec{F} \cdot d\vec{S} = \oint_{\partial P} \vec{F} \cdot d\vec{S}$

for the vector field $\vec{F} = (x^2z, x - z, -z^2x)$ and the paraboloid

$y = x^2 + z^2$ with $y \leq 9$ oriented to the right

Use the following steps:

First the Left Hand Side:

Parametrize the paraboloid as:

$$\vec{R}(r, \theta) = (r \cos \theta, r^2, r \sin \theta)$$

a. Compute the coordinate tangent vectors:

$$\vec{e}_r =$$

$$\vec{e}_\theta =$$

b. Compute the normal vector:

$$\vec{N} =$$


c. Compute the curl:

$$\vec{\nabla} \times \vec{F} =$$

d. Evaluate the curl on the surface:

$$\vec{\nabla} \times \vec{F} \Big|_{\vec{R}(r, \theta)} =$$

e. Compute the left hand side:



Second the Right Hand Side:

f. Parametrize the boundary circle:

$$\vec{r}(\theta) =$$

g. Compute the tangent vector:

$$\vec{v} =$$

h. Evaluate $\vec{F} = (x^2z, x - z, -z^2x)$ on the circle:

$$\vec{F}|_{\vec{r}(\theta)} =$$

i. Compute the right hand side: