

Math 304–504

Linear Algebra

**Lecture 6:  
Inverse matrix (continued).**

## Identity matrix

*Definition.* The **identity matrix** (or **unit matrix**) is a diagonal matrix with all diagonal entries equal to 1.

The  $n \times n$  identity matrix is denoted  $I_n$  or simply  $I$ .

$$I_1 = (1), \quad I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

In general, 
$$I = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix}.$$

**Theorem.** Let  $A$  be an arbitrary  $m \times n$  matrix.

Then  $I_m A = A I_n = A$ .

## Inverse matrix

*Definition.* Let  $A$  be an  $n \times n$  matrix. The **inverse** of  $A$  is an  $n \times n$  matrix, denoted  $A^{-1}$ , such that

$$AA^{-1} = A^{-1}A = I.$$

If  $A^{-1}$  exists then the matrix  $A$  is called **invertible**. Otherwise  $A$  is called **singular**.

*Basic properties of inverse matrices:*

- The inverse matrix (if it exists) is unique.
- If  $A$  is invertible, so is  $A^{-1}$ , and  $(A^{-1})^{-1} = A$ .
- If  $n \times n$  matrices  $A_1, A_2, \dots, A_k$  are invertible, so is  $A_1A_2 \dots A_k$ , and  $(A_1A_2 \dots A_k)^{-1} = A_k^{-1} \dots A_2^{-1}A_1^{-1}$ .

System of  $n$  linear equations in  $n$  variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \dots\dots\dots \\ a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n = b_n \end{cases}$$

$$\iff \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$$

**Theorem** If the matrix  $A = (a_{ij})$  is invertible then the system has a unique solution.

**Theorem** If an  $n \times n$  matrix  $A$  is invertible, then for any  $n$ -dimensional column vector  $\mathbf{b}$  the matrix equation  $A\mathbf{x} = \mathbf{b}$  has a unique solution, which is  $\mathbf{x} = A^{-1}\mathbf{b}$ .

Indeed,  $A(A^{-1}\mathbf{b}) = (AA^{-1})\mathbf{b} = I_n\mathbf{b} = \mathbf{b}$ .

Conversely, if  $A\mathbf{x} = \mathbf{b}$  then

$$\mathbf{x} = I_n\mathbf{x} = (A^{-1}A)\mathbf{x} = A^{-1}(A\mathbf{x}) = A^{-1}\mathbf{b}.$$

**Corollary** If the matrix equation  $A\mathbf{x} = \mathbf{0}$  has a nonzero solution then  $A$  is not invertible.

## Examples

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, \quad C = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.$$

---

$$AB = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

$$BA = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

$$C^2 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Thus  $A^{-1} = B$ ,  $B^{-1} = A$ , and  $C^{-1} = C$ .

## Examples

$$D = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}.$$

---

$$D^2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

It follows that  $D$  is a singular matrix as otherwise

$$D^2 = O \implies D^{-1}D^2 = D^{-1}O \implies D = O.$$

$$E^2 = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix} = 2E.$$

It follows that  $E$  is a singular matrix as otherwise

$$E^2 = 2E \implies E^2E^{-1} = 2EE^{-1} \implies E = 2I.$$

## Inverting diagonal matrices

**Theorem** A diagonal matrix  $D = \text{diag}(d_1, \dots, d_n)$  is invertible if and only if all diagonal entries are nonzero:  $d_i \neq 0$  for  $1 \leq i \leq n$ .

If  $D$  is invertible then  $D^{-1} = \text{diag}(d_1^{-1}, \dots, d_n^{-1})$ .

$$\begin{pmatrix} d_1 & 0 & \dots & 0 \\ 0 & d_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n \end{pmatrix}^{-1} = \begin{pmatrix} d_1^{-1} & 0 & \dots & 0 \\ 0 & d_2^{-1} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n^{-1} \end{pmatrix}$$



## Inverting diagonal matrices

**Theorem** A diagonal matrix  $D = \text{diag}(d_1, \dots, d_n)$  is invertible if and only if all diagonal entries are nonzero:  $d_i \neq 0$  for  $1 \leq i \leq n$ .

If  $D$  is invertible then  $D^{-1} = \text{diag}(d_1^{-1}, \dots, d_n^{-1})$ .

*Proof:* If all  $d_i \neq 0$  then, clearly,  
 $\text{diag}(d_1, \dots, d_n) \text{diag}(d_1^{-1}, \dots, d_n^{-1}) = \text{diag}(1, \dots, 1) = I$ ,  
 $\text{diag}(d_1^{-1}, \dots, d_n^{-1}) \text{diag}(d_1, \dots, d_n) = \text{diag}(1, \dots, 1) = I$ .

Now suppose that  $d_i = 0$  for some  $i$ . Then for any  $n \times n$  matrix  $B$  the  $i$ th row of the matrix  $DB$  is a zero row. Hence  $DB \neq I$ .

## Inverting $2 \times 2$ matrices

*Definition.* The **determinant** of a  $2 \times 2$  matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \text{ is } \det A = ad - bc.$$

**Theorem** A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  is invertible if and only if  $\det A \neq 0$ .

If  $\det A \neq 0$  then

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

**Theorem** A matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  is invertible if and only if  $\det A \neq 0$ . If  $\det A \neq 0$  then

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

*Proof:* It is easy to verify that

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = (ad - bc)I_2.$$

Clearly,  $O$  is not invertible since  $OB = O \neq I$  for any  $2 \times 2$  matrix  $B$ . Besides, if  $A$  is invertible then

$$AB = O \implies A^{-1}AB = A^{-1}O \implies B = O.$$

## Fundamental results on inverse matrices

**Theorem 1** Given a square matrix  $A$ , the following are equivalent:

- (i)  $A$  is invertible;
- (ii)  $\mathbf{x} = \mathbf{0}$  is the only solution of the matrix equation  $A\mathbf{x} = \mathbf{0}$ ;
- (iii) the row echelon form of  $A$  has no zero rows;
- (iv) the reduced row echelon form of  $A$  is the identity matrix.


**Theorem 2** Suppose that a sequence of elementary row operations converts a matrix  $A$  into the identity matrix.

Then the same sequence of operations converts the identity matrix into the inverse matrix  $A^{-1}$ .


**Theorem 3** For any  $n \times n$  matrices  $A$  and  $B$ ,

$$BA = I \iff AB = I.$$

*Row echelon form of a square matrix:*



noninvertible case



invertible case

*Row echelon form of a  $3 \times 3$  matrix:*

$$\begin{pmatrix} \boxed{1} & * & * \\ 0 & \boxed{1} & * \\ 0 & 0 & \boxed{1} \end{pmatrix}, \begin{pmatrix} \boxed{1} & * & * \\ 0 & \boxed{1} & * \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} \boxed{1} & * & * \\ 0 & 0 & \boxed{1} \\ 0 & 0 & 0 \end{pmatrix}, \\ \begin{pmatrix} 0 & \boxed{1} & * \\ 0 & 0 & \boxed{1} \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \boxed{1} & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & \boxed{1} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

*Reduced row echelon form of a  $3 \times 3$  matrix:*

$$\begin{pmatrix} \boxed{1} & 0 & 0 \\ 0 & \boxed{1} & 0 \\ 0 & 0 & \boxed{1} \end{pmatrix}, \begin{pmatrix} \boxed{1} & 0 & * \\ 0 & \boxed{1} & * \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} \boxed{1} & * & 0 \\ 0 & 0 & \boxed{1} \\ 0 & 0 & 0 \end{pmatrix}, \\ \begin{pmatrix} 0 & \boxed{1} & 0 \\ 0 & 0 & \boxed{1} \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \boxed{1} & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & \boxed{1} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

*Example.*  $A = \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}.$

*To check whether  $A$  is invertible, we convert it to row echelon form.*

Interchange the 1st row with the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 3 & -2 & 0 \\ -2 & 3 & 0 \end{pmatrix}$$

Add  $-3$  times the 1st row to the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ -2 & 3 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ -2 & 3 & 0 \end{pmatrix}$$

Add 2 times the 1st row to the 3rd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ 0 & 3 & 2 \end{pmatrix}$$

Multiply the 2nd row by  $-1/2$ :

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 3 & 2 \end{pmatrix}$$



$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 3 & 2 \end{pmatrix}$$

Add  $-3$  times the 2nd row to the 3rd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 0 & -2.5 \end{pmatrix}$$

Multiply the 3rd row by  $-2/5$ :

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} \boxed{1} & 0 & 1 \\ 0 & \boxed{1} & 1.5 \\ 0 & 0 & \boxed{1} \end{pmatrix}$$

*We already know that the matrix  $A$  is invertible.  
Let's proceed towards reduced row echelon form.*

Add  $-3/2$  times the 3rd row to the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Add  $-1$  times the 3rd row to the 1st row:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

To obtain  $A^{-1}$ , we need to apply the following sequence of elementary row operations to the identity matrix:

- interchange the 1st row with the 2nd row,
- add  $-3$  times the 1st row to the 2nd row,
- add 2 times the 1st row to the 3rd row,
- multiply the 2nd row by  $-1/2$ ,
- add  $-3$  times the 2nd row to the 3rd row,
- multiply the 3rd row by  $-2/5$ ,
- add  $-3/2$  times the 3rd row to the 2nd row,
- add  $-1$  times the 3rd row to the 1st row.

A convenient way to compute the inverse matrix  $A^{-1}$  is to merge the matrices  $A$  and  $I$  into one  $3 \times 6$  matrix  $(A | I)$ , and apply elementary row operations to this new matrix.

$$A = \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$(A | I) = \left( \begin{array}{ccc|ccc} 3 & -2 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

$$\left( \begin{array}{ccc|ccc} 3 & -2 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Interchange the 1st row with the 2nd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 3 & -2 & 0 & 1 & 0 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Add  $-3$  times the 1st row to the 2nd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Add 2 times the 1st row to the 3rd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

Multiply the 2nd row by  $-1/2$ :

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

Add  $-3$  times the 2nd row to the 3rd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & -2.5 & 1.5 & -2.5 & 1 \end{array} \right)$$

Multiply the 3rd row by  $-2/5$ :

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

Add  $-3/2$  times the 3rd row to the 2nd row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0.4 & 0 & 0.6 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

Add  $-1$  times the 3rd row to the 1st row:

$$\left( \begin{array}{ccc|ccc} 1 & 0 & 0 & 0.6 & 0 & 0.4 \\ 0 & 1 & 0 & 0.4 & 0 & 0.6 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$



Thus 
$$\begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}^{-1} = \begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix}.$$

That is,

$$\begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix} \begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$\begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix} \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$