

MATH 304

Linear Algebra

**Lecture 22:**

**Matrix of a linear transformation (continued).  
Similar matrices.**

## Linear transformation

*Definition.* Given vector spaces  $V_1$  and  $V_2$ , a mapping  $L : V_1 \rightarrow V_2$  is **linear** if

$$L(\mathbf{x} + \mathbf{y}) = L(\mathbf{x}) + L(\mathbf{y}),$$

$$L(r\mathbf{x}) = rL(\mathbf{x})$$

for any  $\mathbf{x}, \mathbf{y} \in V_1$  and  $r \in \mathbb{R}$ .

## Matrix transformations

**Theorem** Suppose  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a linear map. Then there exists an  $m \times n$  matrix  $A$  such that  $L(\mathbf{x}) = A\mathbf{x}$  for all  $\mathbf{x} \in \mathbb{R}^n$ . Columns of  $A$  are vectors  $L(\mathbf{e}_1), L(\mathbf{e}_2), \dots, L(\mathbf{e}_n)$ , where  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  is the standard basis for  $\mathbb{R}^n$ .

$$\mathbf{y} = A\mathbf{x} \iff \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$$\iff \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix} = x_1 \begin{pmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{pmatrix} + x_2 \begin{pmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{pmatrix} + \dots + x_n \begin{pmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{pmatrix}$$

## Basis and coordinates

If  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is a basis for a vector space  $V$ , then any vector  $\mathbf{v} \in V$  has a unique representation

$$\mathbf{v} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \dots + x_n\mathbf{v}_n,$$

where  $x_i \in \mathbb{R}$ . The coefficients  $x_1, x_2, \dots, x_n$  are called the **coordinates** of  $\mathbf{v}$  with respect to the ordered basis  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ .

The mapping

$$\text{vector } \mathbf{v} \mapsto \text{its coordinates } (x_1, x_2, \dots, x_n)$$


provides a one-to-one correspondence between  $V$  and  $\mathbb{R}^n$ . Besides, this mapping is **linear**.

## Change of coordinates

Let  $V$  be a vector space.

Let  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  be a basis for  $V$  and  $g_1 : V \rightarrow \mathbb{R}^n$  be the coordinate mapping corresponding to this basis.

Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$  be another basis for  $V$  and  $g_2 : V \rightarrow \mathbb{R}^n$  be the coordinate mapping corresponding to this basis.



The composition  $g_2 \circ g_1^{-1}$  is a linear mapping of  $\mathbb{R}^n$  to itself. It is represented as  $\mathbf{x} \mapsto U\mathbf{x}$ , where  $U$  is an  $n \times n$  matrix.

$U$  is called the **transition matrix** from  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  to  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ . Columns of  $U$  are coordinates of the vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  with respect to the basis  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ .

## Matrix of a linear transformation

Let  $V, W$  be vector spaces and  $f : V \rightarrow W$  be a linear map.

Let  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  be a basis for  $V$  and  $g_1 : V \rightarrow \mathbb{R}^n$  be the coordinate mapping corresponding to this basis.

Let  $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_m$  be a basis for  $W$  and  $g_2 : W \rightarrow \mathbb{R}^m$  be the coordinate mapping corresponding to this basis.

$$\begin{array}{ccc} V & \xrightarrow{f} & W \\ g_1 \downarrow & & \downarrow g_2 \\ \mathbb{R}^n & \longrightarrow & \mathbb{R}^m \end{array}$$

The composition  $g_2 \circ f \circ g_1^{-1}$  is a linear mapping of  $\mathbb{R}^n$  to  $\mathbb{R}^m$ . It is represented as  $\mathbf{x} \mapsto A\mathbf{x}$ , where  $A$  is an  $m \times n$  matrix.

$A$  is called the **matrix of  $f$**  with respect to bases  $\mathbf{v}_1, \dots, \mathbf{v}_n$  and  $\mathbf{w}_1, \dots, \mathbf{w}_m$ . Columns of  $A$  are coordinates of vectors  $f(\mathbf{v}_1), \dots, f(\mathbf{v}_n)$  with respect to the basis  $\mathbf{w}_1, \dots, \mathbf{w}_m$ .

*Examples.* •  $D : \mathcal{P}_3 \rightarrow \mathcal{P}_2$ ,  $(Dp)(x) = p'(x)$ .

Let  $A_D$  be the matrix of  $D$  with respect to the bases  $1, x, x^2$  and  $1, x$ . Columns of  $A_D$  are coordinates of polynomials  $D1, Dx, Dx^2$  w.r.t. the basis  $1, x$ .

$$D1 = 0, Dx = 1, Dx^2 = 2x \implies A_D = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

•  $L : \mathcal{P}_3 \rightarrow \mathcal{P}_3$ ,  $(Lp)(x) = p(x + 1)$ .

Let  $A_L$  be the matrix of  $L$  w.r.t. the basis  $1, x, x^2$ .

$L1 = 1$ ,  $Lx = 1 + x$ ,  $Lx^2 = (x + 1)^2 = 1 + 2x + x^2$ .

$$\implies A_L = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}$$

**Problem.** Consider a linear operator  $L$  on the vector space of  $2 \times 2$  matrices given by

$$L \begin{pmatrix} x & y \\ z & w \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} x & y \\ z & w \end{pmatrix}.$$

Find the matrix of  $L$  with respect to the basis

$$E_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, E_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, E_3 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, E_4 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

Let  $M_L$  denote the desired matrix.

It follows from the definition that  $M_L$  is a  $4 \times 4$  matrix whose columns are coordinates of the matrices

$$L(E_1), L(E_2), L(E_3), L(E_4)$$

with respect to the basis  $E_1, E_2, E_3, E_4$ .



$$L(E_1) = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 3 & 0 \end{pmatrix} = 1E_1 + 0E_2 + 3E_3 + 0E_4,$$

$$L(E_2) = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 3 \end{pmatrix} = 0E_1 + 1E_2 + 0E_3 + 3E_4,$$

$$L(E_3) = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 4 & 0 \end{pmatrix} = 2E_1 + 0E_2 + 4E_3 + 0E_4,$$

$$L(E_4) = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 2 \\ 0 & 4 \end{pmatrix} = 0E_1 + 2E_2 + 0E_3 + 4E_4.$$

Therefore

$$M_L = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \\ 3 & 0 & 4 & 0 \\ 0 & 3 & 0 & 4 \end{pmatrix}.$$

Thus the relation

$$\begin{pmatrix} x_1 & y_1 \\ z_1 & w_1 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} x & y \\ z & w \end{pmatrix}$$

is equivalent to the relation

$$\begin{pmatrix} x_1 \\ y_1 \\ z_1 \\ w_1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \\ 3 & 0 & 4 & 0 \\ 0 & 3 & 0 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ w \end{pmatrix}.$$

**Problem.** Consider a linear operator  $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,

$$L \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

Find the matrix of  $L$  with respect to the basis

$$\mathbf{v}_1 = (3, 1), \quad \mathbf{v}_2 = (2, 1).$$

Let  $N$  be the desired matrix. Columns of  $N$  are coordinates of the vectors  $L(\mathbf{v}_1)$  and  $L(\mathbf{v}_2)$  w.r.t. the basis  $\mathbf{v}_1, \mathbf{v}_2$ .

$$L(\mathbf{v}_1) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ 1 \end{pmatrix}, \quad L(\mathbf{v}_2) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \end{pmatrix}.$$

Clearly,  $L(\mathbf{v}_2) = \mathbf{v}_1 = 1\mathbf{v}_1 + 0\mathbf{v}_2$ .

$$L(\mathbf{v}_1) = \alpha\mathbf{v}_1 + \beta\mathbf{v}_2 \iff \begin{cases} 3\alpha + 2\beta = 4 \\ \alpha + \beta = 1 \end{cases} \iff \begin{cases} \alpha = 2 \\ \beta = -1 \end{cases}$$

$$\text{Thus } N = \begin{pmatrix} 2 & 1 \\ -1 & 0 \end{pmatrix}.$$

## Change of basis for a linear operator

Let  $L : V \rightarrow V$  be a linear operator on a vector space  $V$ .

Let  $A$  be the matrix of  $L$  relative to a basis  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$  for  $V$ . Let  $B$  be the matrix of  $L$  relative to another basis  $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n$  for  $V$ .

Let  $U$  be the transition matrix from the basis  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$  to  $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n$ .

$$\begin{array}{ccc} \boxed{\mathbf{a}\text{-coordinates of } \mathbf{v}} & \xrightarrow{A} & \boxed{\mathbf{a}\text{-coordinates of } L(\mathbf{v})} \\ U \downarrow & & \downarrow U \\ \boxed{\mathbf{b}\text{-coordinates of } \mathbf{v}} & \xrightarrow{B} & \boxed{\mathbf{b}\text{-coordinates of } L(\mathbf{v})} \end{array}$$

It follows that  $UA\mathbf{x} = BU\mathbf{x}$  for all  $\mathbf{x} \in \mathbb{R}^n \implies UA = BU$ .  
Then  $A = U^{-1}BU$  and  $B = UAU^{-1}$ .

**Problem.** Consider a linear operator  $L : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,

$$L \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

Find the matrix of  $L$  with respect to the basis  $\mathbf{v}_1 = (3, 1)$ ,  $\mathbf{v}_2 = (2, 1)$ .

Let  $S$  be the matrix of  $L$  with respect to the standard basis,  $N$  be the matrix of  $L$  with respect to the basis  $\mathbf{v}_1, \mathbf{v}_2$ , and  $U$  be the transition matrix from  $\mathbf{v}_1, \mathbf{v}_2$  to  $\mathbf{e}_1, \mathbf{e}_2$ . Then  $N = U^{-1}SU$ .

$$S = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix},$$

$$\begin{aligned} N &= U^{-1}SU = \begin{pmatrix} 1 & -2 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix} \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ -1 & 0 \end{pmatrix}. \end{aligned}$$

## Similarity

*Definition.* An  $n \times n$  matrix  $B$  is said to be **similar** to an  $n \times n$  matrix  $A$  if  $B = S^{-1}AS$  for some nonsingular  $n \times n$  matrix  $S$ .

*Remark.* Two  $n \times n$  matrices are similar if and only if they represent the same linear operator on  $\mathbb{R}^n$  with respect to different bases.

**Theorem** Similarity is an *equivalence relation*, which means that

- (i) any square matrix  $A$  is similar to itself;
- (ii) if  $B$  is similar to  $A$ , then  $A$  is similar to  $B$ ;
- (iii) if  $A$  is similar to  $B$  and  $B$  is similar to  $C$ , then  $A$  is similar to  $C$ .

**Theorem** Similarity is an *equivalence relation*, i.e.,

- (i) any square matrix  $A$  is similar to itself;
- (ii) if  $B$  is similar to  $A$ , then  $A$  is similar to  $B$ ;
- (iii) if  $A$  is similar to  $B$  and  $B$  is similar to  $C$ , then  $A$  is similar to  $C$ .

*Proof:* (i)  $A = I^{-1}AI$ .

(ii) If  $B = S^{-1}AS$  then  $A = SBS^{-1} = (S^{-1})^{-1}BS^{-1} = S_1^{-1}BS_1$ , where  $S_1 = S^{-1}$ .

(iii) If  $A = S^{-1}BS$  and  $B = T^{-1}CT$  then  
 $A = S^{-1}(T^{-1}CT)S = (S^{-1}T^{-1})C(TS) = (TS)^{-1}C(TS) = S_2^{-1}CS_2$ , where  $S_2 = TS$ .

**Theorem** If  $A$  and  $B$  are similar matrices then they have the same (i) determinant, (ii) trace = the sum of diagonal entries, (iii) rank, and (iv) nullity.