

MATH 311

Topics in Applied Mathematics I

Lecture 7:

Inverse matrix (continued).

Transpose of a matrix.

Inverse matrix

Definition. Let A be an $n \times n$ matrix. The **inverse** of A is an $n \times n$ matrix, denoted A^{-1} , such that

$$\boxed{AA^{-1} = A^{-1}A = I.}$$

If A^{-1} exists then the matrix A is called **invertible**.
Otherwise A is called **singular**.

System of n linear equations in n variables:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \dots\dots\dots \\ a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n = b_n \end{cases} \iff \mathbf{Ax} = \mathbf{b},$$

where

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}.$$

Theorem If the matrix A is invertible then the system has a unique solution, which is $\mathbf{x} = A^{-1}\mathbf{b}$.

General results on inverse matrices

Theorem 1 Given an $n \times n$ matrix A , the following conditions are equivalent:

- (i) A is invertible;
- (ii) $\mathbf{x} = \mathbf{0}$ is the only solution of the matrix equation $A\mathbf{x} = \mathbf{0}$;
- (iii) the matrix equation $A\mathbf{x} = \mathbf{b}$ has a unique solution for any n -dimensional column vector \mathbf{b} ;
- (iv) the row echelon form of A has no zero rows;
- (v) the reduced row echelon form of A is the identity matrix.

Theorem 2 Suppose that a sequence of elementary row operations converts a matrix A into the identity matrix.

Then the same sequence of operations converts the identity matrix into the inverse matrix A^{-1} .

Row echelon form of a square matrix:

A 7x7 matrix in row echelon form. The diagonal elements are represented by squares, and the rest of the matrix is filled with asterisks. A blue staircase line connects the bottom-right corner of each square to the top-left corner of the square in the row below it, indicating the positions of the leading ones.

invertible case

A 7x7 matrix in row echelon form, similar to the first one, but with two asterisks circled in red. The first red asterisk is in the third row, fourth column, and the second red asterisk is in the sixth row, seventh column. A blue staircase line is also present, connecting the bottom-right corner of each square to the top-left corner of the square in the row below it.

noninvertible case

For any matrix in row echelon form, the number of columns with leading entries equals the number of rows with leading entries. For a square matrix, also the number of columns *without* leading entries (i.e., the number of free variables in a related system of linear equations) equals the number of rows *without* leading entries (i.e., zero rows).

Row echelon form of a square matrix:

$$\begin{pmatrix} \square & * & * & * & * & * & * \\ & \square & * & * & * & * & * \\ & & \square & * & * & * & * \\ & & & \square & * & * & * \\ & & & & \square & * & * \\ & & & & & \square & * \\ & & & & & & \square \end{pmatrix}$$

invertible case

$$\begin{pmatrix} \square & * & * & * & * & * & * \\ & \square & * & * & * & * & * \\ & & \square & * & * & * & * \\ & & & \square & * & * & * \\ & & & & \square & * & * \\ & & & & & \square & * \\ & & & & & & \square \end{pmatrix}$$

noninvertible case

Hence the row echelon form of a square matrix A is either strict triangular or else it has a zero row. In the former case, the equation $A\mathbf{x} = \mathbf{b}$ *always* has a unique solution. In the latter case, $A\mathbf{x} = \mathbf{b}$ *never* has a unique solution. Also, in the former case the reduced row echelon form of A is I .

Example. $A = \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}.$

To check whether A is invertible, we convert it to row echelon form.

Interchange the 1st row with the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 3 & -2 & 0 \\ -2 & 3 & 0 \end{pmatrix}$$

Add -3 times the 1st row to the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ -2 & 3 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ -2 & 3 & 0 \end{pmatrix}$$

Add 2 times the 1st row to the 3rd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -2 & -3 \\ 0 & 3 & 2 \end{pmatrix}$$

Multiply the 2nd row by -0.5 :

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 3 & 2 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 3 & 2 \end{pmatrix}$$

Add -3 times the 2nd row to the 3rd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 0 & -2.5 \end{pmatrix}$$

Multiply the 3rd row by -0.4 :

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1.5 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} \boxed{1} & 0 & 1 \\ 0 & \boxed{1} & 1.5 \\ 0 & 0 & \boxed{1} \end{pmatrix}$$

We already know that the matrix A is invertible.

Let's proceed towards reduced row echelon form.

Add -1.5 times the 3rd row to the 2nd row:

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Add -1 times the 3rd row to the 1st row:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

To obtain A^{-1} , we need to apply the following sequence of elementary row operations to the identity matrix:

- interchange the 1st row with the 2nd row,
- add -3 times the 1st row to the 2nd row,
- add 2 times the 1st row to the 3rd row,
- multiply the 2nd row by -0.5 ,
- add -3 times the 2nd row to the 3rd row,
- multiply the 3rd row by -0.4 ,
- add -1.5 times the 3rd row to the 2nd row,
- add -1 times the 3rd row to the 1st row.

A convenient way to compute the inverse matrix A^{-1} is to merge the matrices A and I into one 3×6 matrix $(A | I)$, and apply elementary row operations to this new matrix.

$$A = \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$(A | I) = \left(\begin{array}{ccc|ccc} 3 & -2 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

$$\left(\begin{array}{ccc|ccc} 3 & -2 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Interchange the 1st row with the 2nd row:

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 3 & -2 & 0 & 1 & 0 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Add -3 times the 1st row to the 2nd row:

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ -2 & 3 & 0 & 0 & 0 & 1 \end{array} \right)$$

Add 2 times the 1st row to the 3rd row:

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & -2 & -3 & 1 & -3 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

Multiply the 2nd row by -0.5 :

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 3 & 2 & 0 & 2 & 1 \end{array} \right)$$

Add -3 times the 2nd row to the 3rd row:

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & -2.5 & 1.5 & -2.5 & 1 \end{array} \right)$$

Multiply the 3rd row by -0.4 :

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1.5 & -0.5 & 1.5 & 0 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

Add -1.5 times the 3rd row to the 2nd row:

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0.4 & 0 & 0.6 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right)$$

Add -1 times the 3rd row to the 1st row:

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 0.6 & 0 & 0.4 \\ 0 & 1 & 0 & 0.4 & 0 & 0.6 \\ 0 & 0 & 1 & -0.6 & 1 & -0.4 \end{array} \right) = (I | A^{-1})$$

Thus
$$\begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}^{-1} = \begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix}.$$

That is,

$$\begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix} \begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$\begin{pmatrix} \frac{3}{5} & 0 & \frac{2}{5} \\ \frac{2}{5} & 0 & \frac{3}{5} \\ -\frac{3}{5} & 1 & -\frac{2}{5} \end{pmatrix} \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Why does it work?

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} = \begin{pmatrix} a_1 & a_2 & a_3 \\ 2b_1 & 2b_2 & 2b_3 \\ c_1 & c_2 & c_3 \end{pmatrix},$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1+3a_1 & b_2+3a_2 & b_3+3a_3 \\ c_1 & c_2 & c_3 \end{pmatrix},$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} = \begin{pmatrix} a_1 & a_2 & a_3 \\ c_1 & c_2 & c_3 \\ b_1 & b_2 & b_3 \end{pmatrix}.$$

Proposition Any elementary row operation can be simulated as left multiplication by a certain matrix.

Elementary matrices

$$E = \begin{pmatrix} 1 & & & & & & & \\ \vdots & \ddots & & & & & & \\ 0 & \cdots & 1 & & & & & \\ \vdots & & \vdots & \ddots & & & & \\ 0 & \cdots & r & \cdots & 1 & & & \\ \vdots & & \vdots & & \vdots & \ddots & & \\ 0 & \cdots & 0 & \cdots & 0 & \cdots & 1 & \end{pmatrix} \begin{array}{l} \text{row \#}i \\ \\ \text{row \#}j \end{array}$$

To obtain the matrix EA from A , add r times the i th row to the j th row. To obtain the matrix AE from A , add r times the j th column to the i th column.

Why does it work? (continued)

Assume that a square matrix A can be converted to the identity matrix by a sequence of elementary row operations. Then $E_k E_{k-1} \dots E_2 E_1 A = I$, where E_1, E_2, \dots, E_k are elementary matrices simulating those operations.

Applying the same sequence of operations to the identity matrix, we obtain the matrix

$$B = E_k E_{k-1} \dots E_2 E_1 I = E_k E_{k-1} \dots E_2 E_1.$$

Thus $BA = I$. Besides, B is invertible since elementary matrices are invertible (why?). It follows that $A = B^{-1}$, then $B = A^{-1}$.

Transpose of a matrix

Definition. Given a matrix A , the **transpose** of A , denoted A^T , is the matrix whose rows are columns of A (and whose columns are rows of A). That is, if $A = (a_{ij})$ then $A^T = (b_{ij})$, where $b_{ij} = a_{ji}$.

Examples.
$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}^T = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix},$$

$$\begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix}^T = (7, 8, 9), \quad \begin{pmatrix} 4 & 7 \\ 7 & 0 \end{pmatrix}^T = \begin{pmatrix} 4 & 7 \\ 7 & 0 \end{pmatrix}.$$

Properties of transposes:

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(rA)^T = rA^T$
- $(AB)^T = B^T A^T$
- $(A_1 A_2 \dots A_k)^T = A_k^T \dots A_2^T A_1^T$
- $(A^{-1})^T = (A^T)^{-1}$

Definition. A square matrix A is said to be **symmetric** if $A^T = A$.

For example, any diagonal matrix is symmetric.

Proposition For any square matrix A the matrices $B = AA^T$ and $C = A + A^T$ are symmetric.

Proof:

$$B^T = (AA^T)^T = (A^T)^T A^T = AA^T = B,$$

$$C^T = (A + A^T)^T = A^T + (A^T)^T = A^T + A = C.$$