

MATH 311

Topics in Applied Mathematics I

**Lecture 10:**  
**Vector spaces.**

## Linear operations on vectors

Let  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  and  $\mathbf{y} = (y_1, y_2, \dots, y_n)$  be  $n$ -dimensional vectors, and  $r \in \mathbb{R}$  be a scalar.

*Vector sum:*  $\mathbf{x} + \mathbf{y} = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n)$

*Scalar multiple:*  $r\mathbf{x} = (rx_1, rx_2, \dots, rx_n)$

*Zero vector:*  $\mathbf{0} = (0, 0, \dots, 0)$

*Negative of a vector:*  $-\mathbf{y} = (-y_1, -y_2, \dots, -y_n)$

*Vector difference:*

$\mathbf{x} - \mathbf{y} = \mathbf{x} + (-\mathbf{y}) = (x_1 - y_1, x_2 - y_2, \dots, x_n - y_n)$

## Properties of linear operations

$$\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$$

$$(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$$

$$\mathbf{x} + \mathbf{0} = \mathbf{0} + \mathbf{x} = \mathbf{x}$$

$$\mathbf{x} + (-\mathbf{x}) = (-\mathbf{x}) + \mathbf{x} = \mathbf{0}$$

$$r(\mathbf{x} + \mathbf{y}) = r\mathbf{x} + r\mathbf{y}$$

$$(r + s)\mathbf{x} = r\mathbf{x} + s\mathbf{x}$$

$$(rs)\mathbf{x} = r(s\mathbf{x})$$

$$1\mathbf{x} = \mathbf{x}$$

$$0\mathbf{x} = \mathbf{0}$$

$$(-1)\mathbf{x} = -\mathbf{x}$$

## Linear operations on matrices

Let  $A = (a_{ij})$  and  $B = (b_{ij})$  be  $m \times n$  matrices, and  $r \in \mathbb{R}$  be a scalar.

*Matrix sum:*  $A + B = (a_{ij} + b_{ij})_{1 \leq i \leq m, 1 \leq j \leq n}$

*Scalar multiple:*  $rA = (ra_{ij})_{1 \leq i \leq m, 1 \leq j \leq n}$

*Zero matrix  $O$ :* all entries are zeros

*Negative of a matrix:*  $-A = (-a_{ij})_{1 \leq i \leq m, 1 \leq j \leq n}$

*Matrix difference:*  $A - B = (a_{ij} - b_{ij})_{1 \leq i \leq m, 1 \leq j \leq n}$

As far as the linear operations are concerned, the  $m \times n$  matrices have the same properties as  $mn$ -dimensional vectors.

## Vector space: informal description

*Vector space = linear space* = a set  $V$  of objects (called *vectors*) that can be added and scaled.

That is, for any  $\mathbf{u}, \mathbf{v} \in V$  and  $r \in \mathbb{R}$  expressions

$$\boxed{\mathbf{u} + \mathbf{v}} \text{ and } \boxed{r\mathbf{u}}$$

should make sense.

Certain restrictions apply. For instance,

$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u},$$

$$2\mathbf{u} + 3\mathbf{u} = 5\mathbf{u}.$$

That is, addition and scalar multiplication in  $V$  should be like those of  $n$ -dimensional vectors.

## Vector space: definition

*Vector space* is a set  $V$  equipped with two operations  $\alpha : V \times V \rightarrow V$  and  $\mu : \mathbb{R} \times V \rightarrow V$  that have certain properties (listed below).

The operation  $\alpha$  is called *addition*. For any  $\mathbf{u}, \mathbf{v} \in V$ , the element  $\alpha(\mathbf{u}, \mathbf{v})$  is denoted  $\mathbf{u} + \mathbf{v}$ .

The operation  $\mu$  is called *scalar multiplication*. For any  $r \in \mathbb{R}$  and  $\mathbf{u} \in V$ , the element  $\mu(r, \mathbf{u})$  is denoted  $r\mathbf{u}$ .

## Properties of addition and scalar multiplication (brief)

A1.  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$

A2.  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$

A3.  $\mathbf{x} + \mathbf{0} = \mathbf{0} + \mathbf{x} = \mathbf{x}$

A4.  $\mathbf{x} + (-\mathbf{x}) = (-\mathbf{x}) + \mathbf{x} = \mathbf{0}$

A5.  $r(\mathbf{x} + \mathbf{y}) = r\mathbf{x} + r\mathbf{y}$

A6.  $(r + s)\mathbf{x} = r\mathbf{x} + s\mathbf{x}$

A7.  $(rs)\mathbf{x} = r(s\mathbf{x})$

A8.  $1\mathbf{x} = \mathbf{x}$

## Properties of addition and scalar multiplication (detailed)

- A1.  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  for all  $\mathbf{x}, \mathbf{y} \in V$ .
- A2.  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  for all  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$ .
- A3. There exists an element of  $V$ , called the *zero vector* and denoted  $\mathbf{0}$ , such that  $\mathbf{x} + \mathbf{0} = \mathbf{0} + \mathbf{x} = \mathbf{x}$  for all  $\mathbf{x} \in V$ .
- A4. For any  $\mathbf{x} \in V$  there exists an element of  $V$ , denoted  $-\mathbf{x}$ , such that  $\mathbf{x} + (-\mathbf{x}) = (-\mathbf{x}) + \mathbf{x} = \mathbf{0}$ .
- A5.  $r(\mathbf{x} + \mathbf{y}) = r\mathbf{x} + r\mathbf{y}$  for all  $r \in \mathbb{R}$  and  $\mathbf{x}, \mathbf{y} \in V$ .
- A6.  $(r + s)\mathbf{x} = r\mathbf{x} + s\mathbf{x}$  for all  $r, s \in \mathbb{R}$  and  $\mathbf{x} \in V$ .
- A7.  $(rs)\mathbf{x} = r(s\mathbf{x})$  for all  $r, s \in \mathbb{R}$  and  $\mathbf{x} \in V$ .
- A8.  $1\mathbf{x} = \mathbf{x}$  for all  $\mathbf{x} \in V$ .



- Associativity of addition implies that a multiple sum  $\mathbf{u}_1 + \mathbf{u}_2 + \cdots + \mathbf{u}_k$  is well defined for any  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in V$ .

- **Subtraction** in  $V$  is defined as follows:  
 $\mathbf{x} - \mathbf{y} = \mathbf{x} + (-\mathbf{y})$ .

- Addition and scalar multiplication are called **linear operations**.

Given  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in V$  and  $r_1, r_2, \dots, r_k \in \mathbb{R}$ ,

$$\boxed{r_1\mathbf{u}_1 + r_2\mathbf{u}_2 + \cdots + r_k\mathbf{u}_k}$$

is called a **linear combination** of  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ .

## Examples of vector spaces

In most examples, addition and scalar multiplication are natural operations so that properties A1–A8 are easy to verify.

- $\mathbb{R}^n$ :  $n$ -dimensional coordinate vectors
- $\mathcal{M}_{m,n}(\mathbb{R})$ :  $m \times n$  matrices with real entries
- $\mathbb{R}^\infty$ : infinite sequences  $(x_1, x_2, \dots)$ ,  $x_i \in \mathbb{R}$

For any  $\mathbf{x} = (x_1, x_2, \dots)$ ,  $\mathbf{y} = (y_1, y_2, \dots) \in \mathbb{R}^\infty$  and  $r \in \mathbb{R}$  let  $\mathbf{x} + \mathbf{y} = (x_1 + y_1, x_2 + y_2, \dots)$ ,  $r\mathbf{x} = (rx_1, rx_2, \dots)$ . Then  $\mathbf{0} = (0, 0, \dots)$  and  $-\mathbf{x} = (-x_1, -x_2, \dots)$ .

- $\{\mathbf{0}\}$ : the trivial vector space
- $\mathbf{0} + \mathbf{0} = \mathbf{0}$ ,  $r\mathbf{0} = \mathbf{0}$ ,  $-\mathbf{0} = \mathbf{0}$ .

## Functional vector spaces

- $F(\mathbb{R})$ : the set of all functions  $f : \mathbb{R} \rightarrow \mathbb{R}$

Given functions  $f, g \in F(\mathbb{R})$  and a scalar  $r \in \mathbb{R}$ , let  $(f + g)(x) = f(x) + g(x)$  and  $(rf)(x) = rf(x)$  for all  $x \in \mathbb{R}$ .  
Zero vector:  $o(x) = 0$ . Negative:  $(-f)(x) = -f(x)$ .

- $C(\mathbb{R})$ : all continuous functions  $f : \mathbb{R} \rightarrow \mathbb{R}$

Linear operations are inherited from  $F(\mathbb{R})$ . We only need to check that  $f, g \in C(\mathbb{R}) \implies f+g, rf \in C(\mathbb{R})$ , the zero function is continuous, and  $f \in C(\mathbb{R}) \implies -f \in C(\mathbb{R})$ .

- $C^1(\mathbb{R})$ : all continuously differentiable functions  $f : \mathbb{R} \rightarrow \mathbb{R}$

- $C^\infty(\mathbb{R})$ : all smooth functions  $f : \mathbb{R} \rightarrow \mathbb{R}$

- $\mathcal{P}$ : all polynomials  $p(x) = a_0 + a_1x + \cdots + a_nx^n$

## Some general observations

- The zero vector is unique.

Suppose  $\mathbf{z}_1$  and  $\mathbf{z}_2$  are zero vectors. Then  $\mathbf{z}_1 + \mathbf{z}_2 = \mathbf{z}_2$  since  $\mathbf{z}_1$  is a zero vector and  $\mathbf{z}_1 + \mathbf{z}_2 = \mathbf{z}_1$  since  $\mathbf{z}_2$  is a zero vector. Hence  $\mathbf{z}_1 = \mathbf{z}_2$ .

- For any  $\mathbf{x} \in V$ , the negative  $-\mathbf{x}$  is unique.

Suppose  $\mathbf{y}$  and  $\mathbf{y}'$  are both negatives of  $\mathbf{x}$ . Let us compute the sum  $\mathbf{y}' + \mathbf{x} + \mathbf{y}$  in two ways:

$$(\mathbf{y}' + \mathbf{x}) + \mathbf{y} = \mathbf{0} + \mathbf{y} = \mathbf{y},$$

$$\mathbf{y}' + (\mathbf{x} + \mathbf{y}) = \mathbf{y}' + \mathbf{0} = \mathbf{y}'.$$

By associativity of the vector addition,  $\mathbf{y} = \mathbf{y}'$ .

## Some general observations

- (cancellation law)  $\mathbf{x} + \mathbf{y} = \mathbf{x}' + \mathbf{y}$  implies  $\mathbf{x} = \mathbf{x}'$  for any  $\mathbf{x}, \mathbf{x}', \mathbf{y} \in V$ .

If  $\mathbf{x} + \mathbf{y} = \mathbf{x}' + \mathbf{y}$  then  $(\mathbf{x} + \mathbf{y}) + (-\mathbf{y}) = (\mathbf{x}' + \mathbf{y}) + (-\mathbf{y})$ . By associativity,  $(\mathbf{x} + \mathbf{y}) + (-\mathbf{y}) = \mathbf{x} + (\mathbf{y} + (-\mathbf{y})) = \mathbf{x} + \mathbf{0} = \mathbf{x}$  and  $(\mathbf{x}' + \mathbf{y}) + (-\mathbf{y}) = \mathbf{x}' + (\mathbf{y} + (-\mathbf{y})) = \mathbf{x}' + \mathbf{0} = \mathbf{x}'$ . Hence  $\mathbf{x} = \mathbf{x}'$ .

- $0\mathbf{x} = \mathbf{0}$  for any  $\mathbf{x} \in V$ .

Indeed,  $0\mathbf{x} + \mathbf{x} = 0\mathbf{x} + 1\mathbf{x} = (0 + 1)\mathbf{x} = 1\mathbf{x} = \mathbf{x} = \mathbf{0} + \mathbf{x}$ .  
By the cancellation law,  $0\mathbf{x} = \mathbf{0}$ .

- $(-1)\mathbf{x} = -\mathbf{x}$  for any  $\mathbf{x} \in V$ .

Indeed,  $\mathbf{x} + (-1)\mathbf{x} = (-1)\mathbf{x} + \mathbf{x} = (-1)\mathbf{x} + 1\mathbf{x} = (-1 + 1)\mathbf{x} = 0\mathbf{x} = \mathbf{0}$ .

## Counterexample: dumb scaling

Consider the set  $V = \mathbb{R}^n$  with the standard addition and a nonstandard scalar multiplication:

$$\boxed{r \odot \mathbf{x} = \mathbf{0}} \quad \text{for any } \mathbf{x} \in \mathbb{R}^n \text{ and } r \in \mathbb{R}.$$

Properties A1–A4 hold because they do not involve scalar multiplication.

$$\text{A5. } r \odot (\mathbf{x} + \mathbf{y}) = r \odot \mathbf{x} + r \odot \mathbf{y} \quad \iff \mathbf{0} = \mathbf{0} + \mathbf{0}$$

$$\text{A6. } (r + s) \odot \mathbf{x} = r \odot \mathbf{x} + s \odot \mathbf{x} \quad \iff \mathbf{0} = \mathbf{0} + \mathbf{0}$$

$$\text{A7. } (rs) \odot \mathbf{x} = r \odot (s \odot \mathbf{x}) \quad \iff \mathbf{0} = \mathbf{0}$$

$$\text{A8. } 1 \odot \mathbf{x} = \mathbf{x} \quad \iff \mathbf{0} = \mathbf{x}$$

A8 is the only property that fails. As a consequence, property A8 does not follow from properties A1–A7.

## Counterexample: lazy scaling

Consider the set  $V = \mathbb{R}^n$  with the standard addition and a nonstandard scalar multiplication:

$$\boxed{r \odot \mathbf{x} = \mathbf{x}} \text{ for any } \mathbf{x} \in \mathbb{R}^n \text{ and } r \in \mathbb{R}.$$

Properties A1–A4 hold because they do not involve scalar multiplication.

$$\text{A5. } r \odot (\mathbf{x} + \mathbf{y}) = r \odot \mathbf{x} + r \odot \mathbf{y} \iff \mathbf{x} + \mathbf{y} = \mathbf{x} + \mathbf{y}$$

$$\text{A6. } (r + s) \odot \mathbf{x} = r \odot \mathbf{x} + s \odot \mathbf{x} \iff \mathbf{x} = \mathbf{x} + \mathbf{x}$$

$$\text{A7. } (rs) \odot \mathbf{x} = r \odot (s \odot \mathbf{x}) \iff \mathbf{x} = \mathbf{x}$$

$$\text{A8. } 1 \odot \mathbf{x} = \mathbf{x} \iff \mathbf{x} = \mathbf{x}$$

The only property that fails is A6.

## Weird example

Consider the set  $V = \mathbb{R}_+$  of positive numbers with a nonstandard addition and scalar multiplication:

$$\boxed{x \oplus y = xy} \quad \text{for any } x, y \in \mathbb{R}_+.$$

$$\boxed{r \odot x = x^r} \quad \text{for any } x \in \mathbb{R}_+ \text{ and } r \in \mathbb{R}.$$

$$\text{A1. } x \oplus y = y \oplus x \quad \iff xy = yx$$

$$\text{A2. } (x \oplus y) \oplus z = x \oplus (y \oplus z) \quad \iff (xy)z = x(yz)$$

$$\text{A3. } x \oplus \zeta = \zeta \oplus x = x \quad \iff x\zeta = \zeta x = x \text{ (holds for } \zeta = 1)$$

$$\text{A4. } x \oplus \eta = \eta \oplus x = 1 \quad \iff x\eta = \eta x = 1 \text{ (holds for } \eta = x^{-1})$$

$$\text{A5. } r \odot (x \oplus y) = (r \odot x) \oplus (r \odot y) \quad \iff (xy)^r = x^r y^r$$

$$\text{A6. } (r + s) \odot x = (r \odot x) \oplus (s \odot x) \quad \iff x^{r+s} = x^r x^s$$

$$\text{A7. } (rs) \odot x = r \odot (s \odot x) \quad \iff x^{rs} = (x^s)^r$$

$$\text{A8. } 1 \odot x = x \quad \iff x^1 = x$$