

## Sample problems for the final exam: Solutions

Any problem may be altered or replaced by a different one!

**Problem 1** Find the point of intersection of the planes  $x + 2y - z = 1$ ,  $x - 3y = -5$ , and  $2x + y + z = 0$  in  $\mathbb{R}^3$ .

The intersection point  $(x, y, z)$  is a solution of the system

$$\begin{cases} x + 2y - z = 1, \\ x - 3y = -5, \\ 2x + y + z = 0. \end{cases}$$

To solve the system, we convert its augmented matrix into reduced row echelon form using elementary row operations:

$$\begin{aligned} \left( \begin{array}{ccc|c} 1 & 2 & -1 & 1 \\ 1 & -3 & 0 & -5 \\ 2 & 1 & 1 & 0 \end{array} \right) &\rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 1 \\ 0 & -5 & 1 & -6 \\ 2 & 1 & 1 & 0 \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 1 \\ 0 & -5 & 1 & -6 \\ 0 & -3 & 3 & -2 \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 1 \\ 0 & -3 & 3 & -2 \\ 0 & -5 & 1 & -6 \end{array} \right) \\ &\rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 1 \\ 0 & 1 & -1 & \frac{2}{3} \\ 0 & -5 & 1 & -6 \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 1 \\ 0 & 1 & -1 & \frac{2}{3} \\ 0 & 0 & -4 & -\frac{8}{3} \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 1 \\ 0 & 1 & -1 & \frac{2}{3} \\ 0 & 0 & 1 & \frac{2}{3} \end{array} \right) \\ &\rightarrow \left( \begin{array}{ccc|c} 1 & 2 & -1 & 1 \\ 0 & 1 & 0 & \frac{4}{3} \\ 0 & 0 & 1 & \frac{2}{3} \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 2 & 0 & \frac{5}{3} \\ 0 & 1 & 0 & \frac{4}{3} \\ 0 & 0 & 1 & \frac{2}{3} \end{array} \right) \rightarrow \left( \begin{array}{ccc|c} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & \frac{4}{3} \\ 0 & 0 & 1 & \frac{2}{3} \end{array} \right). \end{aligned}$$

Thus the three planes intersect at the point  $(-1, \frac{4}{3}, \frac{2}{3})$ .

*Alternative solution:* The intersection point  $(x, y, z)$  is a solution of the system

$$\begin{cases} x + 2y - z = 1, \\ x - 3y = -5, \\ 2x + y + z = 0. \end{cases}$$

Adding all three equations, we obtain  $4x = -4$ . Hence  $x = -1$ . Substituting  $x = -1$  into the second equation, we obtain  $y = \frac{4}{3}$ . Substituting  $x = -1$  and  $y = \frac{4}{3}$  into the third equation, we obtain  $z = \frac{2}{3}$ . It is easy to check that  $x = -1$ ,  $y = \frac{4}{3}$ ,  $z = \frac{2}{3}$  is indeed a solution of the system. Thus  $(-1, \frac{4}{3}, \frac{2}{3})$  is the unique intersection point.

**Problem 2** Consider a linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by

$$L(\mathbf{v}) = (\mathbf{v} \cdot \mathbf{v}_1)\mathbf{v}_2, \quad \text{where } \mathbf{v}_1 = (1, 1, 1), \mathbf{v}_2 = (1, 2, 2).$$

(i) Find the matrix of the operator  $L$ .

Given  $\mathbf{v} = (x, y, z) \in \mathbb{R}^3$ , we have that  $\mathbf{v} \cdot \mathbf{v}_1 = x+y+z$  and  $L(\mathbf{v}) = (x+y+z, 2(x+y+z), 2(x+y+z))$ . Let  $A$  denote the matrix of the linear operator  $L$ . The columns of  $A$  are vectors  $L(\mathbf{e}_1), L(\mathbf{e}_2), L(\mathbf{e}_3)$ , where  $\mathbf{e}_1 = (1, 0, 0)$ ,  $\mathbf{e}_2 = (0, 1, 0)$ ,  $\mathbf{e}_3 = (0, 0, 1)$  is the standard basis for  $\mathbb{R}^3$ . Therefore

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 2 & 2 & 2 \end{pmatrix}.$$

(ii) Find the dimensions of the range and the kernel of  $L$ .

The range  $\text{Range}(L)$  of the linear operator  $L$  is the subspace of all vectors of the form  $L(\mathbf{v})$ , where  $\mathbf{v} \in \mathbb{R}^3$ . It is easy to see that  $\text{Range}(L)$  is the line spanned by the vector  $\mathbf{v}_2 = (1, 2, 2)$ . Hence  $\dim \text{Range}(L) = 1$ .

The kernel  $\ker(L)$  of the operator  $L$  is the subspace of all vectors  $\mathbf{x} \in \mathbb{R}^3$  such that  $L(\mathbf{x}) = \mathbf{0}$ . Clearly,  $L(\mathbf{x}) = \mathbf{0}$  if and only if  $\mathbf{x} \cdot \mathbf{v}_1 = 0$ . Therefore  $\ker(L)$  is the plane  $x + y + z = 0$  orthogonal to  $\mathbf{v}_1$  and passing through the origin. Its dimension is 2.

(iii) Find bases for the range and the kernel of  $L$ .

Since the range of  $L$  is the line spanned by the vector  $\mathbf{v}_2 = (1, 2, 2)$ , this vector is a basis for the range. The kernel of  $L$  is the plane given by the equation  $x + y + z = 0$ . The general solution of the equation is  $x = -t - s$ ,  $y = t$ ,  $z = s$ , where  $t, s \in \mathbb{R}$ . It gives rise to a parametric representation  $t(-1, 1, 0) + s(-1, 0, 1)$  of the plane. Thus the kernel of  $L$  is spanned by the vectors  $(-1, 1, 0)$  and  $(-1, 0, 1)$ . Since the two vectors are linearly independent, they form a basis for  $\ker(L)$ .

**Problem 3** Let  $\mathbf{v}_1 = (1, 1, 1)$ ,  $\mathbf{v}_2 = (1, 1, 0)$ , and  $\mathbf{v}_3 = (1, 0, 1)$ . Let  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a linear operator on  $\mathbb{R}^3$  such that  $L(\mathbf{v}_1) = \mathbf{v}_2$ ,  $L(\mathbf{v}_2) = \mathbf{v}_3$ ,  $L(\mathbf{v}_3) = \mathbf{v}_1$ .

(i) Show that the vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  form a basis for  $\mathbb{R}^3$ .

Let  $U$  be a  $3 \times 3$  matrix such that its columns are vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ :

$$U = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}.$$

To find the determinant of  $U$ , we subtract the second row from the first one and then expand by the first row:

$$\det U = \begin{vmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} = -1.$$

Since  $\det U \neq 0$ , the vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  are linearly independent. It follows that they form a basis for  $\mathbb{R}^3$ .

(ii) Find the matrix of the operator  $L$  relative to the basis  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ .

Let  $A$  denote the matrix of  $L$  relative to the basis  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ . By definition, the columns of  $A$  are coordinates of vectors  $L(\mathbf{v}_1), L(\mathbf{v}_2), L(\mathbf{v}_3)$  with respect to the basis  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ . Since  $L(\mathbf{v}_1) = \mathbf{v}_2 = 0\mathbf{v}_1 + 1\mathbf{v}_2 + 0\mathbf{v}_3$ ,  $L(\mathbf{v}_2) = \mathbf{v}_3 = 0\mathbf{v}_1 + 0\mathbf{v}_2 + 1\mathbf{v}_3$ ,  $L(\mathbf{v}_3) = \mathbf{v}_1 = 1\mathbf{v}_1 + 0\mathbf{v}_2 + 0\mathbf{v}_3$ , we obtain

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

(iii) Find the matrix of the operator  $L$  relative to the standard basis.

Let  $S$  denote the matrix of  $L$  relative to the standard basis for  $\mathbb{R}^3$ . We have  $S = UAU^{-1}$ , where  $A$  is the matrix of  $L$  relative to the basis  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  (already found) and  $U$  is the transition matrix from  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  to the standard basis (the vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  are consecutive columns of  $U$ ):

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}.$$

To find the inverse  $U^{-1}$ , we merge the matrix  $U$  with the identity matrix  $I$  into one  $3 \times 6$  matrix and apply row reduction to convert the left half  $U$  of this matrix into  $I$ . Simultaneously, the right half  $I$  will be converted into  $U^{-1}$ :

$$\begin{aligned} (U|I) &= \left( \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \left( \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \end{array} \right) \rightarrow \left( \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \end{array} \right) \\ &\rightarrow \left( \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & -1 & 1 & 0 \end{array} \right) \rightarrow \left( \begin{array}{ccc|ccc} 1 & 1 & 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & -1 & 1 & 0 \end{array} \right) \rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & 1 & 1 \\ 0 & -1 & 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & -1 & 1 & 0 \end{array} \right) \\ &\rightarrow \left( \begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 & -1 & 0 \end{array} \right) = (I|U^{-1}). \end{aligned}$$

Thus

$$\begin{aligned} S &= UAU^{-1} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} -1 & 1 & 1 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} -1 & 1 & 1 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 2 & -1 & -1 \end{pmatrix}. \end{aligned}$$

*Alternative solution:* Let  $S$  denote the matrix of  $L$  relative to the standard basis  $\mathbf{e}_1 = (1, 0, 0)$ ,  $\mathbf{e}_2 = (0, 1, 0)$ ,  $\mathbf{e}_3 = (0, 0, 1)$ . By definition, the columns of  $S$  are vectors  $L(\mathbf{e}_1), L(\mathbf{e}_2), L(\mathbf{e}_3)$ . It is easy to observe that  $\mathbf{e}_2 = \mathbf{v}_1 - \mathbf{v}_3$ ,  $\mathbf{e}_3 = \mathbf{v}_1 - \mathbf{v}_2$ , and  $\mathbf{e}_1 = \mathbf{v}_2 - \mathbf{e}_2 = -\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3$ . Therefore

$$\begin{aligned} L(\mathbf{e}_1) &= L(-\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3) = -L(\mathbf{v}_1) + L(\mathbf{v}_2) + L(\mathbf{v}_3) = -\mathbf{v}_2 + \mathbf{v}_3 + \mathbf{v}_1 = (1, 0, 2), \\ L(\mathbf{e}_2) &= L(\mathbf{v}_1 - \mathbf{v}_3) = L(\mathbf{v}_1) - L(\mathbf{v}_3) = \mathbf{v}_2 - \mathbf{v}_1 = (0, 0, -1), \\ L(\mathbf{e}_3) &= L(\mathbf{v}_1 - \mathbf{v}_2) = L(\mathbf{v}_1) - L(\mathbf{v}_2) = \mathbf{v}_2 - \mathbf{v}_3 = (0, 1, -1). \end{aligned}$$

Thus

$$S = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 2 & -1 & -1 \end{pmatrix}.$$

**Problem 4** Let  $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ .

(i) Find all eigenvalues of the matrix  $B$ .

The eigenvalues of  $B$  are roots of the characteristic equation  $\det(B - \lambda I) = 0$ . We obtain that

$$\begin{aligned}\det(B - \lambda I) &= \begin{vmatrix} 1 - \lambda & 1 & 1 \\ 1 & 1 - \lambda & 1 \\ 1 & 1 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^3 - 3(1 - \lambda) + 2 \\ &= (1 - 3\lambda + 3\lambda^2 - \lambda^3) - 3(1 - \lambda) + 2 = 3\lambda^2 - \lambda^3 = \lambda^2(3 - \lambda).\end{aligned}$$

Hence the matrix  $B$  has two eigenvalues: 0 and 3.

(ii) Find a basis for  $\mathbb{R}^3$  consisting of eigenvectors of  $B$ .

An eigenvector  $\mathbf{x} = (x, y, z)$  of  $B$  associated with an eigenvalue  $\lambda$  is a nonzero solution of the vector equation  $(B - \lambda I)\mathbf{x} = \mathbf{0}$ . First consider the case  $\lambda = 0$ . We obtain that

$$B\mathbf{x} = \mathbf{0} \iff \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \iff x + y + z = 0.$$

The general solution is  $x = -t - s$ ,  $y = t$ ,  $z = s$ , where  $t, s \in \mathbb{R}$ . Equivalently,  $\mathbf{x} = t(-1, 1, 0) + s(-1, 0, 1)$ . Hence the eigenspace of  $B$  associated with the eigenvalue 0 is two-dimensional. It is spanned by eigenvectors  $\mathbf{v}_1 = (-1, 1, 0)$  and  $\mathbf{v}_2 = (-1, 0, 1)$ .

Now consider the case  $\lambda = 3$ . We obtain that

$$\begin{aligned}(B - 3I)\mathbf{x} = \mathbf{0} &\iff \begin{pmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \\ &\iff \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \iff \begin{cases} x - z = 0, \\ y - z = 0. \end{cases}\end{aligned}$$

The general solution is  $x = y = z = t$ , where  $t \in \mathbb{R}$ . In particular,  $\mathbf{v}_3 = (1, 1, 1)$  is an eigenvector of  $B$  associated with the eigenvalue 3.

The vectors  $\mathbf{v}_1 = (-1, 1, 0)$ ,  $\mathbf{v}_2 = (-1, 0, 1)$ , and  $\mathbf{v}_3 = (1, 1, 1)$  are eigenvectors of the matrix  $B$ . They are linearly independent since the matrix whose rows are these vectors is nonsingular:

$$\begin{vmatrix} -1 & 1 & 0 \\ -1 & 0 & 1 \\ 1 & 1 & 1 \end{vmatrix} = 3 \neq 0.$$

It follows that  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  is a basis for  $\mathbb{R}^3$ .

(iii) Find an orthonormal basis for  $\mathbb{R}^3$  consisting of eigenvectors of  $B$ .

It is easy to check that the vector  $\mathbf{v}_3$  is orthogonal to  $\mathbf{v}_1$  and  $\mathbf{v}_2$ . To transform the basis  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  into an orthogonal one, we only need to orthogonalize the pair  $\mathbf{v}_1, \mathbf{v}_2$ . Using the Gram-Schmidt process, we replace the vector  $\mathbf{v}_2$  by

$$\mathbf{u} = \mathbf{v}_2 - \frac{\mathbf{v}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 = (-1, 0, 1) - \frac{1}{2}(-1, 1, 0) = (-1/2, -1/2, 1).$$

Now  $\mathbf{v}_1, \mathbf{u}, \mathbf{v}_3$  is an orthogonal basis for  $\mathbb{R}^3$ . Since  $\mathbf{u}$  is a linear combination of the vectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$ , it is also an eigenvector of  $B$  associated with the eigenvalue 0.

Finally, vectors  $\mathbf{w}_1 = \frac{\mathbf{v}_1}{\|\mathbf{v}_1\|}$ ,  $\mathbf{w}_2 = \frac{\mathbf{u}}{\|\mathbf{u}\|}$ , and  $\mathbf{w}_3 = \frac{\mathbf{v}_3}{\|\mathbf{v}_3\|}$  form an orthonormal basis for  $\mathbb{R}^3$  consisting of eigenvectors of  $B$ . We get that  $\|\mathbf{v}_1\| = \sqrt{2}$ ,  $\|\mathbf{u}\| = \sqrt{3/2}$ , and  $\|\mathbf{v}_3\| = \sqrt{3}$ . Thus

$$\mathbf{w}_1 = \frac{1}{\sqrt{2}}(-1, 1, 0), \quad \mathbf{w}_2 = \frac{1}{\sqrt{6}}(-1, -1, 2), \quad \mathbf{w}_3 = \frac{1}{\sqrt{3}}(1, 1, 1).$$

(iv) Find a diagonal matrix  $D$  and an invertible matrix  $U$  such that  $B = UDU^{-1}$ .

The vectors  $\mathbf{v}_1 = (-1, 1, 0)$ ,  $\mathbf{v}_2 = (-1, 0, 1)$ , and  $\mathbf{v}_3 = (1, 1, 1)$  are eigenvectors of the matrix  $B$  associated with eigenvalues 0, 0, and 3, respectively. Since these vectors form a basis for  $\mathbb{R}^3$ , it follows that  $B = UDU^{-1}$ , where

$$D = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad U = \begin{pmatrix} -1 & -1 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}.$$

Here  $U$  is the transition matrix from the basis  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  to the standard basis (its columns are vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ ) while  $D$  is the matrix of the linear operator  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ ,  $L(\mathbf{x}) = B\mathbf{x}$  with respect to the basis  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ .

**Problem 5** Let  $V$  be a subspace of  $\mathbb{R}^4$  spanned by vectors  $\mathbf{x}_1 = (1, 1, 0, 0)$ ,  $\mathbf{x}_2 = (2, 0, -1, 1)$ , and  $\mathbf{x}_3 = (0, 1, 1, 0)$ .

- (i) Find the distance from the point  $\mathbf{y} = (0, 0, 0, 4)$  to the subspace  $V$ .
- (ii) Find the distance from the point  $\mathbf{y}$  to the orthogonal complement  $V^\perp$ .

The vector  $\mathbf{y}$  is uniquely represented as  $\mathbf{y} = \mathbf{p} + \mathbf{o}$ , where  $\mathbf{p} \in V$  and  $\mathbf{o}$  is orthogonal to  $V$ , that is,  $\mathbf{o} \in V^\perp$ . The vector  $\mathbf{p}$  is the orthogonal projection of  $\mathbf{y}$  onto the subspace  $V$ . Since  $(V^\perp)^\perp = V$ , the vector  $\mathbf{o}$  is the orthogonal projection of  $\mathbf{y}$  onto the subspace  $V^\perp$ . It follows that the distance from the point  $\mathbf{y}$  to  $V$  equals  $\|\mathbf{o}\|$  while the distance from  $\mathbf{y}$  to  $V^\perp$  equals  $\|\mathbf{p}\|$ .

The orthogonal projection  $\mathbf{p}$  of the vector  $\mathbf{y}$  onto the subspace  $V$  is easily computed when we have an orthogonal basis for  $V$ . To get such a basis, we apply the Gram-Schmidt orthogonalization process to the basis  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ :

$$\begin{aligned} \mathbf{v}_1 &= \mathbf{x}_1 = (1, 1, 0, 0), & \mathbf{v}_2 &= \mathbf{x}_2 - \frac{\mathbf{x}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 = (2, 0, -1, 1) - \frac{2}{2}(1, 1, 0, 0) = (1, -1, -1, 1), \\ \mathbf{v}_3 &= \mathbf{x}_3 - \frac{\mathbf{x}_3 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{x}_3 \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2 = (0, 1, 1, 0) - \frac{1}{2}(1, 1, 0, 0) - \frac{-2}{4}(1, -1, -1, 1) = (0, 0, 1/2, 1/2). \end{aligned}$$

Now that  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  is an orthogonal basis for  $V$  we obtain

$$\begin{aligned} \mathbf{p} &= \frac{\mathbf{y} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 + \frac{\mathbf{y} \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2 + \frac{\mathbf{y} \cdot \mathbf{v}_3}{\mathbf{v}_3 \cdot \mathbf{v}_3} \mathbf{v}_3 = \\ &= \frac{0}{2}(1, 1, 0, 0) + \frac{4}{4}(1, -1, -1, 1) + \frac{2}{1/2}(0, 0, 1/2, 1/2) = (1, -1, 1, 3). \end{aligned}$$

Consequently,  $\mathbf{o} = \mathbf{y} - \mathbf{p} = (0, 0, 0, 4) - (1, -1, 1, 3) = (-1, 1, -1, 1)$ . Thus the distance from  $\mathbf{y}$  to the subspace  $V$  equals  $\|\mathbf{o}\| = 2$  and the distance from  $\mathbf{y}$  to  $V^\perp$  equals  $\|\mathbf{p}\| = \sqrt{12} = 2\sqrt{3}$ .

**Problem 6** Consider a vector field  $\mathbf{F}(x, y, z) = xyz\mathbf{e}_1 + xy\mathbf{e}_2 + x^2\mathbf{e}_3$ .

(i) Find  $\text{curl}(\mathbf{F})$ .

$$\begin{aligned} \text{curl}(\mathbf{F}) &= \begin{vmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xyz & xy & x^2 \end{vmatrix} = \left( \frac{\partial(x^2)}{\partial y} - \frac{\partial(xy)}{\partial z} \right) \mathbf{e}_1 + \left( \frac{\partial(xyz)}{\partial z} - \frac{\partial(x^2)}{\partial x} \right) \mathbf{e}_2 \\ &\quad + \left( \frac{\partial(xy)}{\partial x} - \frac{\partial(xyz)}{\partial y} \right) \mathbf{e}_3 = (xy - 2x)\mathbf{e}_2 + (y - xz)\mathbf{e}_3. \end{aligned}$$

(ii) Find the integral of the vector field  $\text{curl}(\mathbf{F})$  along a hemisphere  $H = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = 1, z \geq 0\}$ . Orient the hemisphere by the normal vector  $\mathbf{n} = (0, 0, 1)$  at the point  $(0, 0, 1)$ .

According to Stokes' Theorem,

$$\iint_H \text{curl}(\mathbf{F}) \cdot d\mathbf{S} = \oint_{\partial H} \mathbf{F} \cdot d\mathbf{s},$$

where the boundary  $\partial H$  is oriented consistently with  $H$ . The boundary is a circle,  $\partial H = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = 1, z = 0\}$ . It is parametrized (with the right orientation) by a path  $\mathbf{x} : [0, 2\pi] \rightarrow \mathbb{R}^3$ ,  $\mathbf{x}(t) = (\cos t, \sin t, 0)$ . We have  $\mathbf{F}(\mathbf{x}(t)) = (0, \cos t \sin t, \cos^2 t)$  and  $\mathbf{x}'(t) = (-\sin t, \cos t, 0)$ . Therefore

$$\oint_{\partial H} \mathbf{F} \cdot d\mathbf{s} = \int_{\mathbf{x}} \mathbf{F} \cdot d\mathbf{s} = \int_0^{2\pi} \mathbf{F}(\mathbf{x}(t)) \cdot \mathbf{x}'(t) dt = \int_0^{2\pi} \cos^2 t \sin t dt = -\frac{1}{3} \cos^3 t \Big|_{t=0}^{2\pi} = 0.$$

**Problem 7** Find the area of a pentagon with vertices  $(0, 0)$ ,  $(4, 0)$ ,  $(5, 2)$ ,  $(3, 4)$ , and  $(-1, 2)$ .

Segments  $(0, 0) - (3, 4)$  and  $(0, 0) - (5, 2)$  cut the pentagon into three triangles:  $\Delta_1$  with vertices  $(0, 0)$ ,  $(3, 4)$ , and  $(-1, 2)$ ;  $\Delta_2$  with vertices  $(0, 0)$ ,  $(5, 2)$ , and  $(3, 4)$ ; and  $\Delta_3$  with vertices  $(0, 0)$ ,  $(4, 0)$ , and  $(5, 2)$ . Since vectors  $\mathbf{v}_1 = (3, 4)$  and  $\mathbf{v}_2 = (-1, 2)$  are represented by two sides of the triangle  $\Delta_1$ , the area of that triangle equals  $\frac{1}{2}|\det A_1|$ , where

$$A_1 = (\mathbf{v}_1, \mathbf{v}_2) = \begin{pmatrix} 3 & -1 \\ 4 & 2 \end{pmatrix}.$$

Similarly, the area of the triangle  $\Delta_2$  equals  $\frac{1}{2}|\det A_2|$  and the area of  $\Delta_3$  equals  $\frac{1}{2}|\det A_3|$ , where

$$A_2 = \begin{pmatrix} 5 & 3 \\ 2 & 4 \end{pmatrix}, \quad A_3 = \begin{pmatrix} 4 & 5 \\ 0 & 2 \end{pmatrix}.$$

We obtain that  $\det A_1 = 10$ ,  $\det A_2 = 14$ , and  $\det A_3 = 8$ . Therefore the area of the pentagon equals  $\frac{1}{2}(10 + 14 + 8) = 16$ .