

MATH 614

Dynamical Systems and Chaos

Lecture 9:

Compact sets.

Definition of chaos.

Topological conjugacy

Suppose $f : X \rightarrow X$ and $g : Y \rightarrow Y$ are transformations of topological spaces.

Definition. We say that a map $\phi : X \rightarrow Y$ is a **semi-conjugacy** of f with g if ϕ is onto and $\phi \circ f = g \circ \phi$.



$$\begin{array}{ccc} X & \xrightarrow{f} & X \\ \phi \downarrow & & \downarrow \phi \\ Y & \xrightarrow{g} & Y \end{array}$$

The map ϕ is a **conjugacy** if, additionally, it is invertible.

The map ϕ is a **topological conjugacy** if, additionally, it is a homeomorphism, which means that both ϕ and ϕ^{-1} are continuous. In the latter case, we say that the maps f and g are **topologically conjugate**. Note that $f = \phi^{-1}g\phi$ and $g = \phi f \phi^{-1}$.

Unimodal maps

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a unimodal map, Λ be the set of all points $x \in \mathbb{R}$ such that $O_f^+(x) \subset [0, 1]$, and $S : \Lambda \rightarrow \Sigma_2 = \Sigma_{\{0,1\}}$ be the itinerary map.



Then S is a continuous semi-conjugacy of $f|_{\Lambda}$ with the shift. If S is a Cantor set, then S is one-to-one. Is S^{-1} continuous?

Example. $\phi : [0, 1) \cup [2, 3] \rightarrow [0, 2]$, $\phi(x) = x$ for $0 \leq x < 1$, $\phi(x) = x - 1$ for $2 \leq x \leq 3$.

The map ϕ is continuous and invertible, but the inverse is not continuous.

Compact sets

Definition. A subset E of a topological space X is **compact** if any covering of E by open sets admits a finite subcover. The subset E is **sequentially compact** if any sequence of its elements has a subsequence converging to an element of E .

Proposition 1 For any set $E \subset X$, compactness implies sequential compactness. If the topological space X is metrizable, then the converse is true as well.

Proposition 2 Any closed subset of a compact set is also compact.

We say that a topological space X is **Hausdorff** if any two distinct elements of X have disjoint neighborhoods. It is easy to show that any metrizable topological space is Hausdorff.

Proposition 3 In a Hausdorff topological space, every compact set is closed.

Proposition 4 A subset of the Euclidean space \mathbb{R}^n is compact if and only if it is closed and bounded.

Proposition 5 The topological space $\Sigma_{\mathcal{A}}$ of infinite words over a finite alphabet \mathcal{A} is compact.

Proof: Since the topological space $\Sigma_{\mathcal{A}}$ is metrizable, it is enough to prove sequential compactness. Suppose $\mathbf{s}^{(1)}, \mathbf{s}^{(2)}, \dots$ is a sequence of infinite words over the alphabet \mathcal{A} . Note that a subsequence $\mathbf{s}^{(n_1)}, \mathbf{s}^{(n_2)}, \mathbf{s}^{(n_3)}, \dots$ converges to some $\mathbf{s} \in \Sigma_{\mathcal{A}}$ if and only if every finite beginning of \mathbf{s} is also a beginning of $\mathbf{s}^{(n_k)}$ for k large enough.

Since \mathcal{A} is a finite set, the number of finite words over \mathcal{A} of any prescribed length is finite. It follows by induction that there exists a sequence of letters s_1, s_2, \dots such that for any $k \in \mathbb{N}$ the finite word $s_1 s_2 \dots s_k$ occurs as a beginning of $\mathbf{s}^{(n)}$ for infinitely many n 's. Then we choose indices $n_1 < n_2 < \dots$ so that $s_1 s_2 \dots s_k$ is a beginning of $\mathbf{s}^{(n_k)}$ for $k = 1, 2, \dots$. It follows that $\mathbf{s}^{(n_k)} \rightarrow \mathbf{s} = (s_1 s_2 s_3 \dots)$ as $k \rightarrow \infty$.

Compact sets and continuous maps

Proposition 6 The image of a compact set under a continuous map is also compact.

Proposition 7 Any continuous, real-valued function on a compact set attains its maximal and minimal values.

Proposition 8 Suppose that a continuous map $f : X \rightarrow Y$ is invertible. If the topological space X is compact and Y is Hausdorff, then the inverse map f^{-1} is continuous as well.

Proposition 9 Suppose (X, d) and (Y, ρ) are metric spaces. If X is compact then any continuous function $f : X \rightarrow Y$ is **uniformly continuous**, which means that for any $\varepsilon > 0$ there exists $\delta > 0$ such that $d(x, y) < \delta$ implies $\rho(f(x), f(y)) < \varepsilon$ for all $x, y \in X$.

Topological transitivity

Suppose $f : X \rightarrow X$ is a continuous transformation of a topological space X .

Definition. The map f is **topologically transitive** if for any nonempty open sets $U, V \subset X$ there exists a natural number n such that $f^n(U) \cap V \neq \emptyset$.

$$U \ni x \mapsto f(x) \mapsto f^2(x) \mapsto \cdots \mapsto f^n(x) \in V$$

Topological transitivity means that the dynamical system f is, in a sense, indecomposable.

Proposition 1 Topological transitivity is preserved under topological conjugacy.

Proposition 2 If the map f has a dense orbit, then it is topologically transitive provided X is Hausdorff and has no isolated points.

Proposition 3 If X is a metrizable compact space, then any topologically transitive transformation of X has a dense orbit.

Separation of orbits

Suppose $f : X \rightarrow X$ is a continuous transformation of a metric space (X, d) .

Definition. We say that f has **sensitive dependence on initial conditions** if there is $\delta > 0$ such that, for any $x \in X$ and a neighborhood U of x , there exist $y \in U$ and $n \geq 0$ satisfying $d(f^n(y), f^n(x)) > \delta$.

We say that the map f is **expansive** if there is $\delta > 0$ such that, for any $x, y \in X$, $x \neq y$, there exists $n \geq 0$ satisfying $d(f^n(y), f^n(x)) > \delta$.

Proposition If X is compact, then changing the metric d to another metric that induces the same topology cannot affect sensitive dependence on i.c. and expansiveness of the map f .

Corollary For continuous transformations of compact metric spaces, sensitive dependence on initial conditions and expansiveness are preserved under topological conjugacy.

Definition of chaos

Suppose $f : X \rightarrow X$ is a continuous transformation of a metric space (X, d) .

Definition. We say that the map f is **chaotic** if

- f has sensitive dependence on initial conditions;
- f is topologically transitive;
- periodic points of f are dense in X .

The three conditions provide the dynamical system f with unpredictability, indecomposability, and an element of regularity (recurrence).

Examples of chaotic systems

- The shift $\sigma : \Sigma_{\mathcal{A}} \rightarrow \Sigma_{\mathcal{A}}$ is chaotic.
- Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a unimodal map and Λ be the set of all points $x \in \mathbb{R}$ such that $O_f^+(x) \subset [0, 1]$. If Λ is a Cantor set then the restriction $f|_{\Lambda}$ of the map f to Λ is chaotic (otherwise it is not).

Recall that Λ is a Cantor set if and only if the itinerary map $S : \Lambda \rightarrow \Sigma_{\{0,1\}}$ is one-to-one, in which case S is a topological conjugacy of $f|_{\Lambda}$ with the shift on $\Sigma_{\{0,1\}}$.