

MATH 614

Dynamical Systems and Chaos

Lecture 34:

The Fatou components.

The filled Julia set.

The Julia and Fatou sets

Suppose $P : U \rightarrow U$ is a holomorphic map, where U is a domain in \mathbb{C} , the entire plane \mathbb{C} , or the Riemann sphere $\overline{\mathbb{C}}$.

Definition. The **Julia set** $J(P)$ of P is the closure (in U) of the set of repelling periodic points of P . The **Fatou set** $S(P)$ of P is the set of all points $z \in U$ such that the family of iterates $\{P^n\}_{n \geq 1}$ is normal at z .

- $J(P) \cap S(P) = \emptyset$ and $J(P) \cup S(P) = U$.
- $P(J(P)) = J(P)$ and $P^{-1}(J(P)) = J(P)$.
- $P(S(P)) \subset S(P)$ and $P^{-1}(S(P)) = S(P)$.
- If $U \subset \mathbb{C}$ and $\mathbb{C} \setminus U$ contains at least two points, then $S(P) = U$ and $J(P) = \emptyset$.
- If $S(P) \neq \emptyset$, then the Julia set has empty interior.
- If the Julia set is more than one repelling orbit, then it has no isolated points.
- If the Julia set is more than one repelling orbit, then the map P is chaotic on $J(P)$.

The Fatou components

The Fatou set $S(P)$ of a nonconstant holomorphic map $P : U \rightarrow U$ is open. Connected components of this set are called the **Fatou components** of P .

- For any Fatou component D of P , the image $P(D)$ is also a Fatou component of P .
- For any Fatou component D of a rational function P there exist integers $k \geq 0$ and $n \geq 1$ such that the Fatou component $P^k(D)$ is invariant under P^n (Sullivan 1986).
- Some transcendental functions P admit a Fatou component D that is a **wandering domain**, i.e., $D, P(D), P^2(D), \dots$ are disjoint sets.


The Fatou components

There are 5 types of invariant Fatou components for a holomorphic map $P : U \rightarrow U$:

- **immediate basin of attraction** of an attracting fixed point lying inside the component;
- **attracting petal** of a neutral fixed point lying on the boundary of the component;
- **Siegel disc**: the restriction of P to the component is holomorphically conjugate to a rotation of a disc;
- **Herman ring**: the restriction of P to the component is holomorphically conjugate to a rotation of an annulus;
- **Baker domain**: the iterates of P converge (uniformly on compact subsets of the component) to a constant $z_0 \notin U$ that is an essential singularity of P .


The Baker domains cannot occur for a rational function P .
The Herman rings cannot occur for functions $P : \mathbb{C} \rightarrow \mathbb{C}$.

Basin of attraction




$$P(z) = z^2 + \frac{i}{2}$$

Basin of attraction




$$P(z) = z^2 - 1$$

Attracting petal




$$P(z) = z^2 + \frac{1}{4}$$


Attracting petals



Siegel disc



Herman ring



Polynomial maps

From now on, we assume that P is a polynomial map with $\deg P \geq 2$:

$$P(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0,$$

where $a_n \neq 0$, $n \geq 2$. We consider P as a transformation of $\overline{\mathbb{C}}$.

Proposition The point at infinity is a super-attracting fixed point of P .

Proof: Clearly, $P(\infty) = \infty$. To find the derivative $P'(\infty)$, we need to compute the derivative $R'(0)$ of a rational function $R(z) = 1/P(1/z)$. Since $P(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0$, it follows that $R(z) = z^n / (a_n + a_{n-1} z + \cdots + a_1 z^{n-1} + a_0 z^n)$. Since $a_n \neq 0$ and $n \geq 2$, we obtain that $R'(0) = 0$.

The filled Julia set

Definition. The **filled Julia set** of the polynomial P , denoted $K(P)$, is the set of all points $z \in \mathbb{C}$ such that the orbit $z, P(z), P^2(z), \dots$ is bounded.

Proposition 1 The complement of $K(P)$ consists of points whose orbits escape to infinity.

Proposition 2 There is $R_0 > 0$ such that the set $\{z \in \mathbb{C} : |z| > R_0\}$ is contained in the Fatou set.

Proposition 3 The Julia set and the filled Julia set are bounded.

Proposition 4 The Julia set is contained in the filled Julia set.

More properties of the filled Julia set

- The filled Julia set is completely invariant:
 $P(K(P)) \subset K(P)$ and $P^{-1}(K(P)) \subset K(P)$.
- The complement of the filled Julia set is contained in the Fatou set.
- The filled Julia set is closed.
- The filled Julia set is nonempty.
- The interior of the filled Julia set is contained in the Fatou set.