

3. Linear Equations: Method of Integrating Factor (2.1)

1. A first order ODE is called **linear** if it is expressible in the form

$$y' + p(t)y = g(t) \quad (1)$$

where $p(t)$ and $g(t)$ are given functions.

2. If $g(t) \equiv 0$ then

$$y' + p(t)y = 0 \quad (2)$$

is called a **homogeneous** linear ODE. Otherwise (1) is called a **non-homogeneous** linear ODE.

3. The equation (1) is separable if and only if there is a real constant α such that $g(t) = \alpha p(t)$. (We already know how to solve it!)
4. The method to solve (1) for arbitrary $p(t)$ and $q(t)$ is called *Method of Integrating Factors*

Summary of the method of integrating factors

Step 1 Put ODE in the form (1).

Step 2 Find the integrating factor μ by solving the equation $\mu' = p(t)\mu$, i.e.,

$$\mu(t) = e^{\int p(t)dt}$$

Note: Any μ , satisfying $\mu' = p(t)\mu$, will suffice here, thus one can take the constant of integration $C = 0$ when calculating $\int p(t)dt$.

Step 3 Multiply both sides of (1) by μ and use the Product Rule for the left side to express the result as

$$(\mu(t)y(t))' = \mu g(x) \quad (3)$$

Step 4 Integrate both sides of (3). Note: Be sure to include the constant of integration in this step to get all solutions!

Step 5 Solve for the solution $y(t)$. In this step make sure to divide all terms by μ , including the term containing the constant of integration of the previous step.

5. Consider

$$y' - 3xy = -xe^{x^2}.$$

- (a) Find the general solution.
- (b) Find the solution satisfying the initial condition $y(0) = y_0$.
- (c) How do the solutions behave as x becomes large (i.e. $x \rightarrow +\infty$)? Does that behavior depend on the choice of the initial value y_0 ?

4. Models of mixing (from section 2.3) as an example of linear nonhomogeneous ODE

- (a) **Mixing Problems** serve as a model for problem of discharge and filtration of pollutants in a river, injection and absorption of medication in the bloodstream, for example.
- (b) *Chemicals in Tank, or Mixing.* Assume that a tank contains V gal of water and Q_0 lb of some substance (salt, or chemical, for example) is dissolved in it. Suppose that the water containing $\gamma(t)$ lb/gal of substance is entering the tank with the rate r gal/min (the concentration of the chemicals in the entering water varies in time) and then well stirred mixture is draining from the tank at the same rate. Find the amount $Q(t)$ of the substance in tank at any time (Do not solve, just find IVP for $Q(t)$).
- (c) *Chemicals in Pond* (from the textbook) A pond contains 10 million gal of fresh water. Stream water containing an undesirable chemical flows into the pond at the rate 5 million gal/yr, and the mixture in the pond flows out through an overflow culvert at the same rate. The concentration $\gamma(t)$ of chemical in the incoming water varies periodically with time t , measured in years, according to the expression $\gamma(t) = 2 + \sin(2t)$ g/gal.
- Construct a mathematical model of this flow process.
 - Determine the amount of chemical in the pond at any time.
 - Plot the solution and describe in words the effect of the variation of the incoming concentration.