

1 Mathematical Reasoning (Part I)¹

Statements

DEFINITION 1. *A statement is any declarative sentence or assertion that is either true or false.*

A statement cannot be neither true nor false and it cannot be both true and false.

1. The integer 5 is odd.
2. The integer 24277151704311 is prime.
3. $15 + 7 = 22$
4. Substitute the number 7 for x .
5. What is the derivative of $\cos x$?
6. Apple manufactures computers.
7. Apple manufactures the world's best computers.
8. Did you buy IBM?
9. I am telling a lie.
10. What happen when Pinocchio says: "My nose will grow now"?

- **Set Terminology and Notation (very short introduction²)**

DEFINITIONS:

Set is well-defined collection of objects.

Elements are objects or members of the set.

- **Roster notation:**

$A = \{a, b, c, d, e\}$ Read: Set A with elements a, b, c, d, e .

- **Indicating a pattern:**

$B = \{a, b, c, \dots, z\}$ Read: Set B with elements being the letters of the alphabet.

If a is an element of a set A , we write $a \in A$ that read " a belongs to A ." However, if a does not belong to A , we write $a \notin A$.

¹This part is covered in Sections 1.1-1.3 in the textbook.

²We will study SETS in Chapter 2!

Very common sets:

- \mathbf{R} is the set of all *real* numbers;
- $\mathbf{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$, the set of all *integers*;
- $\mathbf{Z}^+ = \{1, 2, 3, \dots\}$, the set of all *positive integers*;
- $\mathbf{N} = \{0, 1, 2, 3, \dots\}$, the set of all *natural* numbers;

Other sets:

- \mathbf{E} is the set of all *even integers*;
- \mathbf{O} is the set of all *odd integers*;
- $n\mathbf{Z}$ is the set of all integers multiples of n ($n \in \mathbf{Z}$);

An **open sentence** is any declarative sentence containing one or more variables, each variable representing a value in some prescribing set, called the **domain** of the variable, and which becomes a statement when values from their respective domains are substituted for these variables.

1. $P(x) : x + 5 = 7$
2. He is a student.

EXAMPLE 2. Discuss $P(x) : (x - 3)^2 \leq 1$ over \mathbf{Z} .

EXAMPLE 3. Discuss $P(x, y) : x^2 + y^2 = 1$ when $x, y \in \mathbf{R}$.

The NEGATION of a Statement

DEFINITION 4. If P is a statement, then the **negation** of P , written $\neg P$ (read “not P ”), is the statement “ P is false”.

Although $\neg P$ could always be expressed as

It is not the case that P .

there are usually better ways to express the statement $\neg P$.

1. P : The integer 77 is even.

2. P : $5^3 = 120$ $\neg P$: _____

3. P : The absolute value of the real number x is less than 5.

Compound Statements

Logical connectivity	write	read	meaning
Conjunction	$P \wedge Q$	P and Q	Both P and Q are true
Disjunction	$P \vee Q$	P or Q	P is true or Q is true

P : Ben is a student.

Q : Ben is a teaching assistant.

TRUTH TABLES

P	Q	$P \wedge Q$	$Q \wedge P$	$P \vee Q$	$Q \vee P$

EXAMPLE 5. Rewrite the following open sentences (over \mathbf{R}) using disjunction or conjunction.

(a) $P(x) : |x| \geq 10.$

(b) $P(x) : |x| < 10.$

(c) $P(x) : |4x + 7| \geq 23.$

Implications

DEFINITION 6. Let P and Q be statements. The **implication** $P \Rightarrow Q$ (read “ P implies Q ”) is the statement “If P is true, then Q is true.”

EXAMPLE 7. If n is odd, then $3n + 7$ is even.

The truth table for implication:

P	Q	$P \Rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

EXAMPLE 8. P : You earn an A on the final exam.

Q : You get an A for your final grade.

$P \Rightarrow Q$:

Different ways of expressing $P \Rightarrow Q$:

If P is true, then Q is true.

Q is true if P is true.

P implies Q .

P is true only if Q is true.

EXAMPLE 9. For a triangle T , let

$P(T)$: T is equilateral $Q(T)$: T is isosceles.

State $P(T) \Rightarrow Q(T)$ in a variety of ways:

Necessary and Sufficient Conditions

$P \Rightarrow Q$ also can be expressed as

P is sufficient for Q .

or

Q is necessary for P .

Equivalently,

In order for Q to be true it is sufficient that P be true.

or

Q must be true in order to P to be true.

EXAMPLE 10. Consider the following open sentences

$P(x)$: x is a multiple of 4. $Q(x)$: x is even. Complete:

- “For every integer integer x , $P(x) \Rightarrow Q(x)$ ” is _____.
- $P(x)$ is a _____ condition for Q to be true.
- $Q(x)$ is a _____ condition for $P(x)$ to be true.
- $Q(x)$ is not a _____ condition for $P(x)$ to be true.

EXAMPLE 11. Consider the following open sentences

$P(f)$: f is a differentiable function.

$Q(f)$: f is a continuous function.

Complete:

- “For every real-valued function f , $P(f) \Rightarrow Q(f)$ ” is _____.
- “For every real-valued function f , $Q(f) \Rightarrow P(f)$ ” is _____.

- $Q(f)$ is a _____ condition for f to be differentiable, but not a _____ condition.
- $P(f)$ is a _____ condition for f to be continuous.

REMARK 12. Note however, if $P \Rightarrow Q$ is true, then it is not necessary that P is true in order for Q to be true. Even if Q is true, P may be false.

Converse

DEFINITION 13. The statement $Q \Rightarrow P$ is called a **converse** of the statement $P \Rightarrow Q$.

EXAMPLE 14. If m and n are odd integers then $m + n$ is even.

Rewrite the statement in symbols. Then write its converse both in symbols and words.

EXAMPLE 15. P : The function $f(x) = \sin x$ is differentiable everywhere.

Q : The function $f(x) = \sin x$ is continuous everywhere.

$$P \Rightarrow Q$$

$$Q \Rightarrow P$$

Biconditional “ \Leftrightarrow ”

For statements P and Q ,

$$(P \Rightarrow Q) \wedge (Q \Rightarrow P)$$

is called the **biconditional** of P and Q and is denoted by $P \Leftrightarrow Q$. The biconditional $P \Leftrightarrow Q$ is stated as

“ P is equivalent to Q .” or “ P if and only if Q .” (or “ P iff Q .”)

or as “ P is a necessary and sufficient condition for Q .”

P	Q	$P \Rightarrow Q$	$Q \Rightarrow P$	$P \Leftrightarrow Q$
T	T			
T	F			
F	T			
F	F			

EXAMPLE 16. Let \vec{a} and \vec{b} be two non zero vectors. Then \vec{a} is orthogonal to \vec{b} iff $\vec{a} \cdot \vec{b} = 0$.

Tautologies and Contradictions

Tautology: statement that is always true

Contradiction: statement that is always false

P	$\neg P$	$P \vee (\neg P)$	$P \wedge (\neg P)$
T			
F			

Logical Equivalence

DEFINITION 17. Two compound statements are **logically equivalent** (write “ \equiv ”) if they have the same truth tables, which means they both are true or both are false.

Question: Are the statements $P \Rightarrow Q$ and $Q \Rightarrow P$ logically equivalent? _____

EXAMPLE 18. Let P and Q be statement forms. Determine whether the compound statements $\neg P \wedge Q$ and $\neg P \vee Q$ are logically equivalent (i.e. both true or both false).

P	Q			

REMARK 19. Let P and Q be statements. The biconditional $P \Leftrightarrow Q$ is a tautology if and only if P and Q are logically equivalent.

Some Fundamental Properties of Logical Equivalence

THEOREM 20. For the statement forms P , Q and R ,

- $\neg(\neg P) \equiv P$
- *Commutative Laws*
 $P \vee Q \equiv Q \vee P$
 $P \wedge Q \equiv Q \wedge P$
- *Associative Laws*
 $P \vee (Q \vee R) \equiv (P \vee Q) \vee R$
 $P \wedge (Q \wedge R) \equiv (P \wedge Q) \wedge R$
- *Distributive Laws*
 $P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)$
 $P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)$
- *De Morgan's Laws*
 $\neg(P \vee Q) \equiv (\neg P) \wedge (\neg Q)$
 $\neg(P \wedge Q) \equiv (\neg P) \vee (\neg Q)$

Proof. Each part of the theorem is verified by means of a truth table.

P	Q					

THEOREM 21. For statements P and Q ,

$$\neg(P \Rightarrow Q) \equiv P \wedge (\neg Q).$$

Proof.

Quantified Statements and Negations

EXAMPLE 22. Consider the following open sentence:

$$P(n) : \frac{2n^2 + 5 + (-1)^n}{2} \text{ is prime.}$$

How to convert this open sentence into a statement?

An open sentence can be made into a statement by using **quantifiers**.

Universal: $\forall x$ means for all/for every assigned value a of x .

Existential: $\exists x$ means that for some assigned values a of x .

Quantified statements

in symbols	in words
$\forall x \in D, P(x).$	For every $x \in D$, $P(x)$. If $x \in D$, then $P(x)$.
$\exists x \in D \ni P(x)$	There exists x such that $P(x)$.

Once a quantifier is applied to a variable, then the variable is called a **bound** variable. The variable that is not bound is called a **free** variable.

1. The area of a rectangle is its length times its width.

Quantifiers:

2. A triangle may be equilateral.

Quantifiers:

3. $15 - 5 = 10$

Quantifiers:

4. A real-valued function that is continuous at 0 is not necessarily differentiable at 0.

Quantifiers:

EXAMPLE 23. Rewrite the following statements in symbols using quantifiers. Introduce variables, where appropriate.

- a) For every real number x , $x + 5 = 7$.
-

- b) All positive real numbers have a square root.
-

- c) The sum of an even integer and an odd integer is even.
-

- d) For every integer n , either $n \leq 1$ or $n^2 \geq 4$.
-

NEGATIONS

(read “not P ”), is the statement “ P is false”.

1. All continuous functions are differentiable.

2. P : Every car on the parking lot #47 was with valid permit.

$\neg P$ _____

3. P : There exist real numbers a and b such that $(a + b)^2 = a^2 + b^2$.

$\neg P$ _____

Rules to negate statements with quantifiers:

$$\neg(\forall x \in D, P(x)) \equiv$$

$$\neg(\exists x \in D \ni P(x)) \equiv$$

$$\neg(\forall x \in D, (P(x) \vee Q(x))) \equiv$$

$$\neg(\forall x \in D, (P(x) \wedge Q(x))) \equiv$$

$$\neg(\exists x \in D \ni (P(x) \vee Q(x))) \equiv$$

$$\neg(\exists x \in D \ni (P(x) \wedge Q(x))) \equiv$$

EXAMPLE 24. Negate the statements below using the following steps:

1. Rewrite P in symbols using quantifiers.
2. Express the negation of P in symbols using the above rules.
3. Express $\neg P$ in words.

- a) P : If n is an odd integer then $3n + 7$ is odd.

P _____

$\neg P$ _____

$\neg P$ _____

b) P : There exists a positive integer n such that $m(n + 5) < 1$ for every integer m .

P _____
 $\neg P$ _____
 $\neg P$ _____

c) P : If n is an integer and n^2 is a multiple of 4 then n is a multiple of 4.

P _____
 $\neg P$ _____
 $\neg P$ _____

d) P : For every even integer n there exists an integer m such that $n = 2m$.

P _____
 $\neg P$ _____
 $\neg P$ _____

e) P : There exists a prime number p which is greater than 7 and less than 10.

P _____
 $\neg P$ _____
 $\neg P$ _____