

14.3: Partial Derivatives

DEFINITION 1. If f is a function of two variables, its **partial derivatives** are the functions f_x and f_y defined by

$$f_x(x, y) = \lim_{h \rightarrow 0} \frac{f(x + h, y) - f(x, y)}{h}$$

$$f_y(x, y) = \lim_{h \rightarrow 0} \frac{f(x, y + h) - f(x, y)}{h}$$

Conclusion: $f_x(x, y)$ represents the *rate of change* of the function $f(x, y)$ as we change x and hold y fixed while $f_y(x, y)$ represents the rate of change of $f(x, y)$ as we change y and hold x fixed.

Notations for partial derivatives: If $z = f(x, y)$, we write

$$f_x(x, y) = f_x = \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} f(x, y) = \frac{\partial z}{\partial x} = f_1 = D_1 f = D_x f$$
$$f_y(x, y) = f_y =$$

RULE FOR FINDING PARTIAL DERIVATIVES OF $z = f(x, y)$:

1. To find f_x , regard y as a constant and differentiate $f(x, y)$ with respect to x .
2. To find f_y , regard x as a constant and differentiate $f(x, y)$ with respect to y .

EXAMPLE 2. If $f(x, y) = x^3 + y^5 e^x$ find $f_x(0, 1)$ and $f_y(0, 1)$.

EXAMPLE 3. Find all of the first order partial derivatives for the following functions:

(a) $z(x, y) = x^3 \sin(xy)$

(c) $u(x, y, z) = ye^{xyz}$


EXAMPLE 4. The temperature at a point (x, y) on a flat metal plate is given by

$$T(x, y) = \frac{80}{1 + x^2 + y^2},$$

where T is measured in $^{\circ}\text{C}$ and x, y in meters. Find the rate of change of temperature with respect to distance at the point $(1, 2)$ in the y -direction.

Geometric interpretation of partial derivatives: Partial derivatives are the *slopes of traces*:

- $f_x(a, b)$ is the slope of the trace of the graph of $z = f(x, y)$ for the plane $y = b$ at the point (a, b) .



- $f_y(a, b)$ is the slope of the trace of the graph of $z = f(x, y)$ for the plane $x = a$ at (a, b) .

EXAMPLE 5. If $f(x, y) = \sqrt{4 - x^2 - 4y^2}$, find $f_x(1, 0)$ and $f_y(1, 0)$ and interpret these numbers as slopes. Illustrate with sketches.

Higher derivatives: Since both of the first order partial derivatives for $f(x, y)$ are also functions of x and y , so we can in turn differentiate each with respect to x or y . We use the following notation:

$$\begin{aligned}
 (f_x)_x &= f_{xx} = f_{11} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 z}{\partial x^2} \\
 (f_x)_y &= f_{xy} = f_{12} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 z}{\partial y \partial x} \\
 (f_y)_x &= f_{yx} = f_{21} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) = \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 z}{\partial x \partial y} \\
 (f_y)_y &= \quad = \quad = \quad = \quad =
 \end{aligned}$$

EXAMPLE 6. Find the second partial derivatives of

$$f(x, y) = y^3 + 5y^2e^{4x} - \cos(x^2).$$

Clairaut's Theorem. Suppose f is defined on a disk D that contains the point (a, b) . If the functions f_{xy} and f_{yx} are both continuous on D then

$$f_{xy}(a, b) = f_{yx}(a, b).$$

Partial derivative of order three or higher can also be defined. For instance,

$$f_{yyx} = (f_{yy})_x = \frac{\partial}{\partial x} \left(\frac{\partial^2 z}{\partial y^2} \right) = \frac{\partial^3 z}{\partial x \partial y^2}.$$

Using Clairaut's Theorem one can show that if the functions f_{yyx} , f_{xyy} and f_{yxy} are continuous then

EXAMPLE 7. Find the indicated derivative for

$$f(x, y, z) = \cos(xy + z).$$

(a) f_{xy}

(b) f_{zxy}

EXAMPLE 8. If f and g are twice differentiable functions of a single variable, show that the function

$$u(x, t) = f(x + at) + g(x - at)$$

is a solution of the wave equation

$$u_{tt} = a^2 u_{xx}.$$