

16.2: Line Integrals

Line integrals on plane: Let C be a plane curve with parametric equations:


$$x = x(t), y = y(t), \quad a \leq t \leq b,$$

or we can write the parametrization of the curve as a vector function:


$$\mathbf{r}(t) = \langle x(t), y(t) \rangle, \quad a \leq t \leq b.$$

DEFINITION 1. The line integral of $f(x, y)$ with respect to arc length, or the **line integral of f along C** is

$$\int_C f(x, y) ds$$



Definite integral - Area of a flat surface



Line integral - Area of a curved surface

<https://brilliant.org/wiki/line-integral/>

Recall that the *arc length* of a curve given by parametric equations $x = x(t), y = y(t), \quad a \leq t \leq b$ can be found as

$$L = \int_a^b ds,$$

where

$$ds = \sqrt{(x'(t))^2 + (y'(t))^2} dt.$$

The line integral is then

$$\int_C f(x, y) ds =$$

If we use the vector form of the parametrization we can simplify the notation up noticing that

$$\mathbf{r}'(t) = \langle x'(t), y'(t) \rangle$$

and then

$$ds = \sqrt{(x'(t))^2 + (y'(t))^2} dt =$$

Using this notation the line integral becomes,

$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t)) |\mathbf{r}'(t)| dt.$$

REMARK 2. The value of the line integral does not depend on the parametrization of the curve, provided that *the curve is traversed exactly once as t increases from a to b .*

Let us emphasize that $ds = |r'(t)| dt = \sqrt{[x'(t)]^2 + [y'(t)]^2} dt$.

EXAMPLE 3. Evaluate the line integral $\int_C y ds$, where $C : x = t^3, y = t^2, 0 \leq t \leq 1$.

Line integrals in space: Let C be a space curve with parametric equations:

$$x = x(t), y = y(t), z = z(t), \quad a \leq t \leq b,$$

or

$$\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}, \quad a \leq t \leq b.$$

The line integral of f along C is

$$\int_C f(x, y, z) ds = \int_a^b f(x(t), y(t), z(t)) |r'(t)| dt.$$

Here

$$ds = |r'(t)| dt = \sqrt{[x'(t)]^2 + [y'(t)]^2 + [z'(t)]^2} dt.$$

EXAMPLE 4. Evaluate the line integral $\int_C (x + y + z) ds$, where C is the line segment joining the points $A(-1, 1, 2)$ and $B(2, 3, 1)$.

Physical interpretation of a line integral: Let $\rho(x, y, z)$ represents the linear density at a point (x, y, z) of a thin wire shaped like a curve C . Then the **mass** m of the wire is:

$$m = \int_C \rho(x, y, z) \, ds.$$

EXAMPLE 5. A thin wire with the linear density $\rho(x, y) = x^2 + 2y^2$ takes the shape of the curve C which consists of the arc of the circle $x^2 + y^2 = 1$ from $(1, 0)$ to $(0, 1)$. Find the mass of the wire.

Line integrals with respect to x, y , and z . Let C be a space curve with parametric equations:

$$x = x(t), y = y(t), z = z(t), \quad a \leq t \leq b,$$

The **line integral of f with respect to x** is,

$$\int_C f(x, y, z) \, dx = \int_a^b f(x(t), y(t), z(t))x'(t) \, dt.$$

The **line integral of f with respect to y** is,

$$\int_C f(x, y, z) \, dy = \int_a^b f(x(t), y(t), z(t))y'(t) \, dt.$$

The **line integral of f with respect to z** is,

$$\int_C f(x, y, z) \, dz =$$

These integrals often appear together by the following notation:

$$\int_C P \, dx + Q \, dy + R \, dz$$

or

$$\int_C P \, dx + Q \, dy.$$

EXAMPLE 6. Compute

$$I = \int_C -\frac{y}{x^2 + y^2} dx + \frac{x}{x^2 + y^2} dy,$$

where C is the circle $x^2 + y^2 = 1$ oriented in the counterclockwise direction.

Line integrals of vector fields.

PROBLEM: Given a continuous force field,

$$\mathbf{F}(x, y, z) = P(x, y, z)\mathbf{i} + Q(x, y, z)\mathbf{j} + R(x, y, z)\mathbf{k},$$

such as a gravitational field. Find the work done by the force \mathbf{F} in moving a particle along a curve

$$C : \quad \mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}, \quad a \leq t \leq b.$$

DEFINITION 7. Let \mathbf{F} be a continuous vector field defined on a curve C given by a vector function $\mathbf{r}(t)$, $a \leq t \leq b$. Then the **line integral of \mathbf{F} along C** is

$$\int_C \mathbf{F} \cdot d\mathbf{r}(t) = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt.$$

REMARK 8. Note that this integral depends on the curve orientation:

$$\int_{-C} \mathbf{F} \cdot d\mathbf{r}(t) = - \int_C \mathbf{F} \cdot d\mathbf{r}(t)$$

Relationship between line integrals of vector fields and line integrals with respect to x, y , and z .

$$\int_C \mathbf{F} \cdot d\mathbf{r}(t) =$$

EXAMPLE 9. Find the work done by the force field $\mathbf{F}(x, y, z) = \langle xy, yz, xz \rangle$ in moving a particle along the curve $C : \mathbf{r}(t) = \langle t, t^2, t^3 \rangle, 0 \leq t \leq 1$.