

## 16.7: Surface Integrals

*Problem:* Find the **mass** of a thin sheet (say, of aluminum foil) which has a shape of a surface  $S$  and the density (mass per unit area) at the point  $(x, y, z)$  is  $\rho(x, y, z)$ .

*Solution:*

If  $S$  is given by  $\mathbf{r}(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k}$ ,  $(u, v) \in D$ , then the **surface integral of  $f$  over the surface  $S$**  is:

$$\iint_S f(x, y, z) \, dS = \iint_D f(\mathbf{r}(u, v)) |\mathbf{N}(u, v)| \, dA =$$

**EXAMPLE 1.** Find the mass of a thin funnel in the shape of a cone  $z = \sqrt{x^2 + y^2}$  inside the cylinder  $x^2 + y^2 \leq 2x$ , if its density is a function  $\rho(x, y, z) = x^2 + y^2 + z^2$ .

- **Oriented surfaces.** We consider only two-sided surfaces.

Let a surface  $S$  has a tangent plane at every point (except at any boundary points). There are two unit normal vectors at  $(x, y, z)$ :  $\hat{\mathbf{n}}$  and  $-\hat{\mathbf{n}}$ .

If it is possible to choose a unit normal vector  $\hat{\mathbf{n}}$  at every point  $(x, y, z)$  of a surface  $S$  so that  $\hat{\mathbf{n}}$  varies continuously over  $S$ , then  $S$  is called **oriented surface** and the given choice of  $\hat{\mathbf{n}}$  provides  $S$  with an **orientation**. There are two possible orientations for any orientable surface:

*Convention: For closed surfaces the positive orientation is outward.*

- **Surface integrals of vector fields.**

DEFINITION 2. If  $\mathbf{F}$  is a continuous vector field defined on an oriented surface  $S$  with unit normal vector  $\hat{\mathbf{n}}$ , then the **surface integral of  $\mathbf{F}$  over  $S$**  is

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot \hat{\mathbf{n}} dS.$$

This integral is also called the **flux of  $\mathbf{F}$  across  $S$** .

Note that if  $S$  is given by  $\mathbf{r}(u, v)$ ,  $(u, v) \in D$ , then

$$\hat{\mathbf{n}} = \frac{\mathbf{n}}{|\mathbf{n}|} =$$

and

$$d\mathbf{S} =$$

Finally,

$$\iint_S \mathbf{F} \cdot d\mathbf{S} =$$

EXAMPLE 3. Find the flux of the vector field

$$\mathbf{F} = \langle x^2, y^2, z^2 \rangle$$

across the surface

$$S = \{z^2 = x^2 + y^2, 0 \leq z \leq 2\}.$$