

due January 26, 2017 at the beginning of class

Topics covered : equations $y' = ay + b$, where a and b are constant, the model of falling object in air resistance (section 1.2 and week 1 lecture notes), separable equations (section 2.2 and week 1 lecture notes), equations that can be reduced to separable equations by an appropriate substitution (beginning of lecture notes of week 2). *You do not need to use calculator for this assignment.*

1. The **half-life** of a radioactive material is the time required for an amount of the material to decay to one-half of its original value. Show that for any radioactive material that decays according to the equation $Q' = -rQ$, $r > 0$, the half-life is equal to $\frac{\ln 2}{r}$ (in particular, it does not depend on the original value $Q(0)$).
2. Assume that the velocity v of the falling object satisfies the following differential equation:

$$v'(t) = 9.8 - \frac{v(t)}{a} \quad (1)$$

where a is a positive constant.

- (a) Find a number v_e such that $v(t) \equiv v_e$ is a solution of equation (1) (in other words find the equilibrium solution of (1)).
 - (b) Solve the equation (1) with initial condition $v(0) = 4.9a$. What is the limit of this solution when $t \rightarrow +\infty$? How this limiting velocity is related to your answer in the item (a)?
 - (c) Find the time (from the initial time moment is 0) that must elapse for the object to reach 90% of the limiting velocity found in the item (b).
 - (d) How far does the object fall in the time found in the item (c).
3. Solve the following differential equations:

(a) Find the general solution of $2t + ty^2 + e^{t^2}yy' = 0$ (please express y as a function of x explicitly).

(b) Find the solution of the initial value problem $(xy^2 + x)\frac{dy}{dx} + x^2y - y = 0$, $y(1) = e$.

4. Before attempting this problem review the lecture notes from week 2, discussing the equations of the type $y' = f(\frac{y}{x})$ (so-called, homogeneous equations) and $y' = f(ax + by + c)$: the main idea here is to make an appropriate substitution to obtain a separable equation: $u(x) = \frac{y(x)}{x}$ in the first case and $u(x) = ax + by(x) + c$ in the second case. Then find the general solution of the following equations:

(a) $y' = \frac{y - 2x}{2x + y}$;

(b) $y' = (3x - 2y - 1)^2$.