

## Homework Assignment 4 in MATH308-Spring 2016, Regular section

due February 23, 2017

Topics covered : *Wronskian, fundamental set of solutions of linear homogeneous equations of second order. linear homogeneous equations of second order with constant coefficients (sections 3.1-3.4).*

- (a) Calculate the Wronskian of the pair of the functions  $\sin^2 t, 1 - \cos 2t$ .  
(b) Assume that  $y_1(t)$  and  $y_2(t)$  are two solutions of the equation

$$y'' + p(t)y' + q(t)y = 0 \quad (1)$$

on the interval  $(-1, 1)$ , where the functions  $p(t)$  and  $q(t)$  are continuous on the same interval. It is known that  $y_1(0) = 2, y_1'(0) = -4, y_2(0) = -3,$  and  $y_2'(0) = 5$ . Is it true that the general solution of (1) is  $C_1y_1(t) + C_2y_2(t)$ ? Justify your answer.

- (c) Answer the same question as in the previous item if  $y_1(0) = 2, y_1'(0) = -4, y_2(0) = -3,$  but  $y_2'(0) = 6$ . Justify your answer. How  $y_1$  and  $y_2$  are related in this case (namely, express  $y_2(t)$  in terms of  $y_1(t)$ , if possible)?  
(d) (**bonus-10 points**) Assume again that  $y_1(t)$  and  $y_2(t)$  are solutions of (1) on the interval  $(-1, 1)$ . Prove that if  $y_1(t)$  and  $y_2(t)$  achieve a maximum or minimum at the same point, then they cannot form a fundamental set of solutions on this interval.  
(e) (**bonus-10 points**) Suppose that the Wronskian of any two solutions of (1) is constant in time. Prove that  $p(t) = 0$ . (Hint: Use the Abel theorem)
- Consider differential equations  $\frac{1}{3}y'' - y' + \alpha y = 0$ , where  $\alpha$  is a parameter.

- (a) In each of the following 3 cases find the general solution of the differential equation corresponding to the given  $\alpha$ , if:

i)  $\alpha = \frac{5}{12}$ ;                      ii)  $\alpha = \frac{3}{4}$ ;                      iii)  $\alpha = \frac{13}{12}$ .

(if the roots of the characteristic equation are complex please find the real-valued general solutions, i.e. the answer in the form  $C_1e^{rt} + C_2e^{\bar{r}t}$ , where  $r$  is complex, will not be accepted)).

- (b) For the case of item (a) i), i.e. when  $\alpha = \frac{5}{12}$ , solve the following problems:
  - find the solution satisfying the initial condition  $y(0) = -2, y'(0) = V$ , where  $V$  is a parameter;
  - find all parameters  $V$  such that the solution  $y(t)$  obtained in the previous subitem for such  $V$  satisfies:  $y(t) < 0$  for any  $t \geq 0$ .
- (c) For the case of item (a) iii), i.e. when  $\alpha = \frac{13}{12}$  solve the following problems:
  - Find the solution of the equation with the initial conditions  $y(\frac{\pi}{2}) = -4e^{\frac{3\pi}{4}}, y'(\frac{\pi}{2}) = -9e^{\frac{3\pi}{4}}$ . Describe the behavior of the solution as  $t \rightarrow -\infty$ ;
  - Determine  $\lambda, \mu > 0, R > 0$  and  $\delta \in [0, 2\pi)$  so that the solution obtained in the previous item can be written in the form  $e^{\lambda t}R \cos(\mu t - \delta)$  (you can use a calculator to determine an approximate value of  $\delta$ ). Then sketch the graph of this solution.

- Consider the equation  $y'' + (1 - \alpha)y' + (2\alpha - 3)(4 - 3\alpha)y = 0$ , where  $\alpha$  is a parameter.

- Determine the values of  $\alpha$ , if any, for which all solutions tend to zero as  $t \rightarrow +\infty$ .
- Determine the values of  $\alpha$ , if any, for which all nonzero solutions become unbounded as  $t \rightarrow +\infty$ .

Hint: It is more efficient here to use the Vieta theorem for roots of quadratic equation instead of the quadratic formula, but be careful on the signs.