

ON LOCAL GEOMETRY OF VECTOR DISTRIBUTIONS WITH GIVEN JACOBI SYMBOLS

BORIS DOUBROV AND IGOR ZELENKO

ABSTRACT. Based on the ideas of Optimal Control, we introduce the new basic characteristic of a bracket generating distribution, the Jacobi symbol. In contrast to the classical Tanaka symbol, the set of Jacobi symbols is discrete and classifiable. We give an explicit and unified algebraic procedure for the construction of the canonical frames (the structure of absolute parallelism) for all distribution with given Jacobi symbol. We describe all Jacobi symbols for which this procedure ends up in a finite number of steps (i.e. all Jacobi symbols of finite type) and distinguish the symplectically flat distributions with given Jacobi symbol, which in the case of finite type are the maximally symmetric distribution among all distribution with given Jacobi symbol. Also, for the most of Jacobi symbols we relate the prolongation procedure and the resulting prolongation algebra, which is the algebra of infinitesimal symmetries of the corresponding symplectically flat distribution, to the standard (in the sense of Kobayashi and Sternberg) prolongation of certain subspace in the algebra of infinitesimal symmetries of the flat (unparametrized) curve of flags associated with the Jacobi symbol. In particular, we show that as such subspace one can take the algebra of infinitesimal symmetries of the flat curve considered as parametrized curve (i.e. only symmetries preserving the parametrization are considered). Finally, we give an upper bounds for the size of the resulting prolongation algebra in terms of certain spaces of polynomials vanishing on certain projective varieties. For some class of Jacobi symbols we are able to identify the resulting prolongation algebra with such spaces of polynomials. In this way we get the description of the prolongation algebra for all Jacobi symbols of finite type appearing in rank 2 and rank 3 distributions and also for most of such Jacobi symbols appearing in rank 4 distributions. In particular, in the case of rank 3 distributions this description is given in terms of the tangential developable and the secant varieties of a rational normal curve.

1. INTRODUCTION

This paper is a culmination of a long-standing program for study of the geometry of non-holonomic vector distributions via a novel approach called the *symplectification/linearization procedure*. This approach takes its origin in Optimal Control and in essence consists of the reduction of the equivalence problem for distributions to the (extrinsic) differential geometry of curves of isotropic and coisotropic subspaces (or, shortly, of curves of symplectic flags) in a linear symplectic space. The extrinsic geometry of such curves is simpler in many respects than the original equivalence problem.

In the earlier articles [15, 16] and the preprint [18] we used this approach to construct the canonical frames (the canonical structures of absolute parallelism) for distributions of rank 2 and rank 3 on manifolds of arbitrarily large dimension under very mild genericity assumptions. Later on, in order to interpret these constructions in more conceptual way and generalize them to distributions of higher rank, we developed several new prolongation techniques such as Tanaka

2000 *Mathematics Subject Classification.* 58A30, 58A17, 53A33.

Key words and phrases. vector distributions, Tanaka prolongation, canonical frames, flag symbol, algebra of infinitesimal symmetries, representation of \mathfrak{sl}_2 .

Research of I. Zelenko is supported by NSF grant DMS 140619.

like theory for curves of flags [19, 20] and for flag structures on manifolds [21]. In the course of our work we realized that we can distinguish a new basic invariant of distributions, the *Jacobi symbol* [21], which is discrete and classifiable in contrast to the Tanaka symbol. In the present paper we combine all this together to describe the uniform (i.e. without any additional branching) construction of canonical frames for distribution with given Jacobi symbol. Our approach significantly extends the scope of the distributions for which the canonical frames can be constructed in explicit algebraic terms, avoiding the problems of the classification of Tanaka symbols (graded nilpotent Lie algebras) and the possible presence of moduli in the set of Tanaka symbols.

1.1. Equivalence problem for vector distributions: statement of the problem. A rank l vector distribution on an n -dimensional manifold M (or, shortly, an (l, n) -distribution) is a rank l subbundle of the tangent bundle TM . The group of germs of diffeomorphisms of M acts naturally on the set of germs of (l, n) -distributions and defines the equivalence relation there.

Vector distributions appear naturally in Geometric Control Theory as the sets of admissible velocities for control systems linear with respect to control parameters and Geometric Theory of Differential Equations as natural distributions on submanifolds of jet spaces.

A simple estimation shows that at least $l(n-l) - n$ functions of n variables are required to describe generic germs of (l, n) -distribution, up to the equivalence. There are only three cases, where $l(n-l) - n$ is not positive: $l = 1$ (line distributions), $l = n - 1$, and $(l, n) = (2, 4)$. Moreover, it is well known that in these cases generic germs of distributions are equivalent. For $l = 1$ it is just a particular case of Frobenius theorem. For $l = n - 1$ all generic germs are equivalent to Darboux's model, while for $(l, n) = (2, 4)$ they are equivalent to so-called Engel's model (see, for example, [6]). In all other cases generic (l, n) -distributions have functional, and, thus, non-trivial differential invariants. Often it is rather difficult to compute such invariants and to interpret them geometrically.

The general way to solve equivalence problems was developed by E. Cartan by mean of his method of "moving frames". The main goal of this method is to assign to a geometric structure on M the (co)frame or the structure of absolute parallelism on some (fiber) bundle $P(M)$ over M in a canonical way. First, it gives the sharp upper bounds for the group of local symmetries of geometric structures and the explicit description of the group of local symmetries for the maximally symmetric models. Second, it reduces the construction of full system of invariants for the original geometric structure to the construction of the full system of invariants for the absolute parallelism and the latter can be done easily [38].

For example, in his classical paper [9] Cartan associates a canonical coframe to any maximally nonholonomic $(2, 5)$ -distribution. This is the case of distributions of smallest rank in the manifold of smallest dimensions where generic distributions have functional invariants. Cartan's method of equivalence was successfully applied to other classes of low-dimensional distributions, for example, for $(3, 6)$ -distributions by R. Bryant ([4, 5]). However, in the original Cartan prolongation procedure one cannot predict the number of prolongation steps and the dimension of the bundle, where the canonical coframe lives, without making the concrete normalizations on each step. The latter often is accompanied with extremely tedious calculations.

Cartan's prolongation procedure has been reformulated by S. Sternberg in the framework of G -structures. However, geometric structures involving nonholonomic distributions do not properly fit into this framework, because the information about the nontrivial commutators of vector fields tangent to these distributions is not encoded in this framework.

1.2. Tanaka approach. N. Tanaka developed a deep generalization of the theory of G -structures which is perfectly adopted to nonholonomic structures, i.e. to nonholonomic distributions themselves or to nonholonomic distribution together with additional structures on them [32, 35, 3, 44]. Here we shortly describe the main notions and results of the Tanaka theory in context of distributions.

First, passing from the natural filtered structure generated by a distribution to the corresponding graded objects, Tanaka assigned to the distribution at a point a special graded nilpotent Lie algebra, called the *symbol of a distribution D at this point*, which contains the information about the principal parts of all commutators of vector fields tangent to D .

More precisely, taking Lie brackets of vector fields tangent to a distribution D (i.e. sections of D) one can define a filtration

$$(1.1) \quad D^{-1} \subset D^{-2} \subset \dots$$

of the tangent bundle, called a *weak derived flag* or a *small flag (of D)*. More precisely, set $D = D^{-1}$ and define recursively $D^{-j} = D^{-j+1} + [D, D^{-j+1}]$, $j > 1$. Let X_1, \dots, X_l be l vector fields constituting a *local basis* of a distribution D , i.e. $D = \text{span}\{X_1, \dots, X_l\}$ in some open set in M . Then $D^{-j}(x)$ is the linear span of all iterated Lie brackets of these vector fields, of length not greater than j , evaluated at a point x . A distribution D is called *bracket-generating* (or *completely nonholonomic*) if for any x there exists $\mu \in \mathbb{N}$ such that $D^{-\mu}(x) = T_x M$. The number μ is called the *degree of nonholonomy* of D at a point x . Let $\mathfrak{g}^{-1}(x) \stackrel{\text{def}}{=} D^{-1}(x)$ and $\mathfrak{g}^j(x) \stackrel{\text{def}}{=} D^j(x)/D^{j+1}(x)$ for $j < -1$. Consider the graded space

$$(1.2) \quad \mathfrak{t}(x) = \bigoplus_{j=-\mu}^{-1} \mathfrak{g}^j(x),$$

corresponding to the filtration

$$D(x) = D^{-1}(x) \subset D^{-2}(x) \subset \dots \subset D^{-\mu+1}(x) \subset D^{-\mu}(x) = T_x M.$$

This space is endowed naturally with the structure of a graded nilpotent Lie algebra, generated by $\mathfrak{g}^{-1}(x)$. Indeed, let $\mathfrak{p}_j : D^j(x) \mapsto \mathfrak{g}^j(x)$ be the canonical projection to a factor space. Take $Y_1 \in \mathfrak{g}^i(x)$ and $Y_2 \in \mathfrak{g}^j(x)$. To define the Lie bracket $[Y_1, Y_2]$ take a local section \tilde{Y}_1 of the distribution D^i and a local section \tilde{Y}_2 of the distribution D^j such that $\mathfrak{p}_i(\tilde{Y}_1(x)) = Y_1$ and $\mathfrak{p}_j(\tilde{Y}_2(x)) = Y_2$. It is clear that $[Y_1, Y_2] \in \mathfrak{g}^{i+j}(x)$. Put

$$(1.3) \quad [Y_1, Y_2] \stackrel{\text{def}}{=} \mathfrak{p}_{i+j}([\tilde{Y}_1, \tilde{Y}_2](x)).$$

It is easy to see that the right-hand side of (1.3) does not depend on the choice of sections \tilde{Y}_1 and \tilde{Y}_2 . Besides, $\mathfrak{g}^{-1}(x)$ generates the whole algebra $\mathfrak{t}(x)$. A graded Lie algebra satisfying the last property is called *fundamental*. The graded nilpotent Lie algebra $\mathfrak{m}(x)$ is called the (*Tanaka symbol of the distribution D at the point x*).

Given a fundamental graded nilpotent Lie algebra $\mathfrak{t} = \bigoplus_{j \leq -1} \mathfrak{g}^j$ Tanaka considered *distributions of constant type \mathfrak{t}* , i.e. distributions D with symbols at any point isomorphic to \mathfrak{t} . Note that in general the assumption of constancy of Tanaka symbol is quite restrictive, see discussions in subsection below. Among all such distributions he distinguish the most simple distribution called the *flat (or standard) distribution* of type \mathfrak{t} , namely, the left-invariant distribution on the connected simply-connected Lie group with Lie algebra \mathfrak{t} such that $D_{\mathfrak{t}}(e) = \mathfrak{g}^{-1}$, where e is the identity of the group.

Remarkably, the prolongation procedure for the construction of canonical frames can be described uniformly, i.e. without any additional branching, for all distribution of constant Tanaka type \mathfrak{t} . This construction can be imitated by the construction of such canonical frame for the flat distribution of type \mathfrak{t} . The latter construction can be described purely algebraically in terms of the so-called universal algebraic prolongation of the symbol \mathfrak{t} , which is in essence the maximal (nondegenerate) graded Lie algebra, containing the graded algebra \mathfrak{t} as its negative part.

More precisely, the *universal algebraic prolongation of the symbol \mathfrak{t}* is the graded Lie algebra $\mathfrak{u}(\mathfrak{t}) = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}^i(\mathfrak{t})$ determined by the following three conditions:

- (1) the graded subalgebra $\bigoplus_{i < 0} \mathfrak{g}^i(\mathfrak{t})$ of $\mathfrak{u}(\mathfrak{t})$, i.e. the negative part of $\mathfrak{u}(\mathfrak{t})$, coincides with \mathfrak{t} ;
- (2) (*non-degenericity assumption*) for any nonzero $x \in \mathfrak{g}^i(\mathfrak{t})$ with $i \geq 0$ there exists $y \in \mathfrak{t}$ such that $[x, y] \neq 0$ (i.e. $\text{ad } x|_{\mathfrak{m}} \neq 0$);
- (3) $\mathfrak{u}(\mathfrak{m})$ is the maximal graded algebra satisfying conditions (1) and (2) above.

Note that if $\dim \mathfrak{u}(\mathfrak{t}) < \infty$, then $\mathfrak{u}(\mathfrak{t})$ is isomorphic to the algebra of infinitesimal symmetries of the flat distribution of type \mathfrak{t} ; if $\dim \mathfrak{u}(\mathfrak{t}) = \infty$, then the completion of $\mathfrak{u}(\mathfrak{t})$ is isomorphic to the algebra of formal power series of infinitesimal symmetries of the flat distribution of \mathfrak{t} ([32] section 6, [42] section 2).

The universal algebraic prolongation can be explicitly realized by constructing $\mathfrak{g}^0(\mathfrak{t})$, $\mathfrak{g}^1(\mathfrak{t})$ etc inductively. Indeed, $\mathfrak{g}^0(\mathfrak{t})$ is realized as the Lie algebra of all derivations a of \mathfrak{t} , preserving the grading (i.e. such that $a\mathfrak{g}^i \subset \mathfrak{g}^i$ for all $i < 0$). Further, the space $\mathfrak{t} \oplus \mathfrak{g}^0(\mathfrak{t})$ is equipped with the structure of Lie algebra in the natural way and we can define $\mathfrak{g}^1(\mathfrak{t})$ as follows:

$$\mathfrak{g}^1(\mathfrak{t}) = \left\{ f \in \bigoplus_{i < 0} \text{Hom}(\mathfrak{g}^i(\mathfrak{t}), \mathfrak{g}^{i+1}(\mathfrak{t})) : f([v_1, v_2]) = [f(v_1), v_2] + [v_1, f(v_2)], \forall v_1, v_2 \in \mathfrak{t} \right\}$$

Now assume that spaces $\mathfrak{g}^l(\mathfrak{t}) \subset \bigoplus_{i < 0} \text{Hom}(\mathfrak{g}^i(\mathfrak{t}), \mathfrak{g}^{i+l}(\mathfrak{t}))$ are defined for all $0 < l < k$. Set

$$(1.4) \quad [f, v] = -[v, f] = f(v) \quad \forall f \in \mathfrak{g}^l(\mathfrak{t}), 0 \leq l < k, \text{ and } v \in \mathfrak{t}.$$

Then let

$$(1.5) \quad \mathfrak{g}^k(\mathfrak{t}) \stackrel{\text{def}}{=} \left\{ f \in \bigoplus_{i < 0} \text{Hom}(\mathfrak{g}^i(\mathfrak{t}), \mathfrak{g}^{i+k}(\mathfrak{t})) : f([v_1, v_2]) = [f(v_1), v_2] + [v_1, f(v_2)] \forall v_1, v_2 \in \mathfrak{t} \right\}.$$

The space $\bigoplus_{i \in \mathbb{Z}} \mathfrak{g}^i(\mathfrak{t})$ can be naturally endowed with the structure of a graded Lie algebra. The brackets of two elements from \mathfrak{t} are as in \mathfrak{t} . The brackets of an element with non-negative weight and an element from \mathfrak{t} are already defined by (1.4). It only remains to define the brackets $[f_1, f_2]$ for $f_1 \in \mathfrak{g}^k(\mathfrak{t})$, $f_2 \in \mathfrak{g}^l(\mathfrak{t})$ with $k, l \geq 0$. The definition is inductive with respect to k and l : if $k = l = 0$ then the bracket $[f_1, f_2]$ is as in $\mathfrak{g}^0(\mathfrak{t})$. Assume that $[f_1, f_2]$ is defined for all $f_1 \in \mathfrak{g}^k(\mathfrak{t})$, $f_2 \in \mathfrak{g}^l(\mathfrak{t})$ such that a pair (k, l) belongs to the set

$$\{(k, l) : 0 \leq k \leq \bar{k}, 0 \leq l \leq \bar{l}\} \setminus \{(\bar{k}, \bar{l})\}.$$

Then define $[f_1, f_2]$ for $f_1 \in \mathfrak{g}^{\bar{k}}(\mathfrak{t})$, $f_2 \in \mathfrak{g}^{\bar{l}}(\mathfrak{t})$ to be the element of $\bigoplus_{i < 0} \text{Hom}(\mathfrak{g}^i(\mathfrak{t}), \mathfrak{g}^{i+\bar{k}+\bar{l}}(\mathfrak{t}))$ given

by

$$(1.6) \quad [f_1, f_2]v \stackrel{\text{def}}{=} [f_1(v), f_2] + [f_1, f_2(v)] \quad \forall v \in \mathfrak{t}.$$

It is easy to see that $[f_1, f_2] \in \mathfrak{g}^{k+l}(\mathfrak{t})$ and that $\bigoplus_{i \in \mathbb{Z}} \mathfrak{g}^i$ with bracket product defined as above is a graded Lie algebra.

One of the main results of Tanaka theory [32] applied to the geometry of distributions can be formulated as follows:

Theorem 1.1. *Assume that $\dim \mathfrak{u}(\mathfrak{t}) < \infty$. Then to a distribution D with constant symbol \mathfrak{t} one can assign in a canonical way a bundle over M of dimension equal to $\dim \mathfrak{u}(\mathfrak{t})$ equipped with a canonical frame. In particular the dimension of algebra of infinitesimal symmetries of D is not greater than $\dim \mathfrak{u}(\mathfrak{t})$. Moreover, this upper bound is sharp and is achieved if and only if a distribution is locally equivalent to the flat distribution $D_{\mathfrak{t}}$.*

Thus, the Tanaka approach allows one to predict the number of prolongations steps and the dimension of the bundle, where the canonical frame lives, without making concrete normalization on each step as the original Cartan method of equivalence suggests. For this it is enough to calculate the universal algebraic prolongation of the symbol \mathfrak{t} , or, equivalently, all algebraic prolongations $\mathfrak{g}^i(\mathfrak{t})$ up to the first vanishing one (if it exists). According to (1.5) the latter consists of solving systems of linear algebraic equations.

Remark 1.1. More generally, given any subalgebra \mathfrak{g}^0 of the algebra $\mathfrak{g}^0(\mathfrak{t})$ one can define the *universal algebraic prolongation*

$$(1.7) \quad \mathfrak{u}(\mathfrak{t}, \mathfrak{g}^0) = \bigoplus_{i \in \mathbb{Z}} \mathfrak{u}^i(\mathfrak{t}, \mathfrak{g}^0)$$

of the pair $(\mathfrak{t}, \mathfrak{g}^0)$ as the maximal (nondegenerate) graded Lie algebra, containing the graded algebra $\mathfrak{t} \oplus \mathfrak{g}^0$ as its non-positive part. This algebra can be constructed by the inductive procedure described by formulas analogous to (1.5): for this set $\mathfrak{u}^i(\mathfrak{t}, \mathfrak{g}^0) = \mathfrak{g}^i$ for $i \leq 0$ and recursively

$$(1.8) \quad \mathfrak{u}^i(\mathfrak{t}, \mathfrak{g}^0) \stackrel{\text{def}}{=} \left\{ f \in \bigoplus_{j < 0} \text{Hom}(\mathfrak{u}^j(\mathfrak{t}, \mathfrak{g}^0), \mathfrak{u}^{j+i}(\mathfrak{t}, \mathfrak{g}^0)) : f([v_1, v_2]) = [f(v_1), v_2] + [v_1, f(v_2)] \quad \forall v_1, v_2 \in \mathfrak{t} \right\}$$

for $i > 0$. The space $\mathfrak{u}^i(\mathfrak{t}, \mathfrak{g}^0)$ is called the *i th Tanaka algebraic prolongation of the pair $(\mathfrak{t}, \mathfrak{g}^0)$* .

The pairs $(\mathfrak{t}, \mathfrak{g}^0)$ appear as basic characteristics of distributions with constant Tanaka symbol \mathfrak{t} and with certain additional structures on them. Such structures are called *Tanaka structures of type $(\mathfrak{t}, \mathfrak{g}^0)$* . To define these structures first to a distribution of type \mathfrak{t} one assigns the principal bundle $P^0(\mathfrak{t})$ of partial frames with the structure group $\text{Aut}(\mathfrak{t})$ of automorphisms of the graded Lie algebra \mathfrak{t} : $P^0(\mathfrak{t})$ is the set of all pairs (x, φ) , where $x \in M$ and $\varphi : \mathfrak{t} \rightarrow \mathfrak{t}(x)$ is an isomorphism of the graded Lie algebras \mathfrak{t} and $\mathfrak{t}(x)$. Since \mathfrak{g}^{-1} generates \mathfrak{t} , the group $\text{Aut}(\mathfrak{t})$ can be identified with a subgroup of $\text{GL}(\mathfrak{g}^{-1})$. By the same reason a point $(x, \varphi) \in P^0(\mathfrak{t})$ of a fiber of $P^0(\mathfrak{t})$ is uniquely defined by $\varphi|_{\mathfrak{g}^{-1}}$. So one can identify $P^0(\mathfrak{t})$ with the set of pairs (x, ψ) , where $x \in M$ and $\psi : \mathfrak{g}^{-1} \rightarrow D(x)$ i.e. it is a partial frame that can be extended to an automorphism of the graded Lie algebras \mathfrak{t} and $\mathfrak{t}(x)$. Let $G^0 \subset \text{Aut}(\mathfrak{t})$ be a Lie group with the Lie algebra \mathfrak{g}^0 . A *Tanaka structure of constant type $(\mathfrak{t}, \mathfrak{g}^0)$* is any G^0 -reduction of the bundle $P^0(\mathfrak{t})$. In [32] Theorem 1.1 (except the last statement about the uniqueness of the maximally symmetric model) was proved not only for distributions but for any Tanaka structure of type $(\mathfrak{t}, \mathfrak{g}^0)$. The last statement of Theorem 1.1 is not true for general Tanaka structures (for example it is not true for the classical Riemannian metrics which is a particular case of them) but it is true for distributions (i.e. when $\mathfrak{g}^0 = \mathfrak{g}^0(\mathfrak{t})$) due to the presence of the grading element in $\mathfrak{g}^0(\mathfrak{t})$.

Remark 1.2. Further, Tanaka showed [33, 35, 7, 42] that if the algebra $\mathfrak{u}(\mathfrak{t}, \mathfrak{g}^0)$ is semisimple, then the normalization conditions in the prolongation procedure can be chosen such that the resulting coframe is the Cartan connection (a coframe with special nice properties) modeled by a parabolic geometry. The E. Cartan and R. Bryant constructions for (2, 5) and (3, 6) distribution are the particular cases of this theory, which includes also conformal and CR structures [39, 8].

1.3. Limits and weak points of Tanaka approach. All constructions of the Tanaka theory depend on the symbol. Therefore in order to apply the Tanaka machinery to all bracket-generating distributions of the given rank l on a manifold of the given dimension n , one has to classify all n -dimensional graded nilpotent Lie algebra with l generators. Note also that the set of all possible Tanaka symbols may contain moduli, i.e. may depend on continuous parameters.

The problem of classification of all Tanaka symbols is quite nontrivial already in dimension 7 (see [30]) and it looks completely hopeless for arbitrary dimensions. For example, as was shown in [30] already in dimension 7 the continuous parameters appear in symbols of rank (3, 6, ...) -distributions (see models $m7_3_3(\alpha)$ and $m7_3_13(\alpha)$ there), and there are 6 more non-isomorphic symbols in addition to that. In the case of (2, n)-distributions with $\dim D^{-3} = 5$ there are 3 non-isomorphic symbols Tanaka symbols if $n = 6$, then 8 non-isomorphic symbols if $n = 7$, and continuous parameters appear in the set of all symbols for $n \geq 8$.

Hence, first, generic distributions may have non-isomorphic Tanaka symbols at different points so that in this case the Tanaka theory cannot be directly applied, although a generalization of the Tanaka prolongation procedure to distributions with non-constant Tanaka symbols is certainly possible (a slightly different aspect concerning the upper bound for the infinitesimal symmetries algebra of distribution with non-constant Tanaka symbols was addressed in [28, 29]). Second, even restricting to distributions with a constant Tanaka symbol, without the classification of these symbols we do not have a complete picture about the geometry of distributions. In particular, even in the case of rank 2 and rank 3 distributions the Tanaka theory gives a complete picture only in few cases of low dimensions.

1.4. Geometry of distribution via symplectification. In [1, 2] Andrei Agrachev put forward the idea that invariants of a submanifold of the tangent bundle of a manifold M (with respect to the natural action of the group of diffeomorphisms of M) can be obtained by studying the flow of extremals of variational problems naturally associated with this submanifold. In the case when this submanifold is a bracket-generating distribution, the Agrachev idea can be briefly described as follows (see section 2 below for more detail).

First recall that a Lipschitzian curve $\alpha(t)$ is called *admissible (horizontal)* curve of a distribution D , if $\alpha'(t) \in \mathcal{D} \cap T_{\alpha(t)}M$ for almost every t . Fixing any two points on M consider any functional of integral type (for example, the length functional with respect to some Riemannian metric) on the space of admissible curves connecting these points. Note that this space is not empty by the Rashevsky-Chow theorem. The Pontryagin extremals of a variational problem are special curves in the cotangent bundle T^*M such that their projections to the base manifold M are candidates for an extremum. They are described by the Pontryagin Maximum Principle of Optimal Control [36]. Among all this extremals there can be extremals with zero Lagrangian multiplier near the functional. Such extremals are called *abnormal*.

Abnormal extremals do not depend on the functional but on the distribution D only and can be described purely geometrically in terms of D without referring to any auxiliary functional on the space of admissible curves of D and using just the canonical symplectic structure on the cotangent bundle T^*M .

For this consider the dual object to the distribution D in T^*M , which is its annihilator D^\perp , i.e. the submanifold of T^*M consisting of all $(p, q) \in T^*M$, where $q \in M$ and $p \in T_q^*M$, such that the covector p annihilates the space $D(q)$. Also let $(TM)^\perp$ be the zero section of T^*M .

An abnormal extremal is a Lipschitzian curve in $D^\perp \setminus (TM)^\perp$ such that the tangent line to it at almost every point belongs to the kernel of the restriction of the canonical symplectic form $\widehat{\sigma}$ of T^*M to D^\perp at this point. Hence, abnormal extremals lie in the *degeneracy locus* \widetilde{W}_D of the form $\widehat{\sigma}|_{D^\perp}$, i.e. in the set of all points, where the form $\widehat{\sigma}|_{D^\perp}$ is degenerate.

It turns out (see section 2) that if $\text{rank } D$ is odd then $\widetilde{W}_D = D^\perp$ and if $\text{rank } D$ is even then for any point $q \in M$ the intersection $\widetilde{W}_D(q)$ of W_D with the fiber $D^\perp(q)$ over q is the zero level set of a homogeneous polynomial of degree $\frac{1}{2} \text{rank } D$, which is the Pfaffian of the so-called Goh matrix. In particular, in the case of rank 2 distributions $\widetilde{W}_D(q)$ is a vector subspace of $D^\perp(q)$. Moreover, in this case \widetilde{W}_D is equal to the annihilator of D^{-2} .

As a consequence of the previous paragraph, for all odd rank distributions D the set \widetilde{W}_D is an odd dimensional submanifold of T^*M and for even rank distributions satisfying very mild genericity assumptions an open and dense subset of \widetilde{W}_D is a smooth submanifold of odd dimension. In particular, in the case of $(2, n)$ -distributions with $n \geq 3$ bracket genericity assumption can be taken as such genericity assumption. Thus in all these cases the form $\widehat{\sigma}|_{\widetilde{W}_D}$ is degenerate at the point of smoothness of \widetilde{W}_D .

Further, consider the “regular part” W_D of the degeneracy locus \widetilde{W}_D consisting of all points of smoothness λ in \widetilde{W}_D such that the kernel of the form $\widehat{\sigma}(\lambda)|_{\widetilde{W}_D}$ is one dimensional. These kernels define the characteristic rank 1 distributions \widehat{C} on W_D and any integral curve of this characteristic distribution is an abnormal extremal. Such abnormal extremals are called *regular*.

The set W_D and the characteristic rank 1 distribution on it can be described explicitly in terms of the weak derived flag (1.1) and a local basis of D and the set W_D is non-empty (and therefore open in \widetilde{W}_D) for generic germs of distributions (see section 2 for details). In particular, $W_D = (D^{-2})^\perp \setminus (D^{-3})^\perp$ if $\text{rank } D = 2$ and $W_D = (D)^\perp \setminus (D^{-2})^\perp$ if $\text{rank } D = 3$ and therefore W_D is non-empty for bracket generating distributions of rank 2 and 3 in an ambient manifold M of dimension greater or equal than 4.

After the projectivization of each fiber of the cotangent bundle T^*M regular abnormal extremals foliate the even-dimensional submanifold $\mathbb{P}W_D$ of the projectivized cotangent bundle $\mathbb{P}T^*M$. The space of regular abnormal extremals can be considered as the quotient manifold \mathcal{N} by the foliation of these extremals. The natural contact distribution Δ is induced on \mathcal{N} by the tautological 1-form (the Liouville form) in T^*M . The natural conformal symplectic structure, i.e. a symplectic form σ_γ is defined, up to a nonzero scalar multiple, on each space $\Delta(\gamma)$, $\gamma \in \mathcal{N}$.

Further, the distributions D can be lifted to W_D by the pullback with respect to the canonical projection $W_D \rightarrow M$. Denote the resulting distribution by \mathcal{J} . Given an extremal $\gamma \in \mathcal{N}$, for any point $\lambda \in \gamma$ consider the images $\mathfrak{J}_\gamma(\lambda)$ of the space $\mathcal{J}(\lambda)$ under the differentials of the canonical map from W_D to the quotient manifold \mathcal{N} . These images define a curve $\lambda \mapsto \mathfrak{J}_\gamma(\lambda)$, $\lambda \in \gamma$ of subspaces in the space $\Delta(\gamma)$, called the *Jacobi curve* of γ .

The motivation for this name comes from the considering the case of rank 2 distributions. In this case all subspaces $\mathfrak{J}_\gamma(\lambda)$ are Lagrangian subspaces of $\Delta(\gamma)$ with respect to the symplectic form σ_γ , i.e. the curve \mathfrak{J}_γ is a curve in a Lagrangian Grassmannian. Actually this curve describes the space of solutions of the Jacobi equations along the extremal γ and the index of the second variation along the extremal γ can be expressed in terms of certain intersection index (a Maslov

type index) of it [1]. In the case of distributions of the rank greater than 2 all subspaces $\mathfrak{J}_\gamma(\lambda)$ are strictly coisotropic, i.e. they strictly contain the skew-orthogonal complements of themselves with respect to the form σ_γ . Although in this case the index of the second variation along an abnormal extremal is generically infinite, so that there is no analogy with the Calculus of Variation as in the case of rank 2 distributions, we still keep to call the curves \mathfrak{J}_γ by the name Jacobi curve and we can exploit them for the local geometry of distributions. Namely, *one can construct canonical frames and invariants of the original distribution D by studying the differential geometry of the Jacobi curves \mathfrak{J}_γ with respect to the action the group $\text{CSp}(\Delta(\gamma))$ of the conformal symplectic transformation of $\Delta(\gamma)$ (i.e. the group of the linear transformation preserving the symplectic form σ_γ , up to a scalar multiple) for generic abnormal extremals γ .*

1.5. Jacobi symbol of distributions. It turns out [19, 20] that the differential geometry of Jacobi curves (and even more general curves in flag varieties) can be treated similarly to the Tanaka theory of filtered structures. The central notion in this treatment is the flag symbol of the Jacobi curve at a given point, which in turn gives a new basic characteristic of distributions, called the Jacobi symbol. Here we give a brief informal description of what is the Jacobi symbols of a distribution, postponing the detailed description to section 5.

First we build a curve of flags of $\Delta(\gamma)$ out of the Jacobi curve \mathfrak{J}_γ by collecting the osculating spaces of the Jacobi curve of any order together with their skew-orthogonal complements with respect to the form σ_γ . The resulting curve of flags is called the *extended Jacobi curve of the extremal γ* and will be denoted by $\widehat{\mathfrak{J}}_\gamma$. Each flag $\widehat{\mathfrak{J}}_\gamma(\lambda)$ consists of isotropic/coisotropic subspaces of $\Delta(\gamma)$ and is stable with respect to the operation of taking the skew-orthogonal complement. Such flags will be called *symplectic flags* for shortness. Note that in this case the graded space $\text{gr } \widehat{\mathfrak{J}}_\gamma(\lambda)$ corresponding to the filtration $\widehat{\mathfrak{J}}_\gamma(\lambda)$ is endowed with the conformal symplectic structure naturally induced by the conformal symplectic structure on $\Delta(\gamma)$.

Moreover the extended Jacobi curve $\widehat{\mathfrak{J}}_\gamma$ satisfies an important additional property. Let us describe it first for the case of distributions of odd rank. If the subspaces of the flags $\widehat{\mathfrak{J}}_\gamma(\lambda)$ are indexed by \mathbb{Z} such that the corresponding filtration is nonincreasing by inclusion, then we have

Osculation property *For any $i \in \mathbb{Z}$ and any $\lambda \in \gamma$ the osculating space at λ of the curve of subspaces of $\widehat{\mathfrak{J}}_\gamma$ indexed by i is contained in the subspaces of $\widehat{\mathfrak{J}}_\gamma(\lambda)$ indexed by $i - 1$.*

The latter implies that the tangent line to the curve $\widehat{\mathfrak{J}}_\gamma$ can be identified with a line of degree -1 endomorphisms of the graded space $\text{gr } \widehat{\mathfrak{J}}_\gamma(\lambda)$, belonging to the Lie algebra of the group of conformal symplectic transformations. This line, up to a conjugation by a conformal symplectic transformation preserving the grading, is called the *flag symbol* or the *Jacobi symbol* of the curve $\widehat{\mathfrak{J}}_\gamma$ (or the curve \mathfrak{J}_γ)

Extended Jacobi curves of rank 2 distributions satisfy the osculation property as well. It is not the case for distributions of even rank greater than 2 if the subspaces of the flags $\widehat{\mathfrak{J}}_\gamma(\lambda)$ are indexed by \mathbb{Z} . However, the osculation property becomes valid also in this case, if the subspaces of the flags $\widehat{\mathfrak{J}}_\gamma(\lambda)$ are indexed by $\frac{1}{2}\mathbb{Z} = \{\frac{i}{2} : i \in \mathbb{Z}\}$ such that the corresponding filtration is nonincreasing by inclusion. So, the flag symbol as a line of degree -1 endomorphisms of the corresponding graded space $\text{gr } \widehat{\mathfrak{J}}_\gamma(\lambda)$ can be defined for even rank distribution as well.

To make things uniform it is more convenient in the case of rank 2 distributions to index the subspace of the extended Jacobi curve $\widehat{\mathfrak{J}}_\gamma$ by $\frac{1}{2}\mathbb{Z}_{\text{odd}} := \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$ (see subsection 4.1 below). Then the flag symbols of extended Jacobi curves of distributions of even rank greater than 2 are actually the direct sum of a flag symbol of curves of flags appearing as Jacobi curves of a distribution of

odd rank and one flag symbol of curves of flags appearing as Jacobi curves of a rank 2 distribution. This choice of the set of indices is also justified by the theory of irreducible representations of \mathfrak{sl}_2 , which plays an important role in the sequel (sections 6-8) : the spectrum of a semisimple element of such representations constitute of an arithmetic progression centered at 0 and in the case when the difference between consecutive terms of this progression is equal to 1 and the dimension of the corresponding \mathfrak{sl}_2 -module is even this spectrum belongs to $\frac{1}{2}\mathbb{Z}_{\text{odd}}$.

In contrast to the Tanaka symbols of distributions the flag symbols of Jacobi curves can be easily classified [20, subsection 7.2] and subsection 4.2 below. Besides, the set of all possible flag symbols is discrete. Therefore, if for any $\lambda \in W_D$ we denote by $\delta(\lambda)$ the flag symbol at λ of the Jacobi curve of the regular abnormal extremal passing through λ , then the map $\lambda \mapsto \delta(\lambda)$ is constant in a neighborhood of a generic point of the bundle W_D . Moreover, as it will be shown in subsection 4.3 for a generic point $q \in M$ there is a neighborhood U of q in M such that the map $\lambda \mapsto \delta(\lambda)$ is constant on the generic subset V of $\tilde{\pi}^{-1}(U) \cap \mathbb{P}W_D$, where $\tilde{\pi} : \mathbb{P}T^*M \rightarrow M$ is the canonical projection. In this case $\delta(\lambda)$, $\lambda \in V$ is called the *Jacobi symbol of the distribution D at the point q* . Jacobi symbol is a new basic characteristic of a distribution D .

1.6. Main results and organization of the paper. Although the definition of the Jacobi symbol is more involved, this is a simpler algebraic object than the Tanaka symbol. The natural task is *to describe the construction of the canonical frames uniformly for all distributions with given Jacobi symbol δ* .

To begin with this task, in subsection 4.4, given any Jacobi symbol δ , we define in a sense the most simple model of a distribution with Jacobi symbol δ . We call it symplectically flat distributions because it plays the same role in the local geometry of distributions with given Jacobi symbol, as the flat distribution of fixed type in the local geometry of distributions with given Tanaka symbol (see the last sentence of Theorem 5.2 below), i.e. it is the maximally symmetric distribution within the considered class (if one assumes the finiteness of type). In the case of distributions of odd rank or rank 2 the symplectically flat distributions are also flat distribution in the Tanaka sense for some fundamental Tanaka symbol. An interesting phenomenon here is that this is not the case for a symplectically flat distribution of even rank greater than 2. However, it can be considered as the flat distribution in the Tanaka sense for some nonfundamental Tanaka symbol (see Example 4.3 below).

The previous constructions assign to any distribution with given Jacobi symbol the following structure on the generic subset of the space of regular abnormal extremals \mathcal{N} : the contact distribution Δ and the curves $\widehat{\mathfrak{J}}_\gamma$ of symplectic flags on each fiber $\Delta(\gamma)$. In the case of distributions of odd rank and rank 2 this is a particular case of the flag structures (with constant flag symbol δ) introduced and studied in our recent preprint [21], while the flag structure in the case of distributions of even rank greater than 2 can be treated similarly. The construction of the canonical frames for distributions with given Jacobi symbol δ is given in section 5, Theorem 5.2. Instead of trying to adopt a prolongation procedure for the construction of a canonical frame directly to an original distribution, as in the classical approaches, we adopt it to the corresponding flag structure. This can be done in terms of natural algebraic operations on the Jacobi symbol δ in the category of graded Lie algebras, using the Tanaka like theory of curves of symplectic flags developed in [19, 20] and the Tanaka like prolongation of flag structures developed in [21]. Theorem 5.2 plays the same role in the local geometry of distributions with given Jacobi symbol as Theorem 1.1 in the Tanaka theory of distributions.

In section 6 we describe all Jacobi symbols for which the prolongation procedure ends up in a finite number of steps, i.e. all Jacobi symbols of finite type (see Theorems 6.1 and 6.2 there). Our

considerations there are based on two general criteria for finiteness of type, the Spencer criterium (see Theorem 6.4 and [37, 23, 31]) and the Tanaka criterium (see Theorem 6.3 and [32]), and some theory of \mathfrak{sl}_2 representations.

Sections 7 and 8 are devoted to more detailed study of the resulting prolongation algebra, which is also isomorphic to the algebra of infinitesimal symmetries of the corresponding symplectically flat distribution with given Jacobi symbol δ . This resulting algebra is closely related to the algebra of infinitesimal symmetries of the flat curve of flags with flag symbol δ , see Definition 5.1 (actually the latter algebra is equal to the degree 0 component in the resulting prolongation algebra). Theorem 7.2 and Corollary 7.1 reduce the calculation of the prolongation algebra to the calculation of the standard prolongation (as described, for example, in [26, 38]) of certain subspace in the algebra of infinitesimal symmetries of the flat (unparametrized) curve of flags associated with the Jacobi symbol. In particular, we show that as such subspace one can take the algebra of infinitesimal symmetries of the flat curve with flag symbol δ considered as parametrized curve (i.e. only symmetries preserving the parametrization are considered).

Finally in section 8 we make an attempt to describe the resulting prolongation algebra more explicitly using the fact that if a subspace in the symplectic algebra can be identified with a space of quadratic polynomials vanishing on a given projective variety (in the projectivization of the ambient linear symplectic space), then the standard prolongations of this subspace can be described as the space of certain polynomials vanishing on the secant variety of certain order of this projective variety. For general Jacobi symbols in this way we are able to show that the resulting prolongation algebra is a subspace in a space of this kind, obtaining an upper bound for the dimension of this resulting prolongation algebra (see Theorem 8.1) and an alternative proof of finiteness of type in the considered case, which is more constructive than the proof of Theorem 7.2. Also, we show (Theorem 8.2) that for a large class of Jacobi symbols the resulting prolongation algebra can be identified with some spaces of polynomials of this kind (and not only with some subspaces of them). In this way we get the description of the prolongation algebra for all Jacobi symbols of finite type appearing in rank 2 and rank 3 distributions and also for most of such Jacobi symbols appearing in rank 4 distributions.

For $(2, n)$ -distributions there was already an explicit description (see Theorem 7.1), based on the classical work of Cartan [9] for $n = 5$ and our previous works [15, 16] for $n > 5$. For rank 3 distributions the Jacobi symbol is described by a skew Young diagram consisting of 2 rows. If this skew Young diagram is nonrectangular (and the Jacobi symbol is of finite type) the prolongation algebra can be described in terms of the tangential developable and the secant varieties of a rational normal curve (Theorem 8.3), while in the case of rectangular Young diagram it is described explicitly by Theorem 8.4. For applicability of Theorem 8.2 in the case of rank 4 distribution see Remark 8.3, item 3 there.

2. ABNORMAL EXTREMALS OF DISTRIBUTIONS

Let D be a distribution on a manifold M . The crucial notion in the symplectification procedure of distributions is the notion of *an abnormal extremal*. To describe them, let $D^{-1} \subset D^{-2} \subset \dots$ be a *weak derived flag of D* , as defined in the beginning of section 1.2. Now denote by $(D^j)^\perp \subset T^*M$ the annihilator of D^j , namely

$$(D^j)^\perp = \{(p, q) \in T_q^*M \mid p \cdot v = 0 \quad \forall v \in D^j(q)\}.$$

Let $\pi: T^*M \mapsto M$ be the canonical projection. For any $\lambda \in T^*M$, $\lambda = (p, q)$, $q \in M$, $p \in T_q^*M$, let $\varsigma(\lambda)(\cdot) = p(\pi_*\cdot)$ be the canonical Liouville form and $\tilde{\sigma} = d\varsigma$ be the standard symplectic structure on T^*M . Also denote by $(TM)^\perp$ the zero section of T^*M .

Definition 2.1. An unparametrized Lipschitzian curve in $D^\perp \setminus (TM)^\perp$ is called an abnormal extremal of a distribution D if the tangent line to it at almost every point belongs to the kernel of the restriction $\hat{\sigma}|_{D^\perp}$ of $\hat{\sigma}$ to D^\perp at this point.

The term abnormal extremals comes from Optimal Control Theory: abnormal extremals of the distribution D are exactly Pontryagin extremals with zero Lagrange multiplier near the functional for any variational problem with constraints, given by the distribution D . We are interested not in all abnormal extremals but in those satisfying certain generic properties. Their description given below is based on elementary properties of antisymmetric matrices and their Pfaffians. Similar description is given in [10] (abnormal extremals of our interest here correspond to the lifts of singular curves of *minimal order* in the terminology of [10]).

Directly from the definition it follows that any abnormal extremal belongs to the set

$$(2.1) \quad \widetilde{W}_D := \{\lambda \in D^\perp : \ker(\hat{\sigma}|_{D^\perp}(\lambda)) \neq 0\}$$

Since $\text{codim} D^\perp = \text{rank } D$ and a skew-symmetric form in an odd dimensional vector space has a nontrivial kernel, one immediately have the following

Proposition 2.1. If $\text{rank } D$ is odd, then $\widetilde{W}_D = D^\perp$.

In order to understand the set \widetilde{W}_D in the case of distributions of even rank, let us introduce some notation from the Hamiltonian Dynamics. Given a function $h : T^*M \mapsto \mathbb{R}$ denote by \vec{h} the corresponding Hamiltonian vector field defined by the relation $i_{\vec{h}}\hat{\sigma} = -dh$. Further, given a vector field X in M define the corresponding ‘‘quasi-impulse’’ $H_X : T^*M \rightarrow \mathbb{R}$, $H_X(p, q) = p(X(q))$, $q \in M$, $p \in T_q^*M$. The corresponding Hamiltonian vector field \vec{H}_X is called the *Hamiltonian lift of X to T^*M* (note that by the definition $\pi_*(\vec{H}_X) = X$, where $\pi : T^*M \rightarrow M$ is the canonical projection). Note that the following identities hold for any vector fields X and Y on M :

$$(2.2) \quad dH_Y(\vec{H}_X) = H_{[X, Y]}$$

(as a matter of fact $dH_Y(\vec{H}_X)$ is exactly the Poisson brackets $\{H_X, H_Y\}$ of the Hamiltonians H_X and H_Y).

Now let (X_1, \dots, X_l) be a local basis of the distribution D , $D(q) = \text{span}\{X_1(q), \dots, X_l(q)\}$ for any q from some open set of M . Then locally

$$(2.3) \quad D^\perp = \{H_{X_1} = H_{X_2} = \dots = H_{X_l} = 0\}.$$

Further, for any $\lambda \in D^\perp$ define the following $l \times l$ antisymmetric matrix

$$(2.4) \quad G(\lambda) = (H_{[X_i, X_j]}(\lambda))_{i, j=1}^l.$$

This matrix is called the *Goh matrix (at the point λ) of the local basis (X_1, \dots, X_l) of the distribution D* .

Using formula (2.2) and Definition 2.1 it is easy to show that locally a curve $\gamma(\cdot) \subset D^\perp$ is an abnormal extremal if and only if

$$(2.5) \quad \gamma'(t) = \sum_{i=1}^l \alpha_i \vec{H}_{X_i}(\gamma(t)) \quad \text{for almost every } t,$$

where the vector $(\alpha_1, \dots, \alpha_l)^T$ belongs to the kernel of the Goh matrix $G(\gamma(t))$,

$$(2.6) \quad (\alpha_1, \dots, \alpha_l)^T \in \ker G(\gamma(t)), \quad \text{for almost every } t.$$

It is convenient to describe the degeneracy conditions and the kernel for a skew-symmetric matrix using the notion of Pfaffian. Recall that the Pfaffian $\text{pf}(A)$ of an $l \times l$ skew-symmetric matrix $A = (a_{ij})_{i,j=1}^l$ is by definition the square root of its determinant. It is equal to zero if l is odd and to a certain homogeneous polynomial of degree k in the entries of A if l is even, where $k = \frac{1}{2}l$. To describe this polynomial let $\omega = \sum_{0 \leq i < j \leq l} a_{ij} dx^i \wedge dx^j$ be the corresponding 2-form on \mathbb{R}^l taken with standard coordinates (x^1, \dots, x^l) . Then

$$(2.7) \quad \omega^k = k! \text{pf}(A) dx^1 \wedge \dots \wedge dx^l,$$

where ω^k is the k th wedge power of ω .

As a consequence of (2.5) and (2.6), we have the following

Proposition 2.2. *If rank D is even, the set \widetilde{W}_D , defined by (2.1), is given locally by the following formula*

$$(2.8) \quad \widetilde{W}_D = \{\lambda \in D^\perp : \text{pf}(G(\lambda)) = 0\}.$$

For example, if D is a rank 2 distributions

$$(2.9) \quad G(\lambda) = \begin{pmatrix} 0 & H_{[X_1, X_2]}(\lambda) \\ -H_{[X_1, X_2]}(\lambda) & 0 \end{pmatrix}, \quad \text{pf}(G(\lambda)) = H_{[X_1, X_2]}(\lambda).$$

Hence

$$(2.10) \quad \text{if } l = 2, \text{ then } \widetilde{W}_D = (D^{-2})^\perp,$$

because Proposition 2.2 implies locally that

$$\widetilde{W}_D = \{H_{X_1} = H_{X_2} = H_{[X_1, X_2]} = 0\} = (D^{-2})^\perp.$$

Since we originally work over \mathbb{R} , for l divisible by 4 the set \widetilde{W}_D might be empty (such distributions are called *fat* ([34])). However, in this case we can complexify the fibers of T^*M . Denote the obtained complex vector bundle (over real manifold M) by $(T^*M)^\mathbb{C}$. In the same way we define the complexified bundles $((D^j)^\perp)^\mathbb{C}$ and the Goh matrix $G(\lambda)$ for any $\lambda \in (D^\perp)^\mathbb{C}$. So, the set $(\widetilde{W}_D)^\mathbb{C} = \{\lambda \in (D^\perp)^\mathbb{C} : \text{pf}(G(\lambda)) = 0\}$ is not empty. In the sequel in case we need to work with complexified objects we will do it without special mentioning and use the same notation as in the real category.

Now define a subset W_D of \widetilde{W}_D as follows:

$$(2.11) \quad W_D := \{\lambda \in \widetilde{W}_D : \ker(\widehat{\sigma}|_{\widetilde{W}_D}(\lambda)) \text{ is one-dimensional}\}.$$

By constructions, the kernels of $\widehat{\sigma}|_{W_D}$ form the *characteristic rank 1 distribution* $\widehat{\mathcal{C}}$ on W_D and the integral curves of this distribution are abnormal extremals of the distribution D . Actually, they are exactly the abnormal extremals which lie entirely in W_D . Such abnormal extremals are called *regular*. Note that in general an abnormal extremal does not need to belong to W_D , however, for our purposes it is enough to work with regular abnormal extremals only.

Now describe the set W_D and the characteristic rank 1 distribution $\widehat{\mathcal{C}}$ in terms of a local basis (X_1, \dots, X_l) of the distribution D . For this recall several facts on kernels of skew-symmetric matrices using the notion of Pfaffian. Let $A = (a_{ij})_{i,j=1}^l$ be an $l \times l$ skew-symmetric matrix. Let A_i be the Pfaffian of the skew-symmetric matrix obtained from A by removing the i th row and the i th column. Further, for $i < j$ let A_{ij} be the Pfaffian of the skew-symmetric matrix obtained

from A by removing the i th and j th row and the i th and j th column. If A was a 2×2 matrix, so that after such removal we get an empty matrix, we set $A_{12} = 1$. To define A_{ji} for any ordered pair (i, j) (with integer i and j running from 1 to l) set

$$(2.12) \quad A_{ji} = -A_{ij}, \quad A_{ii} = 0.$$

Then the following proposition is well known and can be easily obtained from (2.7):

Proposition 2.3. (1) *If l is odd, then A has one-dimensional kernel if and only if there exists an index i_0 such that $\text{pf}(A_{i_0}) \neq 0$. In this case $\ker A$ is spanned by the vector*

$$(2.13) \quad (A_1, -A_2, \dots, (-1)^{i-1}A_i, \dots, A_l)^T.$$

(2) *If l is even then*

$$(2.14) \quad \sum_{j=1}^l (-1)^j a_{sj} A_{ij} = \begin{cases} 0 & i \neq s \\ (-1)^{s-1} \text{pf}(A) & i = s. \end{cases}$$

In particular,

(a) *if $\text{pf}(A) = 0$, then for any index $1 \leq i \leq l$ the vector*

$$(2.15) \quad v_i = (-A_{i1}, A_{i2}, \dots, (-1)^j A_{ij}, \dots, A_{il})^T$$

belongs to $\ker A$.

(b) *The matrix A has 2-dimensional kernel if and only if there exist indices i_0 and j_0 such that $A_{i_0 j_0} \neq 0$. In this case,*

$$(2.16) \quad \ker A = \text{span}\{v_{i_0}, v_{j_0}\}$$

where vectors v_{i_0} and v_{j_0} are as in (2.15).

Now apply this proposition to the Goh matrix $G(\lambda)$ using notations $G_i(\lambda)$ and $G_{ij}(\lambda)$ for the pfaffians of the corresponding skew-symmetric submatrices of $G(\lambda)$ (obtained from $G(\lambda)$ by removing the corresponding rows and columns with the relation analogous to (2.12)).

2.1. The case of distributions of odd rank. From item (1) of Proposition 2.3 it follows that if $l = \text{rank } D$ is odd, then locally

$$(2.17) \quad \begin{aligned} W_D &= \{\lambda \in D^\perp : \text{there exist } i, 1 \leq i \leq l \text{ such that } G_i(\lambda) \neq 0\} = \\ &D^\perp \setminus \{\lambda \in \tilde{D} : G_1(\lambda) = \dots = G_l(\lambda) = 0\}, \end{aligned}$$

$$(2.18) \quad \hat{\mathcal{C}}(\lambda) = \text{span} \left\{ \sum_{i=1}^l (-1)^{i-1} G_i(\lambda) \vec{H}_{X_i}(\lambda) \right\}.$$

From (2.17) it follows that for any $q \in M$ the complement of $W_D(q)$ ($:= W_D \cap D^\perp(q)$) to the fiber $D^\perp(q)$ of D^\perp is an algebraic variety depending on the *second jet* of D at q . It is easy to construct examples of germs of (l, n) -distributions with odd l and $n > l$ such that this complement is a proper algebraic subvariety of the corresponding fiber of the annihilator of the distribution (see subsection 4.4). This implies that *for generic germs of distributions of odd rank l on manifolds of dimension greater than l the set W_D is an open and dense subset of \tilde{W}_D .*

For example, if D is a rank 3 distribution, then

$$G(\lambda) = \begin{pmatrix} 0 & H_{[X_1, X_2]}(\lambda) & H_{[X_1, X_3]}(\lambda) \\ -H_{[X_1, X_2]}(\lambda) & 0 & H_{[X_2, X_3]}(\lambda) \\ -H_{[X_1, X_3]}(\lambda) & -H_{[X_2, X_3]}(\lambda) & 0 \end{pmatrix},$$

$$G_1(\lambda) = H_{[X_2, X_3]}(\lambda), \quad G_2(\lambda) = H_{[X_1, X_3]}(\lambda), \quad G_3(\lambda) = H_{[X_1, X_2]}(\lambda).$$

Note that $\{\lambda \in D^\perp : H_{[X_1, X_2]}(\lambda) = H_{[X_1, X_3]}(\lambda) = H_{[X_2, X_3]}(\lambda) = 0\} = (D^{-2})^\perp$. Consequently, in this case by (2.17) we have

$$(2.19) \quad W_D = D^\perp \setminus (D^{-2})^\perp$$

and by (2.18)

$$(2.20) \quad \widehat{C}(\lambda) = \text{span} \left\{ H_{[X_2, X_3]}(\lambda) \vec{H}_{X_1}(\lambda) - H_{[X_1, X_3]}(\lambda) \vec{H}_{X_2}(\lambda) + H_{[X_1, X_2]}(\lambda) \vec{H}_{X_3}(\lambda) \right\}.$$

2.2. The case of distributions of even rank. Now let us describe the set W_D for distributions of even rank. For any i , $1 \leq i \leq l$, define the following vector field on T^*M :

$$(2.21) \quad \mathcal{Y}_i(\lambda) = \sum_{j=1}^l (-1)^j G_{ij}(\lambda) \vec{H}_{X_j}(\lambda).$$

Then from item (2)(a) of Proposition 2.3 it follows that for any $\lambda \in \widetilde{W}_D$

$$(2.22) \quad \mathcal{Y}_i(\lambda) \in \ker(\widehat{\sigma}|_{D^\perp}(\lambda)).$$

The set W_D does not contain the following subset \mathcal{S}_1 of \widetilde{W}_D :

$$\mathcal{S}_1 = \{\lambda \in \widetilde{W}_D : \dim \ker(\widehat{\sigma}|_{D^\perp}(\lambda)) > 2\}.$$

By item (2)(b) of Proposition 2.3 locally the set \mathcal{S}_1 can be described as follows:

$$(2.23) \quad \mathcal{S}_1 = \{\lambda \in D^\perp : \text{pf } G(\lambda) = 0 \text{ and } G_{ij}(\lambda) = 0 \text{ for all } 1 \leq i < j \leq l\}.$$

Besides, by the same item (2)(b) of Proposition 2.3, for any $\lambda \in \widetilde{W}_D \setminus \mathcal{S}_1$ the kernel of $(\widehat{\sigma}|_{D^\perp}(\lambda))$ is 2-dimensional so that

$$(2.24) \quad \text{if } G_{i_0 j_0}(\lambda) \neq 0 \text{ for some } 1 \leq i_0 < j_0 \leq l, \text{ then } \ker(\widehat{\sigma}|_{D^\perp}(\lambda)) = \text{span}\{\mathcal{Y}_{i_0}(\lambda), \mathcal{Y}_{j_0}(\lambda)\}$$

Note that for $\lambda \in \widetilde{W}_D \setminus \mathcal{S}_1$ the kernel of the form $\widehat{\sigma}|_{\widetilde{W}_D}(\lambda)$ is one-dimensional if and only if the kernel of the form $\widehat{\sigma}|_{D^\perp}(\lambda)$ is not tangent to \widetilde{W}_D . Therefore, locally

$$(2.25) \quad W_D = \widetilde{W}_D \setminus (\mathcal{S}_1 \cup \mathcal{S}_2),$$

where

$$(2.26) \quad \mathcal{S}_2 = \{\lambda \in \widetilde{W}_D : d(\text{pf } G)(\lambda)(\mathcal{Y}_{i_0}(\lambda)) = d(\text{pf } G)(\lambda)(\mathcal{Y}_{j_0}(\lambda)) = 0\},$$

and the characteristic rank 1 distribution $\widehat{C}(\lambda)$ satisfies

$$(2.27) \quad \widehat{C}(\lambda) = \text{span} \left\{ d(\text{pf } G)(\lambda)(\mathcal{Y}_{i_0}(\lambda)) \mathcal{Y}_{j_0}(\lambda) - d(\text{pf } G)(\lambda)(\mathcal{Y}_{j_0}(\lambda)) \mathcal{Y}_{i_0}(\lambda) \right\},$$

where the pair of indices (i_0, j_0) is as in (2.24).

From (2.23) and (2.25) it follows that for any $q \in M$ the complement of $W_D(q)$ ($:= W_D \cap D^\perp(q)$) to the fiber $\widetilde{W}_D(q)$ is an algebraic variety depending on the *third jet* of D at q . It is easy to construct examples of germs at q of (l, n) -distributions with even l and $n > l$ such that

this complement is an algebraic subvariety (of $D^\perp(q)$), which is a proper subset of $\widetilde{W}_D(q)$ (see subsection 4.4). This implies that *for generic germs of distributions of even rank l on manifolds of dimension greater than l the set W_D is an open subset of \widetilde{W}_D .*

Let us consider the case of rank 2 distributions in more detail (see also [16, 43]). In this case the set \mathcal{S}_1 is empty. Also, from (2.2) (2.9), and (2.26) it follows that $\mathcal{S}_2 = (D^{-3})^\perp$. Substituting this into (2.25) we get

$$(2.28) \quad W_D = (D^{-2})^\perp \setminus (D^{-3})^\perp.$$

Besides, $\mathcal{Y}_1 = \vec{H}_{X_1}$ and $\mathcal{Y}_2 = -\vec{H}_{X_2}$. Substituting these formulas into (2.27) and using and using (2.9) we get that

$$(2.29) \quad \widehat{\mathcal{C}}(\lambda) = \text{span} \left\{ H_{[X_1, [X_1, X_2]]}(\lambda) \vec{H}_{X_2}(\lambda) - H_{[X_2, [X_1, X_2]]}(\lambda) \vec{H}_{X_1}(\lambda) \right\}.$$

Finally note that in all cases the characteristic rank 1 distribution $\widehat{\mathcal{C}}$ can be computed explicitly in terms of a local basis of the distribution and it depends polynomially on the fibers of W_D .

Remark 2.1. *Since the set \mathcal{S}_2 , defined by (2.26), depends on brackets of length 3 of a local basis of the distribution D , then if D is a flat distribution in Tanaka sense, of even rank, and of degree of nonholonomy 2 (i.e., such that $D \neq TM$ and $D^2 = TM$) the set \mathcal{S}_2 coincides with \widetilde{W}_D , i.e. the set W_D is empty and D has no regular abnormal extremals. Therefore our subsequent theory does not cover such distributions. For example, this is the case for two possible flat $(4, 7)$ -distributions with $\dim D^2 = 7$.*

3. JACOBI CURVES OF REGULAR ABNORMAL EXTREMALS

From now on we consider distributions D such that the set W_D is an open subset in \widetilde{W}_D . Having the manifold W_D foliated by regular abnormal extremals at our disposal, we can make a symplectification of the problem by lifting the original distribution D to W_D . The dynamics of this lift along a regular abnormal extremal is described by a curve of flags consisting of isotropic and coisotropic subspaces in a linear symplectic space, which is stable under the skew-orthogonal complement (or, shortly, of curves of symplectic flags). Such curves have a nontrivial geometry that illuminates the geometry of the distribution D itself.

To begin with, let $\pi : T^*M \rightarrow M$ be the canonical projection. The *lift* \mathcal{J} of the distribution D to W_D is the pullback of the original distribution D to W_D by the canonical projection π :

$$(3.1) \quad \widehat{\mathcal{J}}(\lambda) = \{v \in T_\lambda W_D : \pi_* v \in D(\pi(\lambda))\}.$$

Further, let $\widehat{\mathcal{V}}$ be the distribution of the tangent spaces to the fibers of W_D or the *vertical distribution*

$$(3.2) \quad \widehat{\mathcal{V}}(\lambda) = \{v \in T_\lambda W_D : \pi_* v = 0\}.$$

Obviously $\widehat{\mathcal{V}} \subset \widehat{\mathcal{J}}$. Also, from (2.5) it follows that $\widehat{\mathcal{C}} \subset \widehat{\mathcal{J}}$. Hence, $\widehat{\mathcal{V}} + \widehat{\mathcal{C}} \subset \widehat{\mathcal{J}}$ so that the distribution $\widehat{\mathcal{J}}$ has at least two distinguished subdistributions and, in particular, a distinguished line subdistribution $\widehat{\mathcal{C}}$, while the original distribution D does not have such additional structures in general. Therefore it is much more simple to study the geometry of distribution $\widehat{\mathcal{J}}$ instead of the geometry of the original distribution D and under our assumptions the geometry of D is recovered from the geometry of $\widehat{\mathcal{J}}$ (see corollary??)

Note that the distribution $\widehat{\mathcal{V}}$ is involutive. Also, given a point $q \in M$ for any λ belonging to the fiber $W_D(q)$ and any vector $X \in \mathcal{J}(\lambda)$ the vector $\pi_* X$ belongs to the same vector space $D(q)$.

The last two facts easily imply that the Lie brackets of a vertical vector field, i.e. a section of the vertical distribution $\widehat{\mathcal{V}}$, with a section of the distribution $\widehat{\mathcal{J}}$ is a section of $\widehat{\mathcal{J}}$. In particular, we have the following relation:

$$(3.3) \quad [\widehat{\mathcal{C}}, \widehat{\mathcal{V}}] \subset \widehat{\mathcal{J}}.$$

Given a subspace Λ of $T_\lambda W_D$ denote by Λ^\perp the skew-orthogonal complement of Λ with respect to the skew-symmetric form $\widehat{\sigma}|_{W_D}(\lambda)$: $\Lambda^\perp = \{v \in \Lambda : \widehat{\sigma}(v, w) = 0 \forall w \in \Lambda\}$. The subspace Λ is called *isotropic* if $\Lambda \subset \Lambda^\perp$ and *coisotropic* if $\Lambda^\perp \subset \Lambda$.

Proposition 3.1. *The following statement hold:*

(1) *The subspace $\widehat{\mathcal{J}}(\lambda)^\perp$ satisfies*

$$(3.4) \quad \widehat{\mathcal{J}}(\lambda)^\perp = \{w \in T_\lambda W_D : \pi_* w \in \pi_* \ker(\widehat{\sigma}|_{D^\perp}(\lambda))\}.$$

(2) *$\widehat{\mathcal{J}}(\lambda)^\perp \subseteq \widehat{\mathcal{J}}(\lambda)$, i.e. the space $\widehat{\mathcal{J}}(\lambda)$ is coisotropic with respect to the form $\widehat{\sigma}$;*

(3) *If the distribution D is of odd rank, then*

$$\widehat{\mathcal{J}}(\lambda)^\perp = \widehat{\mathcal{V}}(\lambda) \oplus \widehat{\mathcal{C}}(\lambda)$$

(4) *If the distribution D is of even rank then*

$$(3.5) \quad \dim(\widehat{\mathcal{J}}(\lambda)^\perp) = \dim(\widehat{\mathcal{V}}(\lambda) \oplus \widehat{\mathcal{C}}(\lambda)) + 1$$

and

$$(3.6) \quad (\widehat{\mathcal{V}}(\lambda) \oplus \widehat{\mathcal{C}}(\lambda))^\perp = \widehat{\mathcal{J}}(\lambda) + [\widehat{\mathcal{C}}, (\widehat{\mathcal{J}}(\lambda))^\perp].$$

(5) *If the distribution D is of rank 2, then*

$$(3.7) \quad \widehat{\mathcal{J}}(\lambda)^\perp = \widehat{\mathcal{J}}(\lambda)$$

and

$$(3.8) \quad [\widehat{\mathcal{C}}, \widehat{\mathcal{J}}(\lambda)] = \{v \in T_\lambda W_D : \pi_* v \in D^{-2}(\pi(\lambda))\}.$$

Proof. If $w \in T_\lambda W_D$ such that $\pi_* w \in \pi_* \ker(\widehat{\sigma}|_{D^\perp}(\lambda))$, then $w = w_1 + v$, where $w_1 \in \ker(\widehat{\sigma}|_{D^\perp}(\lambda))$ and $v \in T_\lambda D^\perp(\pi(\lambda))$. Since $\widehat{\mathcal{J}}(\lambda) \subset T_\lambda D^\perp$, for any $z \in \widehat{\mathcal{J}}(\lambda)$ one has $\widehat{\sigma}(w_1, z) = 0$. Also, $\widehat{\sigma}(v, z) = v(\pi_* z) = 0$, where in the second term of this chain of equalities v is considered as an element of $D^\perp(\pi(\lambda))$ under the natural identification of the tangent space to a vector space with the vector space itself. Therefore, $\widehat{\sigma}(w, z) = 0$, i.e. the righthand side of (3.4) is contained in $\widehat{\mathcal{J}}(\lambda)^\perp$.

The equality in (3.4) follows from the dimension count and the fact that the kernel of $\widehat{\sigma}|_{W_D}(\lambda)$ is one-dimensional. Indeed, let $\text{rank } D = l$ and $\dim M = n$. Also, set $\varepsilon = 0$, if l is odd, and $\varepsilon = 1$ if l is even. Then $\dim T_\lambda W_D = 2n - l - \varepsilon$, $\dim \widehat{\mathcal{V}}(\lambda) = n - l - \varepsilon$, dimension of the righthand side of (3.4) is equal to $\dim \widehat{\mathcal{V}}(\lambda) + 1 + \varepsilon = n - l + 1$, and $\dim \widehat{\mathcal{J}}(\lambda) = \dim \widehat{\mathcal{V}}(\lambda) + \text{rank } D = n - \varepsilon$. The latter implies that $\dim \widehat{\mathcal{J}}(\lambda)^\perp = \dim T_\lambda W_D - \dim \widehat{\mathcal{J}}(\lambda) + 1 = n - l + 1$, i.e. dimension of the righthand side of (3.4) is equal to $\dim \widehat{\mathcal{J}}(\lambda)^\perp$, which together with the conclusion of the last paragraph implies (3.4), i.e. item (1).

Item (2) of the lemma follows from the definition of the subspace $\widehat{\mathcal{J}}(\lambda)$ given by (3.1) and the fact that $\pi_* \ker(\widehat{\sigma}|_{D^\perp}(\lambda)) \subset D(\pi(\lambda))$. Item (3) follows from the fact that if l is odd then

$\ker(\widehat{\sigma}|_{D^\perp}(\lambda)) = \widehat{\mathcal{C}}(\lambda)$. Formula (3.5) of item (4) follows from the fact that if l is even, then $\dim(\ker(\widehat{\sigma}|_{D^\perp}(\lambda))) = 2$.

Now let us prove formula (3.6) of item (4). We shall use a local basis X_1, \dots, X_l of the distribution D . Let the matrix $G(\lambda)$ be as in (2.4), and G_{ij} are defined as in the paragraph before Proposition 2.3. From item (1) and formulas (2.24), (2.27) it follows that if $G_{i_0 j_0}(\lambda) \neq 0$ for some $1 \leq i_0 < j_0 \leq l$, then

$$(3.9) \quad \widehat{\mathcal{J}}(\lambda) + [\widehat{\mathcal{C}}, (\widehat{\mathcal{J}}(\lambda))^\leftarrow] = \{w \in T_\lambda W_D : \pi_* w \in D(\pi(\lambda)) + \pi_*([\mathcal{Y}_{i_0}, \mathcal{Y}_{j_0}] (\lambda))\}.$$

Then from (2.21) it follows by direct computations that

$$(3.10) \quad [\mathcal{Y}_{i_0}, \mathcal{Y}_{j_0}] = \sum_{1 \leq s_1, s_2 \leq l} (-1)^{s_1+s_2} G_{i s_1} G_{j s_2} \vec{H}_{[X_{s_1}, X_{s_2}]} \pmod{\text{span}\{\vec{H}_{X_1}, \dots, \vec{H}_{X_l}\}}.$$

This implies that for $v \in \widehat{\mathcal{V}}(\lambda)$ we have

$$(3.11) \quad \widehat{\sigma}([\mathcal{Y}_{i_0}, \mathcal{Y}_{j_0}], v) = - \sum_{s_1=1}^l (-1)^{s_1} G_{i s_1} \sum_{s_2=1}^l (-1)^{s_2} G_{j_0 s_2} dH_{[X_{s_1}, X_{s_2}]}(v).$$

Using (2.14) and the fact that $d(\text{pf}G)(v) = 0$, we get

$$\sum_{s_2=1}^l (-1)^{s_2} G_{j_0 s_2} dH_{[X_{s_1}, X_{s_2}]}(v) = - \sum_{s_2=1}^l (-1)^{s_2} H_{[X_{s_1}, X_{s_2}]} dG_{j_0 s_2}(v)$$

Substituting this into (3.11) we obtain that

$$(3.12) \quad \widehat{\sigma}([\mathcal{Y}_{i_0}, \mathcal{Y}_{j_0}] (\lambda), v) = \sum_{s_2=1}^l (-1)^{s_2} \left(\sum_{s_1=1}^l (-1)^{s_1} H_{[X_{s_1}, X_{s_2}]} G_{i_0 s_1} \right) dG_{j_0 s_2}(v).$$

Using (2.14) again and the fact that $\text{pf}G$ vanishes for on W_D , we obtain that

$$\sum_{s_1=1}^l (-1)^{s_1} H_{[X_{s_1}, X_{s_2}]} (\lambda) G_{i_0 s_1} (\lambda) = 0, \quad \forall \lambda \in W_D.$$

Therefore,

$$(3.13) \quad \widehat{\sigma}([\mathcal{Y}_{i_0}, \mathcal{Y}_{j_0}] (\lambda), v) = 0.$$

Taking into account relation (3.9) and the fact that the restriction of the form $\widehat{\sigma}$ to the $\widehat{\mathcal{V}}(\lambda)$ vanishes we get that $\widehat{\mathcal{V}} + \mathcal{C}$ is contained in the skew-orthogonal complement of $\widehat{\mathcal{J}}(\lambda) + [\widehat{\mathcal{C}}, (\widehat{\mathcal{J}}(\lambda))^\leftarrow]$. Counting the dimension, we obtain in fact that it is equal to this skew-orthogonal complement, which completes the proof of item (4).

Relation (3.7) of item (5) follows immediately from formula (3.5) of item (4) and the fact that $\dim \widehat{\mathcal{J}} - \dim \widehat{\mathcal{V}}(\lambda) = 2$ for $l = 2$. Finally, relation (3.8) of item (5) follows from (3.10). \square

It is more convenient to work with the projectivization of $\mathbb{P}T^*M$ rather than with T^*M . Here $\mathbb{P}T^*M$ is the fiber bundle over M with the fibers that are the projectivizations of the fibers of T^*M . The canonical projection $\Pi: T^*M \rightarrow \mathbb{P}T^*M$ sends the characteristic distribution $\widehat{\mathcal{C}}$ on W_D to the line distribution \mathcal{C} on $\mathbb{P}W_D$ ($:= \Pi(W_D)$), which will be also called the *characteristic distribution* of the latter manifold.

Note that the corank 1 distribution on T^*M annihilating the tautological Liouville form ς on T^*M induces a contact distribution on $\mathbb{P}T^*M$, which in turns induces the even-contact (quasi-contact) distribution $\tilde{\Delta}$ on $\mathbb{P}(D^2)^\perp \setminus \mathbb{P}(D^3)^\perp$. The characteristic line distribution \mathcal{C} is exactly the Cauchy characteristic distribution of $\tilde{\Delta}$, i.e. it is the maximal subdistribution of $\tilde{\Delta}$ such that

$$(3.14) \quad [\mathcal{C}, \tilde{\Delta}] \subset \tilde{\Delta}.$$

Further set $\mathcal{J} = \Pi_* \hat{\mathcal{J}}$, $\mathcal{V} = \Pi_* \hat{\mathcal{V}}$. Each fiber of the even-contact distribution $\tilde{\Delta}$ is endowed with the canonical skew-symmetric form $\tilde{\sigma}$, up to multiplication by a non-zero constant. This form is equal to the restriction to $\tilde{\Delta}$ of the differential of any non-zero 1-form annihilating this distribution.

Now let γ be a segment of a regular abnormal extremal, O_γ be a neighborhood of γ in $\mathbb{P}W_D$ such that the quotient

$$(3.15) \quad \mathcal{N} = O_\gamma / (\text{the charactrestic one-foliation of regular abnormal extremals})$$

is a well defined smooth manifold. Further, let $\Phi : O_\gamma \rightarrow \mathcal{N}$ be the canonical projection to the quotient manifold. Since \mathcal{C} is the Cauchy characteristic distribution of the even-contact distribution $\tilde{\Delta}$, then the push-forward $\Delta := \Phi_* \tilde{\Delta}$ of $\tilde{\Delta}$ by Φ is a contact distribution on \mathcal{N} . Each fiber of this contact distribution is endowed with the canonical symplectic form σ , defined up to up to multiplication by a non-zero constant, i.e. with the canonical conformal symplectic structure. As before, one can define the notion of skew-orthogonal complement and also of isotropic and coisotropic subspaces of the fibers of Δ with respect to this symplectic form. The subspaces of the fibers of Δ that are both isotropic and coisotropic (i.e. coincide with their skew-symmetric complement) are called *Lagrangian*.

For any $\lambda \in \gamma$ define two subspaces of $\Delta(\gamma)$ as follows:

$$(3.16) \quad J(\lambda) := d\Phi(\mathcal{J}(\lambda)), \quad V(\lambda) := d\Phi(\mathcal{V}(\lambda)).$$

Note that by constructions

$$(3.17) \quad \dim J(\lambda) - \dim V(\lambda) = \text{rank } D - 1.$$

Now we need to introduce some terminology. Let $\Lambda(\cdot)$ be a curve in a Grassmannian of subspaces of a vector space L . Let $t \mapsto \Lambda(t)$ be a parametrization of the curve $\Lambda(\cdot)$. Denote by $C(\Lambda)$ the *tautological bundle over the curve* $\Lambda(\cdot)$: the fiber of $C(\Lambda)$ over the point $\Lambda(t)$ is the vector space $\Lambda(t)$. Let $\Gamma(\Lambda)$ be the space of all smooth sections of $C(\Lambda)$. Set $\Lambda^{(0)}(t) := \Lambda(t)$ and define inductively

$$\Lambda^{(j)}(t) = \text{span} \left\{ \frac{d^k}{dt^k} \ell(t) : \ell \in \Gamma(\Lambda), 0 \leq k \leq j \right\}$$

for $j \geq 0$. The space $\Lambda^{(j)}(t)$ is called the *jth extension* or the *jth osculating subspace* of the curve $\Lambda(\cdot)$ at the point $\Lambda(t)$. Obviously, this definition is independent of the parametrization of the curve $\Lambda(\cdot)$.

Recall that for any two smooth vector fields X and Y on a manifold, one has $[X, Y] = \frac{d}{dt}(e^{-tX})_* Y|_{t=0}$, where e^{tX} denotes the flow generated by the field X . This together with (3.16) implies that

$$(3.18) \quad J^{(1)}(\lambda) = d\Phi([\mathcal{C}, \mathcal{J}](\lambda)), \quad V^{(1)}(\lambda) := d\Phi([\mathcal{C}, \mathcal{V}](\lambda)).$$

From relation (3.3) and the last relation it follows that

$$(3.19) \quad V^{(1)}(\lambda) \subset J(\lambda),$$

$$(3.20) \quad \dim J^{(1)}(\lambda) - \dim J(\lambda) \leq \text{rank} D - 1.$$

As a direct consequence of Proposition 3.1, we get

Proposition 3.2. *Assume that $\lambda \in \gamma$. The following statements hold:*

(1) $J(\lambda)^\perp \subseteq J(\lambda)$, i.e. the space $J(\lambda)$ is a coisotropic subspace of $\Delta(\gamma)$ with respect to the canonical conformal symplectic structure;

(2) If the distribution D is of odd rank then

$$(3.21) \quad J(\lambda)^\perp = V(\lambda)$$

(3) If the distribution D is of even rank then

$$\dim(J(\lambda)^\perp) = \dim(V(\lambda)) + 1$$

and

$$(3.22) \quad V(\lambda) = \left(J(\lambda) + \left((J(\lambda))^\perp \right)^{(1)} \right)^\perp.$$

(4) If the distribution D is of rank 2, then $J(\lambda)^\perp = J(\lambda)$, i.e. the space $J(\lambda)$ is a Lagrangian subspace of $\Delta(\gamma)$ with respect to the canonical conformal symplectic structure.

The curve of flags $\lambda \mapsto J(\lambda)$, $\lambda \in \gamma$, of the space $\Delta(\gamma)$ is called the (*non-extended*) *Jacobi curve of the regular abnormal extremals* γ . Since the construction of the Jacobi curves is intrinsic, any invariant of the Jacobi curve with respect to the natural action on the isotropic and coisotropic subspaces of $\Delta(\gamma)$ of the conformal symplectic group $CSp(\Delta(\gamma))$ produces an invariant of the distribution D .

Remark 3.1. Note that from item (2) of Proposition 3.2, if rank D is odd, and formula (3.22) of item (3) of Proposition 3.2, if rank D is even, it follows that the curve $\lambda \mapsto V(\lambda)$, $\lambda \in \gamma$ can be recovered from the Jacobi curve $\lambda \mapsto J(\lambda)$, $\lambda \in \gamma$.

Our next goal is to generate more curves of isotropic and coisotropic spaces from the Jacobi curves using recursively the following two basic operations with curves $\lambda \mapsto V(\lambda)$ and $\lambda \mapsto J(\lambda)$, $\lambda \in \gamma$: the osculations and taking the skew-orthogonal complement. For this let $\frac{1}{2}\mathbb{Z} := \{\frac{i}{2} : i \in \mathbb{Z}\}$ and $\frac{1}{2}\mathbb{Z}_{\text{odd}} := (\frac{1}{2}\mathbb{Z}) \setminus \mathbb{Z}$. Depending on the rank of a distribution D , to the Jacobi curve $\lambda \mapsto \{V(\lambda) \subset J(\lambda)\}$, $\lambda \in \gamma$, of a regular abnormal extremal γ we assign the curve of monotonically non-increasing (by inclusion) flags $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$, where

$$(3.23) \quad I = \begin{cases} \mathbb{Z}, & \text{if rank } D \text{ is odd,} \\ \frac{1}{2}\mathbb{Z}_{\text{odd}}, & \text{if rank } D = 2, \\ \frac{1}{2}\mathbb{Z}, & \text{if rank } D \text{ is even and greater than 2.} \end{cases}$$

This curve of flags is called the *extended Jacobi curve of the regular abnormal extremal* γ . The construction of this curve is slightly different in each of the aforementioned cases. Therefore, we describe it separately in each these cases:

- (1) **D has odd rank.** In this case the extended Jacobi curve of a regular abnormal extremal γ is the curve of flags $\lambda \mapsto \{J^i(\lambda)\}_{i \in \mathbb{Z}}$, $\lambda \in \gamma$, defined as follows: First, we set $J^0(\lambda) := J(\lambda)$. Second, let $J^{-i}(\lambda) := J^{(i)}(\lambda)$ for integer $i > 0$ (i.e. the i th osculating subspace of the curve $\lambda \mapsto J(\lambda)$, $\lambda \in \gamma$ at the point λ). Finally, assume that $J^i(\lambda) := (J^{(1-i)}(\lambda))^\angle$ for integer $i > 0$.

By construction, $J^i(\lambda) \subset J^{i-1}(\lambda)$ for all $i \in \mathbb{Z}$, subspaces $J^i(\lambda)$ are coisotropic for integer $i \leq 0$, and isotropic for integer $i > 0$. Besides, using (3.19) and the definitions of $J^i(\lambda)$ it is easy to see that

$$(3.24) \quad (J^i)^{(1)} \subset J^{i-1}$$

for all $i \in \mathbb{Z}$. Finally by (3.21) we have $V = J^1$.

- (2) **D is a rank 2 distribution.** In this case the extended Jacobi curve of a regular abnormal extremal γ is the curve of flags $\lambda \mapsto \{J^i(\lambda)\}_{i \in \frac{1}{2}\mathbb{Z}_{\text{odd}}}$, $\lambda \in \gamma$, defined as follows: First, we set $J^{\frac{1}{2}}(\lambda) := J(\lambda)$. Second, let $J^{\frac{1}{2}-i}(\lambda) := J^{(i)}(\lambda)$ for integer $i > 0$. Finally, assume that $J^{\frac{1}{2}+i}(\lambda) := (J^{\frac{1}{2}-i}(\lambda))^\angle$ for integer $i > 0$.

By construction, $J^i(\lambda) \subset J^{i-1}(\lambda)$ for all $i \in \mathbb{Z}$, subspaces $J^{\frac{1}{2}+i}(\lambda)$ are coisotropic for integer $i \leq 0$, and isotropic for integer $i \geq 0$. In particular, item (4) of Proposition 3.2 says that subspaces $J^{\frac{1}{2}}(\lambda)$ is Lagrangian. Besides, relation (3.24) holds for all $i \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$.

Finally, from (3.8) and (2.28) it follows easily that

$$(3.25) \quad V = (J^{-\frac{1}{2}})^\angle = J^{\frac{3}{2}}.$$

- (3) **D has an even rank greater than 2.** The extended Jacobi curve of a regular abnormal extremal γ is the curve of flags $\lambda \mapsto \{J^i(\lambda)\}_{i \in \frac{1}{2}\mathbb{Z}}$, $\lambda \in \gamma$, defined as follows: First, we set $J^0(\lambda) := J(\lambda)$ and $J^{\frac{1}{2}}(\lambda) := J(\lambda)^\angle$. Second, let

$$J^{-\frac{1}{2}} = (J^{\frac{1}{2}})^{(1)} + J^0.$$

Further, define recursively $J^{i-1}(\lambda) := (J^i)^{(1)}(\lambda)$ for all negative $i \in \frac{1}{2}\mathbb{Z}$. Finally, set $J^i(\lambda) := (J^{\frac{1}{2}-i}(\lambda))^\angle$ for all positive $i \in \frac{1}{2}\mathbb{Z}$.

By construction, $J^i(\lambda) \subset J^{i-\frac{1}{2}}(\lambda)$ for all $i \in \frac{1}{2}\mathbb{Z}$, subspaces $J^i(\lambda)$ are coisotropic for all nonnegative $i \in \frac{1}{2}\mathbb{Z}$ and isotropic for all positive $i \in \frac{1}{2}\mathbb{Z}$. Besides, relation (3.24) holds for all $i \in \frac{1}{2}\mathbb{Z}$.

Remark 3.2. *Note that, in contrast to the first two cases, in the third case the first osculation of each curve $\lambda \mapsto J^i(\lambda)$ does not belong in general to the next larger subspace in the flag but to the second next larger subspace in the flag. In particular, it does not satisfy the so-called compatibility with respect to differentiation assumption of [20, 21]. However, these case will be treated similarly to the first two.*

We say that a point $\lambda_0 \in \gamma$ is a *regular point of the extended Jacobi curve* $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$ if for every $i \in I$ subspaces $J^i(\lambda)$ have the same dimension in a neighborhood of λ_0 . The set of regular points is open and dense in γ . Also, if λ is a regular point, then

$$(3.26) \quad \dim J^{i-2}(\lambda) - \dim J^{i-1}(\lambda) \leq \dim J^{i-1}(\lambda) - \dim J^i(\lambda), \quad \forall i \in I, i \geq 0.$$

The last inequality together with (3.17) and (3.20) imply that

$$(3.27) \quad \dim J^{i-1}(\lambda) - \dim J^i(\lambda) \leq \text{rank } D - 1, \quad \forall i \in I.$$

Finally, we say that a point $\lambda_0 \in \gamma$ is a *point of maximal class of the extended Jacobi curve* $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$, if there exists $\nu \in I$ such that $J^\nu(\lambda_0) = \Delta(\gamma)$. If λ_0 is not a point of maximal class then there exist $\nu \in I$ such that $J^{\nu'}(\lambda_0) = J^\nu(\lambda_0) \subsetneq \Delta(\gamma)$ for all $\nu' \in I$ such that $\nu' \leq \nu$. If the point λ_0 is regular and $\nu \in I$ is the maximal index with the property above, then the subspace $J^\nu(\lambda)$ is independent of λ in a neighborhood $\tilde{\gamma}$ of λ_0 in γ . Hence, the subspace $J^\nu(\lambda_0)^\subset$ is the (maximal) common subspace for all spaces $J^i(\lambda)$, $i \in I$, $\lambda \in \tilde{\gamma}$ and the whole differential-geometric information on the extended Jacobi curve $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \tilde{\gamma}$, is contained in the curve of flags $\lambda \mapsto \{J^i(\lambda)/J^\nu(\lambda_0)^\subset\}_{i \in I}$, $\lambda \in \tilde{\gamma}$ in the conformal symplectic space $J^\nu(\lambda_0)/J^\nu(\lambda_0)^\subset$.

4. FLAG SYMBOLS OF JACOBI CURVES AND JACOBI SYMBOLS OF DISTRIBUTIONS

4.1. Flag symbols of Jacobi curves. Now we are ready to define the flag symbol of the (extended) Jacobi curve at a point. Informally speaking, the flag symbol at a point represents the tangent line to this curve at this point. To begin with the precise definition, recall that the velocity of a parametrized curve $t \mapsto \Lambda(t)$ in a Grassmannian of subspaces of a vector space L at $t = t_0$ can be naturally identified with an element of $\text{Hom}(\Lambda(t_0), \Lambda^{(1)}(t_0)/\Lambda(t_0))$. For this, take $l \in \Lambda(t_0)$ and consider any smooth section $\ell(t)$ of the tautological bundle $C(\Lambda)$ such that $\ell(t_0) = l$. Then the assignment

$$l \mapsto \frac{d}{dt}\ell(t_0) \bmod \Lambda(t_0)$$

is independent of a choice of such section and defines the desired element of $\text{Hom}(\Lambda(t_0), \Lambda^{(1)}(t_0)/\Lambda(t_0))$.

Using this construction and relation (3.24) one can identify the tangent line to the curve $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$, at λ with the line in the space $\text{Hom}(J^i(\lambda), J^{i-1}(\lambda)/J^i(\lambda))$. Moreover, again from (3.24) this line factors through the line in the space $\text{Hom}(J^i(\lambda)/J^{i+1}(\lambda), J^{i-1}(\lambda)/J^i(\lambda))$.

Therefore the tangent line to the (extended) Jacobi curve $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$ at λ can be identified with a line in the space

$$\bigoplus_{i \in I} \text{Hom}(J^i(\lambda)/J^{i+1}(\lambda), J^{i-1}(\lambda)/J^i(\lambda)).$$

Now consider the following two cases separately (the reason for the difference between these cases is explained in Remark 3.2 above):

- (1) **rank D is odd or equal to 2.** In this case let $\text{Gr}^i(\lambda) := J^i(\lambda)/J^{i+1}(\lambda)$, $i \in I$, and $\text{Gr}(\lambda) = \bigoplus_{i \in I} \text{Gr}^i(\lambda)$ be the graded space corresponding to the filtration $\{J^i(\lambda)\}_{i \in I}$. Here the set I is as in (3.23). Then the tangent line to the (extended) Jacobi curve $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$ at λ can be identified with a line in the space of degree -1 endomorphism of the graded space $\text{Gr}(\lambda)$, which will be denoted by δ_λ .

- (2) **D has an even rank greater than 2.** In this case let $\text{Gr}^i(\lambda) := J^i(\lambda)/J^{i+\frac{1}{2}}(\lambda)$, $i \in \frac{1}{2}\mathbb{Z}$, and $\text{Gr}(\lambda) = \bigoplus_{i \in \frac{1}{2}\mathbb{Z}} \text{Gr}^i(\lambda)$ be the graded space corresponding to the filtration $\{J^i(\lambda)\}_{i \in \frac{1}{2}\mathbb{Z}}$.

By relation (3.24) for any $i \in \frac{1}{2}\mathbb{Z}$ the line in the space $\text{Hom}(J^i(\lambda)/J^{i+1}(\lambda), J^{i-1}(\lambda)/J^i(\lambda))$, representing the tangent line to the curve $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$, at λ , factors through the line in the space $\text{Hom}(\text{Gr}^i(\lambda), \text{Gr}^{i-1}(\lambda))$, which will be denoted by δ_λ^i . Note that passing

from the line in $\text{Hom}(J^i(\lambda)/J^{i+1}(\lambda), J^{i-1}(\lambda)/J^i(\lambda))$ to the line δ_λ^i , some information about the tangent line to the curve $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$, at λ is lost, however it is not important for our purposes. Let δ_λ be the degree -1 endomorphism of the graded space $\text{Gr}(\lambda)$ such that $\delta_\lambda|_{\text{Gr}^i(\lambda)} = \delta_\lambda^i$.

In both cases the degree -1 endomorphism δ_λ of the corresponding graded space $\text{Gr}(\lambda)$ is defined. Further note that the conformal symplectic structure on $\Delta(\gamma)$, represented by the symplectic form σ , induces the conformal symplectic structure on the space $\text{Gr}(\lambda)$ represented by the following symplectic form $\bar{\sigma}$: if $\bar{x} \in \text{Gr}^i(\lambda)$ and $\bar{y} \in \text{Gr}^j(\lambda)$ with $i + j = 0$, then $\bar{\sigma}(\bar{x}, \bar{y}) = \sigma(x, y)$, where x and y are representatives of \bar{x} and \bar{y} in $J^i(\lambda)$ and $J^j(\lambda)$ respectively; if $i + j \neq 0$, then $\bar{\sigma}(\bar{x}, \bar{y}) = 0$.

By construction the line of endomorphisms δ_λ belongs to conformal symplectic algebra $\mathfrak{csp}(\text{Gr}(\lambda))$ (with respect to the conformal symplectic structure generated by the form $\bar{\sigma}$). The group of conformal symplectic endomorphisms of $\text{Gr}(\lambda)$, preserving the grading, acts on the lines of degree -1 endomorphisms by the conjugation. The orbit of the line δ_λ is called the *symplectic flag symbol of the (extended) Jacobi curve* $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$, at λ .

Two flag symbols act in general on different graded spaces endowed with conformal symplectic structures. They are called *isomorphic*, if they are conjugated by a conformal symplectic isomorphism between these spaces (preserving the grading). Therefore, we can fix a model graded space $X = \bigoplus_{i \in I} X^i$ with conformal symplectic structure, where, as before, $I = \mathbb{Z}$, $\frac{1}{2}\mathbb{Z}_{\text{odd}}$, or $\frac{1}{2}\mathbb{Z}$,

such that the symplectic product of elements of X^i and X^j with $i + j \neq 0$ are equal to zero. The graded space X with these properties will be called the *graded conformal symplectic space*. Further, one can look on the flag symbol as an orbit of a line of degree -1 endomorphisms from $\mathfrak{csp}(X)$ with respect to the action of the subgroup of degree 0 endomorphisms from $\mathfrak{csp}(X)$ by the conjugation. Such object will be called a *symplectic flag symbol*. In the sequel we usually refer to the flag symbol as to a line of degree -1 endomorphisms from $\mathfrak{csp}(X)$ representing this orbit.

4.2. Classification of flag symbols of Jacobi curves. Symplectic flag symbols can be easily classified. This classification is a little bit finer than the Jordan classification of nilpotent endomorphisms of a vector space. One has to take into account the presence of the grading and of the symplectic structure. When $I = \mathbb{Z}$ or $\frac{1}{2}\mathbb{Z}_{\text{odd}}$, all symplectic flag symbols were classified in [20, subsection 7.2] and this classification includes as particular case all possible flag symbols of Jacobi curves in the case of distribution of odd rank or rank 2. The classification of flag symbols of Jacobi curves in the case of distributions of even rank can be done similarly. In all cases symplectic flag symbols can be described by means of skew Young diagrams as described below.

Let δ_1 and δ_2 be symplectic flag symbols of the symplectic spaces X_1 and X_2 , which are graded by sets I_1 and I_2 respectively (here in general we do not assume that $I_1 = I_2$). The direct sum $X_1 \oplus X_2$, graded by $I_1 \cup I_2$, is equipped with the natural symplectic grading such that its i th component is the direct sum of i th components of X_1 and X_2 and the symplectic form on $X_1 \oplus X_2$ is defined as follows: the restriction of this form to X_i coincides with the symplectic form on X_i for each $i = 1, 2$ and the spaces X_1 and X_2 are skew-orthogonal in $X_1 \oplus X_2$. The direct sum $\delta_1 \oplus \delta_2$ is the degree -1 endomorphism of $X_1 \oplus X_2$ such that the restriction of it to X_i is equal to δ_i for each $i = 1, 2$. A symplectic flag symbol of a graded conformal symplectic space X is called *symplectically indecomposable* if it cannot be represented as a direct sum of two symplectic flag symbols acting on (nonzero) graded conformal symplectic spaces.

Any symplectic flag symbol is a direct sum of a finite number of symplectically indecomposable flag symbols. Therefore, in order to describe all symplectic flag symbols, it is enough to describe

all symplectically indecomposable ones. Let us consider the same three cases as in the previous section separately:

- (1) In the case of \mathbb{Z} -graded conformal symplectic spaces (i.e., when $I = \mathbb{Z}$) symplectically indecomposable symbols can be described as follows [20, subsection 7.2]): Given a non-negative integer s and an integer l such that $0 \leq l \leq 2s$ let $X_{s,l}$ be a linear symplectic space with a basis $\{e_{s-l}, \dots, e_s, f_{-s}, \dots, f_{l-s}\}$,

$$(4.1) \quad X_{s,l} = \text{span}\{e_{s-l}, \dots, e_s, f_{-s}, \dots, f_{l-s}\},$$

such that $\sigma(e_i, e_j) = \sigma(f_i, f_j) = 0$, $\sigma(e_i, f_{-i}) = (-1)^{s-i}$ and $\sigma(e_i, f_j) = 0$ if $j \neq -i$. We turn the space $X_{s,l}$ into the symplectic graded spaces by defining the subspaces of degree i to be equal to the span of all vectors with index i appearing in the tuple (4.1). Then denote by $\delta_{s,l}$ a degree -1 endomorphism of $X_{s,l}$ from the symplectic algebra such that $\delta_{s,l}(e_i) = e_{i-1}$ for $s-l+1 \leq i \leq s$, $\delta_{s,l}(e_{s-l}) = 0$, $\delta_{s,l}(f_i) = f_{i-1}$ for $-s+1 \leq i \leq l-s$, and $\delta_{s,l}(f_{-s}) = 0$. Then *any indecomposable symplectic flag symbol of a \mathbb{Z} -graded conformal symplectic space is isomorphic to the symplectic flag symbol represented by the line $\mathbb{R}\delta_{s,l}$ for a nonnegative integer s and an integer l such that $0 \leq l \leq 2s$. Moreover, symplectic flag symbols corresponding to different pair (s, l) are not isomorphic.*

It is convenient to visualize a symplectically indecomposable symbol on \mathbb{Z} -graded conformal symplectic spaces via the following skew Young diagrams, consisting of two rows, having central symmetry and with the most left box in the first row being located either above or from the left to the most left box in the second row:

$$(4.2) \quad \begin{array}{cccccc} \square & \dots & \square & \dots & \square & \\ & & \square & \dots & \square & \dots & \square \end{array} \quad \text{or} \quad \begin{array}{ccc} \square & \dots & \square \\ & & \square & \dots & \square \end{array}$$

The left diagram in (4.2) corresponds to the case $s \leq l \leq 2s$. In this case the first $2s-l$ columns from the left consist of one box (situated at the top row), the next $2(l-s)+1$ columns consist of two boxes, and the last $2s-l$ columns consist of one box (situated at the bottom row). The right diagram in (4.2) corresponds to the case $s > l$. In this case the first $l+1$ columns from the left consist of one box (situated at the top row), next $2(s-l)-1$ columns are just void, and the last $l+1$ columns consist of one box (situated at the bottom row).

Remark 4.1. *Note that in both cases the middle part (of columns with two or no boxes) consist of odd columns.*

In order to assign to a diagram of type (4.2) the symplectic flag symbol one fills out the boxes of the diagram by vectors of the bases (4.1) such that the top row is filled from the left to the right with $e_s, \dots, e_{s-1}, \dots, e_{s-l}$ and the bottom row is filled from the left to the right with $f_{l-s}, f_{l-s-1}, \dots, f_{-s}$. For example, in the case $s \leq l$ the diagram is filled as follows:

$$(4.3) \quad \begin{array}{cccccc} e_s & \dots & e_{l-s+1} & e_{l-s} & \dots & e_{s-l} \\ & & & f_{l-s} & \dots & f_{s-l} & f_{s-l-1} & \dots & f_{-s} \end{array}$$

Then the endomorphism $\delta_{s,l}$ corresponds to the *right shift* in the obtained skew Young tableaux (4.3) i.e. a vector in the basis (4.1) is sent either to the vector located in the box to the right to it in the tableaux or to zero if there is no such box. Also note that the skew Young diagram (4.2) contains all information on the grading of the corresponding vector spaces: the subspaces of the same degree are spanned by the vectors situated in the same column. Finally, the (conformal) symplectic structure is encoded by the skew Young diagram as well: the symplectic product of the vectors situated in the boxes which are not centrally symmetric is equal to zero; it is equal to ± 1 for the vectors situated in the boxes which are not centrally symmetric such that it is equal to 1 for two vectors situated in the most left and the most right boxes of the diagram, i.e. vectors e_s and f_{-s} , and it alternates signs when we took symmetric pairs by moving with the same speed from the those boxes toward the middle of the diagram.

Now let us discuss what symplectic flag symbols may appear at regular points of Jacobi curves of a distribution D of odd rank. From the relation (3.17), the fact that the subspaces $J^i(\lambda)$ are obtained by osculations of the curve $\lambda \mapsto J(\lambda)$, regularity, and the inequalities (3.27) it follows that *a symplectic flag symbol at a regular point of Jacobi curves of a distribution D of odd rank is isomorphic to a direct sum of $\frac{1}{2}(\text{rank} D - 1)$ of indecomposable flag symbols with $s \leq l \leq 2s$, i.e. given by skew Young diagrams as in the left of (4.2).*

In particular, for rank 3 distributions a symplectic flag symbol at a regular point λ of a Jacobi curve is isomorphic to an indecomposable one given by a skew Young diagrams as in the left of (4.2), i.e. $\delta_{s,l}$ with $s \leq l \leq 2s$. If, in addition, λ is a point of maximal class and the dimension of the ambient manifold M is equal to n , then the number of boxes in this skew diagram must be equal to the rank of the contact distribution Δ , i.e. to $2n - 6$. On the other hand, if the symbol is represented by the endomorphism $\delta_{s,l}$ then the number of box in the skew diagram is equal to $2l + 2$. Consequently, in the considered case $l = n - 4$ and $\frac{n-4}{2} \leq s \leq n - 4$.

Remark 4.2. *Note that a rank 3 distribution D has the Jacobi symbol $\delta_{s,l}$ with $s < l \leq 2s$ if and only if its square $D^2(q)$ is 6-dimensional at a generic point q , $\dim D^2(q) = 6$. It easily follows from the expression (2.20) for the characteristic rank 1 distribution. Also from this one can show that $\text{rank } D^2 = 4$ if and only if the Jacobi symbol of D is equal to $\mathbb{R}\delta_{0,0}$, i.e. its Young diagram consists of one column with two boxes and $\text{rank } D^2 = 5$ if and only if the Jacobi symbol of D is equal top $\mathbb{R}\delta_{s,s}$ with $s > 0$.*

- (2) In the case of $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ -graded conformal symplectic spaces (i.e., when $I = \frac{1}{2}\mathbb{Z}_{\text{odd}}$) there are two types of symplectically indecomposable symbols. First type is represented by an endomorphism $\delta_{s,l}$ as in the previous item with s being positive and belonging to $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ and l being integer and satisfying $0 \leq l \leq 2s - 1$. It is encoded again by a skew Young diagram consisting of two rows as before.

Second type can be described as follows: Given a positive $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$ let

$$(4.4) \quad \mathcal{L}_m = \bigoplus_{\substack{-m \leq i \leq m, \\ i \in \frac{1}{2}\mathbb{Z}_{\text{odd}}}} E_i$$

be a symplectic graded spaces such that $\dim E_i = 1$ for every $-m \leq i \leq m$ and let τ_m be a degree -1 endomorphism of \mathcal{L}_m which sends E_i onto E_{i-1} for every admissible $-m \leq i < m$ and $\tau_m(E_{-m}) = 0$. An indecomposable symplectic flag symbol of second type on a $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ -graded conformal symplectic space is isomorphic to the symplectic flag symbol represented by the line \mathbb{R} for a positive $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$.

It is convenient to visualize the symplectically indecomposable symbol, which is isomorphic to the symbol represented by the endomorphism τ_m , via the Young diagrams consisting of one row with $2m + 1$ boxes (note that $2m + 1$ is even, because $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$). By analogy with the previous case the endomorphism τ_m corresponds to the right shift in this diagram where the boxes from the left to the right are filled by vectors generated lines E_i , i.e.

$$(4.5) \quad \begin{array}{|c|c|c|c|c|} \hline \varepsilon_m & \varepsilon_{m-1} & \dots & \varepsilon_{1-m} & \varepsilon_{-m} \\ \hline \end{array}, \quad \text{where } E_i = \text{span}\{\varepsilon_i\}$$

By the same arguments as in the previous item for rank 2 distributions a symplectic flag symbol at a regular point of its Jacobi curves is isomorphic to the indecomposable symbol, represented by the endomorphism τ_m for some positive $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$. If, in addition, λ is a point of maximal class and the dimension of the ambient manifold M is equal to n , then the number of boxes in the corresponding skew diagram must be equal to the rank of the contact distribution Δ , i.e. to $2n - 6$. In this case all symplectic flag symbol at regular points are isomorphic to the symbol represented by the endomorphism $\tau_{n-\frac{7}{2}}$.

- (3) Now consider the case of $\frac{1}{2}\mathbb{Z}$ -graded symplectic space X . Obviously, this space is the direct sum of its \mathbb{Z} -graded component $X_{\mathbb{Z}}$ and its $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ -graded component $X_{\frac{1}{2}\mathbb{Z}_{\text{odd}}}$. Therefore, any symplectic flag symbol δ on V is the direct sum of a symplectic flag symbol on $\delta_{\mathbb{Z}}$ on $X_{\mathbb{Z}}$ and of a symplectic flag symbol $\delta_{\frac{1}{2}\mathbb{Z}_{\text{odd}}}$ on $X_{\frac{1}{2}\mathbb{Z}_{\text{odd}}}$, i.e the classification of symplectic flag symbols in the considered case is reduced to the classification of symplectic flag symbols described in the previous two items. The flag symbols $\delta_{\mathbb{Z}}$ and $\delta_{\frac{1}{2}\mathbb{Z}_{\text{odd}}}$ will call the \mathbb{Z} -graded and $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ -graded component of the flag symbol δ , respectively.

Now let us discuss what symplectic flag symbols may appear at regular points of Jacobi curves of a distribution D of even rank greater than 2. From item (3) of Proposition 3.2 it follows that $\dim \text{Gr}^{\frac{1}{2}}(\lambda) = 1$ (where $\text{Gr}^{\frac{1}{2}}(\lambda) = J^{\frac{1}{2}}(\lambda)/J^1(\lambda)$). This implies that the $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ -graded component of a symplectic flag symbol at regular points of its Jacobi curves is isomorphic to the indecomposable symbol, represented by the endomorphism τ_m for some positive $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$. Besides, by the same arguments as in item (1) the \mathbb{Z} -graded component of symplectic flag symbol at a regular point of Jacobi curves is isomorphic to a direct sum of $\frac{1}{2}(\text{rank}D - 2)$ of indecomposable flag symbols with $s \leq l \leq 2s$, i.e. given by skew Young diagrams as in the left of (4.2). In other words, in the considered case the symplectic flag symbol at a regular point of Jacobi curves of a distribution D of even rank greater than 2 is the direct sum of a symplectic flag symbol appearing at regular points of of Jacobi curves of distributions of rank equal to $\text{rank}D - 1$ and of a symplectic flag symbol appearing at regular points of Jacobi curves of rank 2 distributions.

Remark 4.3. *As a matter of fact all three cases considered separately can be unified in item (3): the case of item (1) corresponds to $X_{\frac{1}{2}\mathbb{Z}_{\text{odd}}} = 0$ and the case of item (2) corresponds to $X_{\mathbb{Z}} = 0$.*

Remark 4.4. *To an arbitrary symplectic symbol δ we can assign the skew Young diagram $\text{Diag}(\delta)$ of δ by attaching the skew Young diagrams of its indecomposable components such that the boxes corresponding to elements of the same degree are located in one column. Obviously, $\text{Diag}(\delta)$ is defined up to a permutation of sub-diagrams corresponding to indecomposable components. Note that by our convention in the $\text{Diag}(\mathbb{R}\delta_{s,l})$ the most left box of the upper row is located from the left or above of the most left box of the bottom row. Filling the boxes of the skew Young diagrams by vectors of the model graded symplectic space X as in (4.3) and (4.5) (for each undecomposable*

component) one gets a skew Young tableaux $\text{Tab}(\delta)$ of δ . In this way to any row of $\text{Tab}(\delta)$ we can assign the invariant subspace of δ in X , spanned by the vectors appearing in the boxes of this row.

4.3. Jacobi symbols of distributions. In all three cases above the set of symplectic flag symbols is discrete. Moreover, by construction the symplectic flag symbol δ_λ of the extended Jacobi curve $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$ is determined algebraically by the jets of the (non-extended) Jacobi curve $\lambda \mapsto \{V(\lambda) \subset J(\lambda)\}$, $\lambda \in \gamma$. This implies that for any point $q \in M$ there is a Zariski open subset of the fiber of the bundle $\mathbb{P}W_D(q) \rightarrow M$ over the point q such that the symplectic flag symbols of the extended Jacobi curves at all points of this subset are isomorphic to one symplectic flag symbol δ . In this case we will say that δ is the *Jacobi symbol* of a distribution D at the point q .

Moreover, in contrast to Tanaka symbols, *Jacobi symbols of distributions are always locally constant, i.e. in a neighborhood of a generic point they are isomorphic one to each other.* This leads to the following new formulation: *to construct canonical frames for distributions with given constant Jacobi symbol.*

Note that for a point λ_0 of nonmaximal class the symplectic flag symbol of the Jacobi curve at λ_0 consists of the endomorphisms acting on the graded conformal symplectic space $\text{Gr}(\lambda_0)$ of dimension less than the rank of the contact distribution Δ . Thereby, it will be more convenient in the sequel to modify the definition of the Jacobi symbol as follows. Let $\nu \in I$ be the maximal index for which the osculating subspaces of the Jacobi curve stabilize, i.e. such that $J^{\nu'}(\lambda_0) = J^\nu(\lambda_0) \subsetneq \Delta(\gamma)$ for all $\nu' \in I$ such that $\nu' \leq \nu$. This implies that $\text{Gr}^{\nu'}(\lambda_0) = 0$ for all $\nu' < \nu$ and $\nu' \geq \varepsilon - \nu$, where $\varepsilon = 1$ if D has odd rank and $\varepsilon = \frac{1}{2}$ if D has even rank. In other words, $\text{Gr}(\lambda_0) = \bigoplus_{i \in I, \nu \leq i < \varepsilon - \nu} \text{Gr}^i(\lambda_0)$. Assume that λ_0 belongs to the regular abnormal extremal γ and define the graded conformal symplectic space

$$\widetilde{\text{Gr}}(\lambda_0) = \Delta(\gamma)/J^\nu(\lambda_0) \oplus \text{Gr}(\lambda_0) \oplus J^\nu(\lambda_0)^\perp$$

such that $\Delta(\gamma)/J^\nu(\lambda_0)$ has weight preceding ν in I and $J^\nu(\lambda_0)^\perp$ has the weight $\varepsilon - \nu$. The *modified symplectic flag symbol* of the extended Jacobi curve at λ_0 is represented by the line of endomorphisms δ_{mod} from $\mathbf{csp}(\widetilde{\text{Gr}}(\lambda_0))$ such that the restrictions of these endomorphisms to $\text{Gr}(\lambda_0)$ form the line of endomorphisms of $\mathbf{csp}(\text{Gr}(\lambda_0))$, representing the symplectic flag symbol of the extended Jacobi curve at λ_0 and the restriction of the endomorphisms δ_{mod} to the added spaces $\Delta(\gamma)/J^\nu(\lambda_0)$ and $J^\nu(\lambda_0)^\perp$ is equal to zero. In other words the modified symplectic flag symbol is the direct sum of the original flag symbol with $\dim J^\nu(\lambda_0)^\perp$ copies of the symbol $\mathbb{R}\delta_{\varepsilon-\nu;0}$. Note that for the points of maximal class (and only for them) the modified symbol coincides with the original one. Finally note that in the same way one defines the modified Jacobi symbol of a distribution D at a point. In the sequel the Jacobi symbol of a distribution will be denoted by δ and the modified Jacobi symbol by δ_{mod} .

4.4. Symplectically flat distributions with given Jacobi symbols. We finish the section with an example of a distribution with given Jacobi symbol δ . For this fix a skew Young tableaux $\text{Tab}(\delta)$ and consider the subtableaux $\text{Tab}_-(\delta)$ consisting of all columns with vectors of weight not greater than $\frac{1}{2}$. Consider the Lie algebra \mathfrak{f}_δ spanned by the vectors in the boxes of $\text{Tab}_-(\delta)$ and two additional vectors, denoted by x and z , such that the only nonzero Lie brackets in the product table of the Lie algebra \mathfrak{f}_δ with respect to the chosen basis come from the following properties:

- (1) The operator $\text{ad } x$ acts as the right shift in the tableaux $\text{Tab}_-(\delta)$;
- (2) For any subtableaux of $\text{Tab}(\delta)$, corresponding to an indecomposable symbol represented by $\delta_{s,l}$ (if exists), the Lie brackets of two vectors situated in the column corresponding to weight 0 is equal to $\pm z$;
- (3) For a subtableaux of $\text{Tab}(\delta)$, corresponding to an indecomposable symbol represented by τ_m (if exists), the Lie brackets of two vectors in the box of weight $\frac{1}{2}$ and $-\frac{1}{2}$ is equal to $\pm z$.

Then consider the distribution D_δ^{SP} which is the left invariant distribution on the connected simply connected Lie group with the Lie algebra \mathfrak{f}_δ such that at the identity e of the group the subspace $D_\delta^{\text{SP}}(e)$ is equal to the span of x and the vectors from the tableaux $\text{Tab}_-(\delta)$ with the nonnegative weights (or, equivalently, with weight 0 or $\frac{1}{2}$). It can be shown by direct computations that the distribution D_δ^{SP} has Jacobi symbol δ and is of maximal class. The distribution D_δ^{SP} is called the *symplectically flat distribution with Jacobi symbol δ* . The reason for this name is that these distributions play the same role in local geometry of distribution with fixed Jacobi symbol, as flat distribution of fixed type in the local geometry of distributions with given Tanaka symbol (see the last sentence of Theorem 5.2 below).

Let us describe the symplectically flat distributions of indecomposable symbols more explicitly

Example 4.1. Assume that $\delta = \mathbb{R}\delta_{s,l}$ with nonnegative s and an integer l satisfying $s \leq l \leq 2s$. Also assume that the Young tableaux $\text{Tab}(\mathbb{R}\delta_{s,l})$ is as in (4.3). Then the Lie algebra $\mathfrak{f}_{\mathbb{R}\delta_{s,l}}$ is spanned by $e_0, \dots, e_{s-l}, f_0, \dots, f_{-s}, x, z$ (in particular it is $l+4$ -dimensional) and the only nonzero Lie brackets in the product table of the Lie algebra $\mathfrak{f}_{\mathbb{R}\delta_{s,l}}$ with respect to the chosen basis are

$$(4.6) \quad \begin{aligned} [x, e_i] &= e_{i-1}, & \forall i : s-l+1 \leq i \leq 0, i \in \mathbb{Z}; \\ [x, f_i] &= f_{i-1}, & \forall j : -s+1 \leq j \leq 0, j \in \mathbb{Z}; \\ [e_0, f_0] &= z. \end{aligned}$$

The symplectically flat distribution $D_{\mathbb{R}\delta_{s,l}}^{\text{SP}}$ with Jacobi symbol $\mathbb{R}\delta_{s,l}$ is the left invariant rank 3 distribution on the connected simply connected Lie group with the Lie algebra $\mathfrak{f}_{\mathbb{R}\delta_{s,l}}$ such that

$$D_{\mathbb{R}\delta_{s,l}}^{\text{SP}}(e) = \text{span}\{x, e_0, f_0\}.$$

We can introduce another grading by negative integers on the Lie algebra $\mathfrak{f}_{\mathbb{R}\delta_{s,l}}$ such that $\mathfrak{f}_{\mathbb{R}\delta_{s,l}}^{-1} = \text{span}\{x, e_0, f_0\}$ and the Lie algebra $\mathfrak{f}_{\mathbb{R}\delta_{s,l}}$ is fundamental with respect to this grading, i.e. it is generated by $\mathfrak{f}_{\mathbb{R}\delta_{s,l}}^{-1}$. In other words, the distribution $D_{\mathbb{R}\delta_{s,l}}^{\text{SP}}$ is the flat distribution in the Tanaka sense for the fundamental Tanaka symbol given by $\mathfrak{f}_{\mathbb{R}\delta_{s,l}}$ with the new grading.

Similarly to Example 4.1 symplectically flat distributions of odd rank are flat in Tanaka sense for the corresponding Tanaka symbol.

Example 4.2. Assume that $\delta = \mathbb{R}\tau_m$ where m is positive, belongs to $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ and the Young tableaux $\text{Tab}(\mathbb{R}\tau_m)$ is as in (4.5). Then the Lie algebra $\mathfrak{f}_{\mathbb{R}\tau_m}$ is spanned by $\varepsilon_{\frac{1}{2}}, \dots, \varepsilon_{-m}, x, z$ (in particular it is $m + \frac{7}{2}$ -dimensional) and the only nonzero Lie brackets in the product table of the Lie algebra $\mathfrak{f}_{\mathbb{R}\tau_m}$ with respect to the chosen basis are

$$(4.7) \quad \begin{aligned} [x, \varepsilon_i] &= \varepsilon_{i-1}, & \forall i : -m+1 \leq i \leq \frac{1}{2}, i \in \frac{1}{2}\mathbb{Z}_{\text{odd}}; \\ [\varepsilon_{\frac{1}{2}}, \varepsilon_{-\frac{1}{2}}] &= z. \end{aligned}$$

The symplectically flat distribution $D_{\mathbb{R}\tau_m}^{\text{SP}}$ with Jacobi symbol $\mathbb{R}\tau_m$ is the left invariant rank 2 distribution on the connected simply connected Lie group with the Lie algebra $\mathfrak{f}_{\mathbb{R}\tau_m}$ such that

$$D_{\mathbb{R}\tau_m}^{\text{SP}}(e) = \text{span}\{x, e_{\frac{1}{2}}\}.$$

As in the previous example, we can introduce another grading by negative integers on the Lie algebra $\mathfrak{f}_{\mathbb{R}\tau_m}$ such that $\mathfrak{f}_{\mathbb{R}\tau_m}^{-1} = \text{span}\{x, e_{\frac{1}{2}}\}$ and the Lie algebra $\mathfrak{f}_{\mathbb{R}\tau_m}$ is fundamental with respect to this grading. In other words, the distribution $D_{\mathbb{R}\delta_{s,l}}^{\text{SP}}$ is the flat distribution (in the Tanaka sense) for the fundamental Tanaka symbol given by $\mathfrak{f}_{\mathbb{R}\tau_m}$ with the new grading.

Example 4.3. Assume that $\delta = \mathbb{R}(\delta_{s,l} \oplus \tau_m)$ where s is a nonnegative integer, l is an integer satisfying $s \leq l \leq 2s$, and m is positive and belongs to $\frac{1}{2}\mathbb{Z}_{\text{odd}}$. Also assume that the Young tableaux $\text{Tab}(\delta)$ is the union of Young tableaux appearing in (4.3) and (4.5). Then the Lie algebra \mathfrak{f}_{δ} is spanned by $e_0, \dots, e_{s-l}, f_0, \dots, f_{-s}, \varepsilon_{\frac{1}{2}}, \dots, \varepsilon_{-m}, x, z$ (in particular it is $l + m + \frac{1}{2}$ -dimensional) and the only nonzero Lie brackets in the product table of the Lie algebra \mathfrak{f}_{δ} with respect to the chosen basis are as in (4.6) and (4.7). The symplectically flat distribution D_{δ}^{SP} with Jacobi symbol δ is the left invariant rank 4 distribution on the connected simply connected Lie group with the Lie algebra \mathfrak{f}_{δ} such that

$$(4.8) \quad D_{\delta}^{\text{SP}}(e) = \text{span}\{x, e_0, f_0, \varepsilon_{\frac{1}{2}}\}.$$

An interesting phenomenon here is that in contrast to previously discussed cases of distributions of odd rank and rank 2 the symplectically flat distribution D_{δ}^{SP} is not the flat distribution in Tanaka sense for a fundamental Tanaka symbol. Indeed, if it was the case all vectors appearing in (4.8) must have weight -1 . From this, on one hand, $z = [e_0, f_0]$ has weight -2 , but on the other hand, $\varepsilon_{-\frac{1}{2}} = [x, \varepsilon_{\frac{1}{2}}]$ has weight -2 and therefore $z = [\varepsilon_{\frac{1}{2}}, \varepsilon_{-\frac{1}{2}}]$ must have weight -3 . We got a contradiction.

However, we can take another grading of the Lie algebra \mathfrak{f}_{δ} by negative integers such that the Lie algebra \mathfrak{f}_{δ} is generated by $\mathfrak{f}_{\delta}^{-1} \oplus \mathfrak{f}_{\delta}^{-2}$ and the distribution D_{δ}^{SP} is the left invariant distribution with $D_{\delta}^{\text{SP}}(e) = \mathfrak{f}_{\delta}^{-1} \oplus \mathfrak{f}_{\delta}^{-2}$. For this set

$$\mathfrak{f}_{\delta}^{-1} = \text{span}\{\varepsilon_{\frac{1}{2}}\}, \quad \mathfrak{f}_{\delta}^{-2} = \text{span}\{x, e_0, f_0\}.$$

Note that the same phenomenon holds for all symplectically flat distribution of even rank greater than 2.

5. CANONICAL FRAME FOR DISTRIBUTIONS WITH GIVEN JACOBI SYMBOLS

To construct the canonical frame for a distribution D with given Jacobi symbol we use the general theory of curves of flags developed in [20] (with a slight modification in the case of distributions of even rank greater than 2) and the prolongation procedure for flag structures developed in our previous paper [21]. By a flag structure we mean a manifold endowed with a distribution and a submanifold of a flag variety of each fiber of this distribution. In our case the flag structure associated with the distribution D is obtained as follows: the ambient manifold is the space of regular abnormal extremals \mathcal{N} of D as in (3.15), the distribution on \mathcal{N} is the contact distribution Δ and the submanifold of flags on the fibers $\Delta(\gamma)$ are the extended Jacobi curve $\lambda \mapsto \{J^i(\lambda)\}_{i \in I}$, $\lambda \in \gamma$. Note that although \mathcal{N} may not be a manifold in general we can always overcome this issue by restricting ourselves to local considerations on W_D . Besides, in the case of distributions of odd rank or rank 2 the symplectic flag symbols of the modified extended Jacobi curves are exactly the flag symbols of this flag structure in the sense of [21].

Assume that the modified Jacobi symbol δ_{mod} of the distribution D is represented by a line δ_{mod} of degree -1 endomorphisms from $\mathbf{csp}(X)$, where $X = \bigoplus_{i \in I} X^i$ is a graded conformal symplectic space, $\dim X = \text{rank } \Delta$. Note that the grading on X induces the natural filtration $\{X_i\}_{i \in I}$ with

$$(5.1) \quad X_i = \bigoplus_{k \geq i} X^k.$$

Furthermore, the grading and the filtration on X induce the natural grading and filtration on $\mathbf{csp}(X)$. It is important to stress that the graded space $\text{gr } X = \bigoplus_{i \in I} X_i/X_{n(i)}$ associated with this filtration, where $n(i)$ is the index next to the index i in I with respect to the natural order, is canonically identified with X via the canonical identification of $X_i/X_{n(i)}$ with X^i .

The grading on $\mathbf{csp}(X)$ is defined as follows: $\mathbf{csp}(X) = \bigoplus_{j \in J} \mathbf{csp}^j(X)$, where

$$\mathbf{csp}^j(X) = \{A \in \mathbf{csp}(X) : A(X^i) \in X^{i+j} \quad \forall i \in I\}$$

and $J = \{i_1 - i_2 : i_1, i_2 \in I\}$, i.e. $J = \mathbb{Z}$, if $I = \mathbb{Z}$ or $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ and $J = \frac{1}{2}\mathbb{Z}$, if $I = \frac{1}{2}\mathbb{Z}$. The filtration on $\mathbf{csp}(X)$ is defined as follows: $\{\mathbf{csp}_j(X)\}_{j \in J}$, where

$$\mathbf{csp}_j(X) = \{A \in \mathbf{csp}(X) : A(X_i) \in X_{i+j} \quad \forall i \in I\} = \bigoplus_{k \geq j} \mathbf{csp}^k(X).$$

Now the filtration on $\mathbf{csp}(X)$ induces the filtration on any its subspace L : $\{L_j\}_{j \in J}$ with $L_j = L \cap \mathbf{csp}_j(X)$. Let $\text{gr } L$ be the corresponding graded space, $\text{gr } L = \bigoplus_{j \in J} L_j/L_{n(j)}$, where, similarly to above, $n(j)$ is the index next to the index j in J with respect to the natural order. Note that we have the following canonical identifications:

$$(5.2) \quad \text{gr } \mathbf{csp}(X) \cong \mathbf{csp}(\text{gr } X) \cong \mathbf{csp}(X).$$

The first identification is standard, in the last identification we use the identification of $\text{gr } X$ with X described above. Using these identifications the space $\text{gr } L$ can be seen as a subspace of $\mathbf{csp}(X)$, i.e. of the same space as the original subspace L . Note that in general $\text{gr } L$ differs from L .

Further note that the Tanaka symbol of the contact distribution Δ is isomorphic to the Heisenberg algebra of the corresponding dimension that will be denoted by η with the natural grading $\eta = \eta^{-1} \oplus \eta^{-2}$. The model graded conformal symplectic space X can be identified with η^{-1} . Then the algebra $\mathfrak{g}^0(\eta)$ of all derivations of η preserving the grading is equal to $\mathbf{csp}(X)$.

As in Remark 1.1 denote by $P^0(\eta)$ the principal bundle over \mathcal{N} such that the fiber $P^0(\eta)_\gamma$ over a point $\gamma \in \mathcal{N}$ consists of all isomorphisms between the conformal symplectic spaces X and $\Delta(\gamma)$, i.e. linear isomorphisms preserving the conformal symplectic structures on X and $\Delta(\gamma)$. In other words, $P^0(\eta)$ is the bundle of conformal symplectic frames of each fiber of the contact distribution Δ .

At the first step of our constructions, following [20], we would like to encode the presence of the Jacobi curves on the fibers of the contact distribution Δ by adopting to them a fiber subbundle of $P^0(\eta)$ of the minimal possible dimension in a canonical way. In contrast to Tanaka structures of Remark 1.1 such subbundle cannot be chosen in general as a principal reduction of the bundle $P^0(\eta)$. However, this subbundle belongs to more general class of fiber subbundles of P^0 that we call *quasi-principal bundles*.

To define the latter notion first note that, since the tangent spaces to the fibers of $P^0(\eta)$ are canonically identified with the Lie algebra $\mathfrak{csp}(X)$, the tangent space to a fiber subbundle P^0 of $P^0(\eta)$ at a point ψ is identified canonically with a subspace L_ψ of $\mathfrak{csp}(X)$. If P^0 is a principal reduction of $P^0(\eta)$ these subspaces are the same for different points ψ and equal to the Lie subalgebra of the structure Lie group of P . In general these subspaces vary when one varies the points ψ . However, we can weaken the assumption of constancy of these subspaces by assuming that the corresponding graded space $\text{gr } L_\psi$, seen again as the subspace of $\mathfrak{csp}(X)$ under the identifications (5.2), is independent of ψ and equal to a subalgebra \mathfrak{g}^0 of $\mathfrak{csp}(X)$. In this case we say that the fiber subbundle P^0 of $P^0(\eta)$ is a *quasi-principal bundle of type* (η, \mathfrak{g}^0) .

In our case the algebra \mathfrak{g}^0 can be described purely algebraically in terms of the modified Jacobi symbol δ_{mod} , namely in terms of the *universal algebraic prolongation* $\mathfrak{u}^F(\delta_{\text{mod}})$ of δ_{mod} (in $\mathfrak{csp}(X)$), which plays the same role in the geometry of curves of flags as the Tanaka universal algebraic prolongation on the geometry of filtered structures [20]. This is by definition the largest graded Lie subalgebra of the Lie algebra $\mathfrak{csp}(X)$ such that its negative part coincides with δ_{mod} . The algebra $\mathfrak{u}^F(\delta_{\text{mod}})$ can be explicitly constructed by induction.

First let us describe this inductive construction in the case when $I = \mathbb{Z}$ or $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ (corresponding to the cases when the original distributions D is of odd rank or of rank 2, respectively). Set $\mathfrak{u}_{-1}^F(\delta_{\text{mod}}) := \delta_{\text{mod}}$ and define by induction in $k \in \mathbb{Z}$, $k \geq 0$

$$(5.3) \quad \mathfrak{u}_k^F(\delta_{\text{mod}}) := \{v \in \mathfrak{csp}^k(X) : [v, \delta_{\text{mod}}] \in \mathfrak{u}_{k-1}^F(\delta)\}.$$

The space $\mathfrak{u}_k^F(\delta_{\text{mod}})$ is called the *kth algebraic prolongation of the symbol* δ_{mod} . Then

$$(5.4) \quad \mathfrak{u}^F(\delta_{\text{mod}}) = \bigoplus_{k \in \mathbb{Z}, k \geq -1} \mathfrak{u}_k^F(\delta_{\text{mod}})$$

In the case when $I = \frac{1}{2}\mathbb{Z}_{\text{odd}}$ (corresponding to the cases when the original distributions D is of even rank greater than 2) the following modification is needed: Set $\mathfrak{u}_{-1}^F(\delta_{\text{mod}}) := \delta_{\text{mod}}$ and $\mathfrak{u}_{-\frac{1}{2}}^F(\delta_{\text{mod}}) := 0$ and define $\mathfrak{u}_k^F(\delta_{\text{mod}})$ as in (5.3) where k are nonnegative and belong $\frac{1}{2}\mathbb{Z}_{\text{odd}}$. Then the algebra $\mathfrak{u}_k^F(\delta_{\text{mod}})$ satisfies (5.4) with \mathbb{Z} replaced by $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ in the summation of the righthand side.

As in the theory of Tanaka structures, one can define the flat curve with given symplectic flag symbol, which is in a sense the simplest (the most symmetric) curve among all curves of symplectic flags with this flag symbol. In our case consider the filtration/ the flag $\{X_i\}_{i \in I}$ on the graded conformal symplectic space X , where subspaces X_i are as in (5.1).

Definition 5.1. *Then the flat curve $\mathcal{F}_{\delta_{\text{mod}}}$ of flags with flag symbol δ_{mod} is, up to the action of the conformal symplectic group $\mathfrak{csp}(X)$ on the corresponding flag variety, the closure of the orbit of this filtration under the action of the one-parametric subgroup generated by the line δ_{mod} , namely the closure of the curve $x \mapsto \{e^x X_i\}_{i \in I}$, $x \in \delta_{\text{mod}}$.*

It can be shown by direct computations that for the symplectically flat distribution D_δ^{SP} the germ of Jacobi curves at a generic point are germs of flat curves with flag symbol δ . Moreover, the corresponding flag structure is locally isomorphic to the left-invariant flag structure on the Heisenberg group of dimension equal to $\dim \mathcal{N}$ (where \mathcal{N} is the space of regular abnormal extremals) such that at the identity of the group the curve of flags in the corresponding fiber of the contact distribution is the flat curve with flag symbol δ .

The building blocks for a more geometric description of flat curves of flags with given flag symbol are rational normal curves in projective spaces. Recall that a rational normal curve in r -dimensional projective space \mathbb{P}^r is a curve represented as

$$t \mapsto [1 : t : t^2 : \dots : t^r]$$

in some homogeneous coordinates. If $\delta_{\text{mod}} = \mathbb{R}\tau_m$ for some positive $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$ (as in the case of $(2, \frac{2m+1}{2})$ -distributions of maximal class), then the flat curve $\mathcal{F}_{\mathbb{R}\tau_m}$ is the curve of complete flags such that the curve of one dimensional subspaces of the flags in $\mathcal{F}_{\mathbb{R}\tau_m}$ is a rational normal curve in the $(2m+1)$ -dimensional projective space $\mathbb{P}X$ and all other subspaces of the flags in $\mathcal{F}_{\mathbb{R}\tau_m}$ are obtained by osculation of the rational normal curve.

For a general symplectic flag symbol δ_{mod} fix a skew Young tableaux $\text{Tab}(\delta_{\text{mod}})$ of δ_{mod} , assume that this tableaux consists of \tilde{k} rows (recall that $\tilde{k} = \text{rank } D - 1$), and let \tilde{L}_i , $1 \leq i \leq \tilde{k}$, be the subspaces of X spanned by the vectors appearing in the boxes of the i th row of $\text{Tab}(\delta_{\text{mod}})$. Then intersecting the subspaces of flags of the flat curve $\mathcal{F}_{\delta_{\text{mod}}}$ with the subspace \tilde{L}_i one gets again the curve of osculating subspaces of a rational normal curve in the projective space $\mathbb{P}\tilde{L}_i$. Informally speaking, the flat curve $\mathcal{F}_{\delta_{\text{mod}}}$ is a direct sum of \tilde{k} curves of osculating subspaces sitting in \tilde{L}_i of rational normal curves such that the grading is taken into account.

Remark 5.1. (Geometric interpretation of the universal algebraic prolongation in terms of flat curves) Finally note ([14]) that the algebra $\mathfrak{u}^F(\delta_{\text{mod}})$ is isomorphic to the algebra of infinitesimal symmetries (in $\mathfrak{csp}(X)$) of the flat curve $\mathcal{F}_{\delta_{\text{mod}}}$.

Theorem 5.1. To a germ of distribution D with constant Jacobi symbol δ one can assign in a canonical way a quasi-principal subbundle of $P^0(\eta)$ of type $(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$, where δ_{mod} is the modified Jacobi symbol.

Note that the assignment in Theorem 5.1 is uniquely determined by a choice of a so-called normalization condition for the structure equation of the moving frames associated with the extended Jacobi curves. The normalization conditions are given by a subspace W complementary to the subspace $\mathfrak{u}_+^F(\delta_{\text{mod}}) + [\delta_{\text{mod}}, \mathfrak{csp}_+(X)]$ in $\mathfrak{csp}_+(X)$, where $\mathfrak{csp}_+(X)$ is the nonnegative part of the graded algebra $\mathfrak{csp}(X)$ and $\mathfrak{u}_+^F(\delta_{\text{mod}}) = \mathfrak{u}^F(\delta_{\text{mod}}) \cap \mathfrak{csp}_+(X)$. This theorem is a direct consequence of [20, Theorem 4.4] applied to the extended Jacobi curve of each regular abnormal extremal γ in the case of distributions of odd rank or rank 2 and a straightforward generalization of this result in the case of distributions of even rank greater than 2. As was already mentioned in Remark 3.2 the latter case does not satisfy the compatibility with respect to differentiation assumption. However the constructions in this case are literally the same as in the proof of Theorem 4.4 of [20].

The result of Theorem 5.1 can be seen as the zero step of the prolongation procedure for the construction of the canonical frame for the distribution D if compared with the prolongation procedure for Tanaka structures (as in Remark 1.1 and Theorem 1.1). In [21] we have shown that the analog of Theorem 1.1 is valid for quasi-principle bundle as well. This implies the following

Theorem 5.2. Assume that D is a distribution with constant Jacobi symbol δ and δ_{mod} is the modified Jacobi symbol of D . Assume that the universal algebraic prolongation $\mathfrak{u}(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ of the pair $(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ is finite dimensional and $k \geq 0$ is the maximal integer such that $\mathfrak{u}^k(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ is not zero. Then to the distribution D one can assign, in a canonical way, a sequence of bundles $\{P^i\}_{i=0}^k$ such that P^0 is as in Theorem 5.1 and P^i for $i \geq 0$ is an affine bundle over P^{i-1} with fibers being affine spaces over the linear space of dimension equal to $\dim \mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ for any $i = 1, \dots, k$ and such that P^k is endowed with the canonical frame. In particular, the dimension

of the algebra of infinitesimal symmetries of a distribution D with Jacobi symbol δ is not greater than $\dim \mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$. This upper bound is sharp and is achieved if and only if a distribution is locally equivalent to the symplectically flat distribution D_δ^{sp} for which the algebra of infinitesimal symmetries is isomorphic to $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$.

Note that the statement of this Theorem about sharpness of the bounds and uniqueness of maximally symmetric model up to an equivalence does not directly follow from Theorem 2.3 of [21] but it can be proved without difficulties using the arguments of existence of so-called grading element in the algebra $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ similarly to [33],[42].

6. JACOBI SYMBOLS δ WITH FINITE DIMENSIONAL ALGEBRA $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$

Now the natural question is for what Jacobi symbols δ is the algebra $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ finite dimensional?

Theorem 6.1. *Assume that D is a distribution with constant Jacobi symbol δ and δ_{mod} is the modified Jacobi symbol of D .*

- (1) *If $\text{rank } D$ is odd then $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ is finite dimensional if and only if δ_{mod} is isomorphic to the direct sum of symplectically indecomposable symbols generated by $\delta_{s,l}$ with $s < l \leq 2s$, s, l are positive integers. In other words, each symplectically indecomposable component of δ_{mod} has the skew Young diagram consisting of two rows as in (4.2) and having at least three columns with two boxes;*
- (2) *If $\text{rank } D = 2$ then $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ is finite dimensional if and only if δ_{mod} is isomorphic to the symbol generated by τ_m with $m > \frac{1}{2}$, $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$. In other words, the skew Young diagram of δ_{mod} consists of one row with at least 4 boxes.*
- (3) *If $\text{rank } D$ is even then $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ is finite dimensional if and only if δ_{mod} is isomorphic to the direct sum of symplectically indecomposable symbols generated by $\delta_{s,l}$ with $s < l \leq 2s$, s, l are positive integers, and of exactly one symplectic symbol generated by τ_m with $m \geq \frac{1}{2}$, $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$. In other words, the symplectically indecomposable components of δ_{mod} have the skew Young diagrams consisting of either two rows as in (4.2) with at least 3 columns with two boxes or of one row and the latter component appears exactly ones.*

Note that in item (3) of this theorem, in contrast to item 2, the Young diagram of the symplectically indecomposable component consisting of one row may have 2 boxes.

As a direct consequence of Theorem 6.1 we get the following

Corollary 6.1. *If $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ is finite dimensional then $\delta = \delta_{\text{mod}}$, i.e. the distribution D is of maximal class.*

Also as a direct consequence of item (1) of Theorem 6.1 and Remark 4.2 one has the following

Corollary 6.2. *For rank a 3 distribution D the algebra $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ is finite dimensional if and only if D is of maximal class and $\dim D^2(q) = 6$ for generic point q .*

Using Remark 4.1, Theorem 6.1 can be equivalently reformulated in the following shorter way:

Theorem 6.2. *Assume that D is a distribution with constant Jacobi symbol δ and δ_{mod} is the modified Jacobi symbol of D . Then the algebra $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ is finite dimensional if and only if the skew Young diagram $\text{Diag}(\delta_{\text{mod}})$ of δ_{mod} satisfies both of the following two conditions:*

- (1) *It does not contain two rows such that the most left box of one of the rows is located in the same column or from the right to the most right box of another row;*
- (2) *It does not consist of one row with two boxes.*

Proof of Theorem 6.2 The proof is based on two general criteria for finiteness of prolongations of Tanaka and Spencer, respectively, and on some theory of representation of \mathfrak{sl}_2 .

Let us first describe these criteria in general setting. Let as before $\mathfrak{t} = \bigoplus_{j \leq -1} \mathfrak{g}^j$ be a graded

nilpotent Lie algebra, $\mathfrak{g}^0(\mathfrak{t})$ be the algebra of differentiations of \mathfrak{t} preserving the grading, and \mathfrak{g}^0 be a subalgebra of $\mathfrak{g}^0(\mathfrak{t})$. Tanaka criterium reduces the question whether the Tanaka prolongation $\mathfrak{u}(\mathfrak{t}, \mathfrak{g}^0)$ of a pair $(\mathfrak{t}, \mathfrak{g}^0)$ is finite-dimensional or not to the same question for the standard prolongation (as described, for example, in [26, 38]) of the following subalgebra $\mathfrak{r}(\mathfrak{g}^0)$ of \mathfrak{g}^0 .

$$(6.1) \quad \mathfrak{r}(\mathfrak{g}^0) = \{f \in \mathfrak{g}^0 \mid f(x) = 0 \text{ for all } x \in \mathfrak{g}^i, i \leq -2\}.$$

Recall that $\mathfrak{g}^0(\mathfrak{t})$ (and hence $\mathfrak{r}(\mathfrak{g}^0)$) can be considered as a subalgebra of $\mathfrak{gl}(\mathfrak{g}^{-1})$. The standard prolongation of $\mathfrak{r}(\mathfrak{g}^0)$ is nothing but the Tanaka prolongation of the pair $(\mathfrak{g}^{-1}, \mathfrak{r}(\mathfrak{g}^0))$, where \mathfrak{g}^{-1} is considered as a commutative graded Lie algebra with the trivial grading, i.e. such that all elements of \mathfrak{g}^{-1} have degree -1 . Note that the standard prolongation can be defined not only for a subalgebra but for any subspace of $\mathfrak{gl}(\mathfrak{g}^{-1})$ ([26, 38]). Tanaka criterium can be formulated as follows:

Theorem 6.3. Tanaka criterium ([32]). *The Tanaka prolongation $\mathfrak{u}(\mathfrak{t}, \mathfrak{g}^0)$ of the pair $(\mathfrak{t}, \mathfrak{g}^0)$ is finite-dimensional if and only if so is the standard prolongation of $\mathfrak{r}(\mathfrak{g}^0)$.*

Remark 6.1. *In the case when $\mathfrak{t} = \eta (= \eta^{-1} \oplus \eta^{-2})$, the Heisenberg algebra, $\mathfrak{g}^0(\eta)$ is the conformal symplectic algebra $\mathfrak{csp}(\eta^{-1})$ of η^{-1} , and the $\mathfrak{r}(\mathfrak{g}^0(\eta))$ is equal to the symplectic algebra $\mathfrak{sp}(\eta^{-1})$. This implies also that the algebra $\mathfrak{r}(\mathfrak{u}^F(\delta_{\text{mod}}))$ is equal to the universal algebraic prolongation of the symbol δ in the algebra $\mathfrak{sp}(\eta^{-1})$, i.e. the largest graded Lie subalgebra of the Lie algebra $\mathfrak{sp}(\eta^{-1})$ such that its negative part coincides with δ . Obviously,*

$$(6.2) \quad \mathfrak{u}^F(\delta_{\text{mod}}) = \mathfrak{r}(\mathfrak{u}^F(\delta_{\text{mod}})) \oplus \mathbb{R} \text{Id}_{\eta^{-1}}.$$

In its turn, Spencer criterium provides a computationally efficient method of detecting whether the standard prolongation of a linear subspace $A \subset \mathfrak{gl}(Y)$ of a vector space Y over the field k is finite-dimensional or not. Let \bar{k} be the algebraic closure of the field k and by $A^{\bar{k}}$ be the subspace in $\mathfrak{gl}(Y^{\bar{k}})$ obtained from A by field extension.

Theorem 6.4. Spencer criterium ([37, 23, 31]) *The standard prolongation of a subspace $A \subset \mathfrak{gl}(Y)$ is finite-dimensional if and only if $A^{\bar{k}} \subset \mathfrak{gl}(Y^{\bar{k}})$ does not contain endomorphisms of rank 1.*

The universal algebraic prolongations of symplectic flag symbols in the symplectic algebra (and therefore, according to (6.2), in the conformal symplectic algebra) were described quite explicitly in [20, section 8] using the theory of \mathfrak{sl}_2 representation. Let us briefly list the main features of this algebras in context of the Spencer criterium.

We start with some terminology. We say that a graded space X which is also \mathfrak{sl}_2 -module is a *nice \mathfrak{sl}_2 -module*, if the corresponding embedding of \mathfrak{sl}_2 into $\mathfrak{gl}(X)$ is spanned by endomorphisms of degree -1 , 0 , and 1 . Let Y_1 and Y_2 be two nice \mathfrak{sl}_2 -modules. Then $\text{Hom}(Y_1, Y_2) = Y_2 \otimes Y_1^*$ is the \mathfrak{sl}_2 -module and a graded space in a natural way. Denote by $\mathfrak{n}(Y_1, Y_2)$ the maximal \mathfrak{sl}_2 -submodule of $\text{Hom}(X, Y)$ concentrated in the non-negative degree part. If $Y_1 = Y_2$ we simply will write $\mathfrak{n}(Y_1)$ instead of $\mathfrak{n}(Y_1, Y_2)$.

Further, assume that X is a graded conformal symplectic space which is also a nice \mathfrak{sl}_2 -module such that the corresponding embedding of \mathfrak{sl}_2 into $\mathfrak{gl}(X)$ belongs to $\mathfrak{sp}(X)$. In this case we will say that X is a *nice symplectic \mathfrak{sl}_2 -module*. The conformal symplectic algebra $\mathfrak{csp}(X)$ is a \mathfrak{sl}_2 -module and a graded space in a natural way. Denote by $\mathfrak{l}(X)$ the maximal \mathfrak{sl}_2 -submodule of $\mathfrak{sp}(X)$ concentrated in the non-negative degree part. From the maximality assumption it follows that $\mathfrak{l}(X)$ is a subalgebra of $\mathfrak{sp}(X)$.

Given a symplectic flag symbol ρ on a graded conformal symplectic space X we can make X to a nice symplectic \mathfrak{sl}_2 -module with ρ being degree -1 component of the corresponding embedding φ_ρ of \mathfrak{sl}_2 into $\mathfrak{sp}(X)$. Moreover, if one fix a skew Young tableaux $\text{Tab}(\rho)$ of ρ (as described in Remark 4.4), then each row in $\text{Tab}(\rho)$ defines an irreducible \mathfrak{sl}_2 -submodule. From [20, subsection 8.3] it follows that the universal algebraic prolongation $\mathfrak{r}(\mathbf{u}^F(\rho))$ of the symbol ρ in $\mathfrak{csp}(X)$ is equal to the semidirect sum of the constructed embedding φ_ρ of \mathfrak{sl}_2 into $\mathfrak{sp}(X)$ and the algebra $\mathfrak{l}(X)$,

$$(6.3) \quad \mathfrak{r}(\mathbf{u}^F(\rho)) = \mathfrak{l}(X) \rtimes_{\varphi_\rho} \mathfrak{sl}_2.$$

Remark 6.2. *By Remarks 5.1 and 6.1 the algebra $\mathfrak{r}(\mathbf{u}^F(\rho))$ is isomorphic to the subalgebra of $\mathfrak{sp}(X)$ consisting of all infinitesimal symmetries of the flat curve \mathcal{F}_ρ with the symbol ρ . Hence, to any element of $\mathfrak{r}(\mathbf{u}^F(\rho))$ we can assign the vector field on the curve \mathcal{F}_ρ . It turns out that the ideal $\mathfrak{l}(X)$ consists of all elements of $\mathfrak{r}(\mathbf{u}^F(\rho))$ for which the corresponding vector fields on \mathcal{F}_ρ are identically zero. In other words, $\mathfrak{l}(X)$ is the subalgebra of $\mathfrak{sp}(X)$ consisting of the infinitesimal symmetries of the flat curve \mathcal{F}_ρ considered as a parametrized curve. This implies that $A \in \mathfrak{l}(X)$ if and only if*

$$(6.4) \quad A(e^x(X_i)) \subset e^x(X_i), \quad \forall x \in \rho, \quad i \in I,$$

where $\{X_i\}_{i \in I}$ is the filtration of X as in (5.1).

Remark 6.3. *Note that analogous statements are true if we forget about the presence of symplectic structure on X and consider curves of flags with respect to the action of $GL(X)$. In all definitions and statements the algebras $\mathfrak{csp}(X)$ or $\mathfrak{sp}(X)$ should be replaced by $\mathfrak{gl}(X)$, the algebra $\mathfrak{l}(X)$ is replaced by $\mathfrak{n}(X)$ and the latter satisfies relation (6.4).*

1. Rank 1 elements in $\mathfrak{l}(x)$

Our first goal is to prove the following

Proposition 6.1. *In the case when $\rho = \delta_{mod}$ the algebra $\mathfrak{l}(X)$ contains elements of rank 1 if and only the corresponding skew Young diagram $\text{Diag}(\delta_{mod})$ does not satisfy condition (1) of Theorem 6.2, i.e. it contains two rows such that the most left box of one of the rows is located in the same column or from the right to the most right box of another row*

Proof. In this case ρ is isomorphic to the direct sum of $N(s, l)$ copies of the indecomposable symbols $\mathbb{R}\delta_{s, l}$ for some finite number of pairs (s, l) with $s \geq 0$ and $0 \leq l \leq 2s$ and of at most one copy of $\mathbb{R}\tau_m$ for some $m \in \frac{1}{2}\mathbb{Z}_{\text{odd}}$. Then we can split X as

$$(6.5) \quad X = \bigoplus_{i=1}^k L_i,$$

for some $k \in \mathbb{N}$, where $N(s, l)$ of subspaces L_i are isomorphic to $X_{s, l}$ and at most one subspace L_i is isomorphic to \mathcal{L}_m , where spaces $X_{s, l}$ and \mathcal{L}_m are as in (4.1) and (4.4), respectively. The

restriction of ρ to L_i is isomorphic to the indecomposable symbols $\mathbb{R}\delta_{s,l}$ in the first case and to $\mathbb{R}\tau_m$ in the second case. Further, the restriction $\sigma|_{L_i}$ to L_i of the symplectic form σ of V defines natural identification between L_i and its dual space L_i^* . Here σ denotes the symplectic form on X . Splitting (6.5) defines the following (in general non-canonical) splitting of $\mathfrak{gl}(X)$:

$$(6.6) \quad \mathfrak{gl}(X) = \bigoplus_{i,j=1}^k \text{Hom}(L_i, L_j).$$

An endomorphism $A \in \mathfrak{gl}(V)$, having the decomposition $A = \sum_{i,j=1}^k A_{ij}$ with respect to the splitting (6.6), belongs to $\mathfrak{sp}(X)$ if and only if $A_{ii} \in \mathfrak{sp}(L_i)$ for all $1 \leq i \leq k$ and $A_{ij} = -A_{ji}^*$ for all $1 \leq i \neq j \leq k$ (here the dual linear map A_{ji}^* is considered as a map from L_i to L_j under the aforementioned identification $L_i \sim L_i^*$ and $L_j \sim L_j^*$). Therefore the map $A \mapsto \sum_{i=1}^k A_{ii} + \sum_{1 \leq i < j \leq k} A_{ij}$ defines the identification

$$(6.7) \quad \mathfrak{sp}(X) \cong \bigoplus_{i=1}^k \mathfrak{sp}(L_i) \oplus \bigoplus_{1 \leq i < j \leq k} \text{Hom}(L_i, L_j).$$

This implies that

$$(6.8) \quad \mathfrak{l}(X) \cong \bigoplus_{i=1}^k \mathfrak{l}(L_i) \oplus \bigoplus_{1 \leq i < j \leq k} \mathfrak{n}(L_i, L_j).$$

Note that by constructions the space $\mathfrak{l}(X)$ is an invariant subspace with respect to the projections on each component of the decomposition (6.7). Therefore the question of the existence of rank 1 elements in $\mathfrak{l}(X)$ can be reduced to the same question regarding each component of the decomposition (6.8). Moreover, let us show that in the question of existence of rank 1 elements in $\mathfrak{l}(X)$ one can get rid of multiple copies of $X_{s,l}$ in the splitting (6.5) of X .

Lemma 6.1. *The algebra $\mathfrak{l}(X)$ contains rank 1 elements if and only if at least one of the following algebras does:*

- (1) $\mathfrak{n}(X_{s_1,l_1}, X_{s_2,l_2})$, if X_{s_1,l_1} and X_{s_2,l_2} appear as components in the splitting (6.5);
- (2) $\mathfrak{n}(\mathcal{L}_m, X_{s,l})$, if \mathcal{L}_m and $X_{s,l}$ appear as components in the splitting (6.5);
- (3) $\mathfrak{l}(X_{s,l})$, if $X_{s,l}$ appears as a component in the splitting (6.5);
- (4) $\mathfrak{l}(\mathcal{L}_m)$, if \mathcal{L}_m appears as a component in the splitting (6.5);

Proof. Applying formula (6.7) to the graded conformal symplectic space $X_{s,l} \otimes \mathbb{R}^{N(s,l)}$ instead of X it is easy to get the following natural identification:

$$(6.9) \quad \mathfrak{sp}(X_{s,l} \otimes \mathbb{R}^{N(s,l)}) \cong \left(\mathfrak{sp}(X_{s,l}) \otimes \mathbb{R}^{N(s,l)} \right) \oplus \left(\mathfrak{gl}(X_{s,l}) \otimes \wedge^2 \mathbb{R}^{N(s,l)} \right),$$

where $\wedge^2 \mathbb{R}^{N(s,l)}$ denotes a skew-symmetric square of $\mathbb{R}^{N(s,l)}$. Consequently,

$$(6.10) \quad \mathfrak{l}(X_{s,l} \otimes \mathbb{R}^{N(s,l)}) \cong \left(\mathfrak{l}(X_{s,l}) \otimes \mathbb{R}^{N(s,l)} \right) \oplus \left(\mathfrak{n}(X_{s,l}) \otimes \wedge^2 \mathbb{R}^{N(s,l)} \right).$$

Besides, for any two nice \mathfrak{sl}_2 -modules X_1 and X_2

$$(6.11) \quad \mathfrak{n}(X_1 \otimes \mathbb{R}^{N_1}, X_2 \otimes \mathbb{R}^{N_2}) \cong \mathfrak{n}(X_1, X_2) \otimes \text{Hom}(\mathbb{R}^{N_1}, \mathbb{R}^{N_2}).$$

Combining (6.10) and (6.11) with (6.8), one obtains the statement of the lemma. \square

Now analyze the algebras appearing in Lemma 6.1 separately:

a) Algebras from the first two items of Lemma 6.1

The calculation of algebras $\mathfrak{n}(X_{s_1, l_1}, X_{s_2, l_2})$ and $\mathfrak{n}(\mathcal{L}_m, X_{s, l})$ from the first two items of Lemma 6.1 is reduced to the calculation of $\mathfrak{n}(Y_1, Y_2)$ where Y_1 and Y_2 are two disjoint irreducible nice \mathfrak{sl}_2 -modules graded by the sets I_1 and I_2 , where each I_i is either \mathbb{Z} or $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ (note that I_1 and I_2 might be different). For each $i \in \{1, 2\}$, let r_i and s_i be the lowest and the highest weights of element of Y_i respectively and let $l_i = s_i - r_i$ ($= \dim Y_i - 1$). Recall that the corresponding \mathfrak{sl}_2 -submodule $\text{Hom}(Y_1, Y_2) = Y_2 \otimes Y_1^*$ is decomposed into the irreducible \mathfrak{sl}_2 -modules as follows:

$$(6.12) \quad \text{Hom}(Y_1, Y_2) \cong \bigoplus_{i=0}^{\min\{l_1, l_2\}} \Pi_{l_1+l_2-2i},$$

where Π_j denotes an irreducible \mathfrak{sl}_2 -module of dimension $j + 1$ (see, for example, [22]). Note that the following lemma immediately follows from the construction of the decomposition (6.12)

Lemma 6.2. *The only submodule in the decomposition (6.12) containing rank 1 elements is the submodule $\Pi_{l_1+l_2}$ of the largest dimension.*

Furthermore, the following formula can be easily verified (see [20, subsection 8.2]) using the decomposition (6.12):

$$(6.13) \quad \mathfrak{n}(Y_1, Y_2) = \begin{cases} \bigoplus_{i \in \mathbb{Z} \cap [\max\{0, s_1 - r_2\}, \min\{l_1, l_2\}]} \Pi_{l_1+l_2-2i}, & s_2 \geq s_1 \text{ and } r_2 \geq r_1 \\ 0, & \text{otherwise.} \end{cases}$$

From this formula and Lemma 6.2 it follows that $\mathfrak{n}(Y_1, Y_2)$ contains elements of rank 1 if and only if $\mathfrak{n}(Y_1, Y_2) = \text{Hom}(Y_1, Y_2)$ or, equivalently, all elements of $\text{Hom}(Y_1, Y_2)$ have nonnegative degrees, or, equivalently, $s_1 \leq r_2$, i.e. the maximal degree of elements of Y_1 does not exceed the minimal degree of elements of Y_2 . Consider the skew Young diagram consisting of two rows with $l_1 + 1$ and $l_2 + 1$ boxes, respectively, such that the columns of it are indexed by the set I in the decreasing order from left to right and for any $i \in I_1 \cup I_2$ the number of boxes in the column indexed by i is equal to the dimension of the subspace of degree i in $Y_1 \oplus Y_2$. Then the conditions above are equivalent to the fact that the obtained skew Young diagram does not satisfy condition (1) of Theorem 6.2.

b) Algebras from item (3) of Lemma 6.1 Now consider the algebra $\mathfrak{l}(X_{s, l})$ from item (3) of Lemma 6.1. The space $X_{s, l}$ is the sum of two irreducible nice \mathfrak{sl}_2 -submodules

$$(6.14) \quad E_{s, l} = \text{span}\{e_i\}_{s-l \leq i \leq s}, \quad F_{s, l} = \text{span}\{f_i\}_{-s \leq i \leq l-s},$$

where $\{e_i\}_{s-l \leq i \leq s}$ and $\{f_i\}_{-s \leq i \leq l-s}$ are as in (4.1). Note that $E_{s, l}$ and $F_{s, l}$ are two transversal Lagrangian subspaces of $X_{s, l}$. Therefore the symplectic form on $X_{s, l}$ defines the following natural identifications: $E_{s, l} \cong (F_{s, l})^*$ and $F_{s, l} \cong (E_{s, l})^*$. Keeping in mind these identifications, an endomorphism $A \in \mathfrak{gl}(V)$, which has the decomposition $A = A_{11} + A_{22} + A_{12} + A_{21}$ with respect to the splitting

$$(6.15) \quad \mathfrak{gl}(X_{s, l}) = \mathfrak{gl}(E_{s, l}) \oplus \mathfrak{gl}(F_{s, l}) \oplus \text{Hom}(F_{s, l}, E_{s, l}) \oplus \text{Hom}(E_{s, l}, F_{s, l}),$$

belongs to $\mathfrak{sp}(X_{s, l})$ if and only if

$$(6.16) \quad A_{12}^* = A_{12}, \quad A_{21}^* = A_{21}, \quad \text{and} \quad A_{22} = -A_{11}^*.$$

Therefore the map $F_{s,l} : A \mapsto A_{11} + A_{22} + A_{12}$ defines the following identifications

$$(6.17) \quad \mathfrak{sp}(X_{s,l}) \cong \mathfrak{gl}(E_{s,l}) \oplus S^2(E_{s,l}) \oplus S^2(F_{s,l}).$$

Here $S^2(E_{s,l})$ is a subspace of $\text{Hom}(F_{s,l}, E_{s,l})$ and $S^2(F_{s,l})$ is a subspace of $\text{Hom}(E_{s,l}, F_{s,l})$. Keeping this in mind, we define $\mathfrak{s}(E_{s,l})$ as the maximal \mathfrak{sl}_2 -submodule of $S^2(E_{s,l})$ concentrated in the non-negative degree part of $S^2(E_{s,l})$. Similarly, let $\mathfrak{s}(F_{s,l})$ be the maximal \mathfrak{sl}_2 -submodule of $S^2(F_{s,l})$ concentrated in the non-negative degree part of $S^2(F_{s,l})$.

Then from (6.17) it follows that

$$(6.18) \quad \mathfrak{l}(X_{s,l}) \cong \mathfrak{n}(E_{s,l}) \oplus \mathfrak{s}(E_{s,l}) \oplus \mathfrak{s}(F_{s,l})$$

Obviously, $\mathfrak{l}(X_{s,l})$ contains an element of rank 1 if and only if at least one of the spaces $\mathfrak{s}(E_{s,l})$, $\mathfrak{s}(F_{s,l})$ or $F^{-1}(\mathfrak{n}(E_{s,l}))$ contains an element of rank 1. Now analyze each of this spaces. Start with $\mathfrak{s}(E_{s,l})$. By analogy with (6.12), \mathfrak{sl}_2 -module $S^2(E_{s,l})$ is decomposed into the irreducible \mathfrak{sl}_2 -modules in the following way ([22]):

$$(6.19) \quad S^2(E_{s,l}) = \bigoplus_{i=0}^{\lfloor \frac{l}{2} \rfloor} \Pi_{2l-4i}.$$

Using the last decomposition, the following formula can be easily verified (see [20, subsection 8.3.1]):

$$(6.20) \quad \mathfrak{s}(E_{s,l}) = \bigoplus_{i=\max\{0, l-s\}}^{\lfloor \frac{l}{2} \rfloor} \Pi_{2l-4i},$$

From this formula and the obvious analogues of Lemma 6.2 it follows that $\mathfrak{l}_{s,l}^1$ contains element of rank 1 if and only if $\mathfrak{s}(E_{s,l}) = S^2(E_{s,l})$ or, equivalently, all elements of $S^2(E_{s,l})$ have nonnegative degrees or, equivalently, $l \geq s$, i.e. the maximal degree of elements of $F_{s,l}$ does not exceed the minimal degree of elements of $E_{s,l}$. The latter condition in turn is equivalent to the fact that the corresponding skew Young diagram as in (4.2) does not satisfy condition (1) of Theorem 6.2.

Further, the \mathfrak{sl}_2 -module $S^2(F_{s,l})$ has the decomposition to irreducible submodules similar to (6.19). However, the grading on these irreducible submodules is different from the case of $S^2(E_{s,l})$, because it is induced from $\text{Hom}(E_{s,l}, F_{s,l})$ instead of $\text{Hom}(F_{s,l}, E_{s,l})$. One can easily show [20, subsection 8.3.1] that

$$(6.21) \quad \mathfrak{s}(F_{s,l}) = \begin{cases} \mathbb{R} & \text{if } l \text{ is even and } s = \frac{l}{2}, \\ 0 & \text{otherwise.} \end{cases}$$

and it contains a rank 1 element if and only if $s = l = 0$, i.e. when the corresponding Young diagram consists of one column and two rows. The latter condition trivially does not satisfy condition (1) of Theorem 6.2. Note that the first option in (6.21) is equivalent to the fact that the corresponding Young diagram is rectangular.

Finally, note that from (6.13) applied to the case $Y_1 = Y_2$ it follows that

$$(6.22) \quad \mathfrak{n}(E_{s,l}) \cong \Pi_0 \cong \mathbb{R} \text{Id}_{E_{s,l}}$$

and, taking into account the last relation of (6.16), it follows that the space $(F_{s,l})^{-1}(\mathfrak{n}(E_{s,l}))$ has no element of rank 1 or, equivalently, elements A in $\mathfrak{l}(X_{s,l})$ of the form $A = A_{11} + A_{22}$ with respect to decomposition (6.15) are never of rank 1.

Remark 6.4. *As the matter of fact $(F_{s,l})^{-1}(\mathfrak{n}(E_{s,l}))$ is spanned by the endomorphism Z such that $Z|_{E_{s,l}} = \text{Id}|_{E_{s,l}}$ and $Z|_{F_{s,l}} = -\text{Id}|_{F_{s,l}}$.*

c) Algebras from item (4) of Lemma 6.1 Note that $\mathfrak{l}(\mathcal{L}_m) \subset \mathfrak{n}(\mathcal{L}_m) \cong \mathbb{R} \text{Id}_{\mathcal{L}_m}$, but $\text{Id} \notin \mathfrak{sp}(\mathcal{L}_m)$. Therefore $\mathfrak{l}(\mathcal{L}_m) = 0$.

Combining results of items a), b), and c) above with Lemma 6.1 we obtain the Proposition 6.1.

□

2. Subspaces \mathfrak{a} , \mathfrak{z} , and \mathfrak{p} . If we consider the embedding $\varphi_{\delta_{\text{mod}}}(\mathfrak{sl}_2)$ of \mathfrak{sl}_2 into $\mathfrak{sp}(X)$ appearing in the decomposition (6.3), then from the basic representation theory of \mathfrak{sl}_2 it follows immediately that $\varphi_{\rho}(\mathfrak{sl}_2)$ has a rank 1 element if and only if the skew Young diagram of δ_{mod} does not satisfy condition (2) of Theorem 6.2, i.e. it consists of one row with two boxes. Note that this and the “only if” part of Proposition 6.1 implies the “only if” part of Theorem 6.2. However, this together with the “if” part of Proposition 6.1 is not enough to finish the proof of the “if” part of Theorem 6.2.

For this goal, we need a decomposition of $\mathfrak{r}(\mathbf{u}^F(\delta_{\text{mod}}))$, which is finer than (6.3). Let \mathfrak{a} be the subalgebra of $\mathfrak{r}(\mathbf{u}^F(\delta_{\text{mod}}))$ consisting of those elements of $\mathfrak{r}(\mathbf{u}^F(\delta_{\text{mod}}))$ for which any irreducible \mathfrak{sl}_2 -submodule of the \mathfrak{sl}_2 -module X is an invariant subspace. The embedding $\varphi_{\delta_{\text{mod}}}(\mathfrak{sl}_2)$ clearly belongs to \mathfrak{a} , which also implies that \mathfrak{a} is an \mathfrak{sl}_2 -submodule of $\mathfrak{r}(\mathbf{u}^F(\delta_{\text{mod}}))$. Therefore, there exists the unique \mathfrak{sl}_2 -module \mathfrak{p} which is complementary to \mathfrak{a} in $\mathfrak{r}(\mathbf{u}^F(\delta_{\text{mod}}))$,

$$(6.23) \quad \mathfrak{r}(\mathbf{u}^F(\delta_{\text{mod}})) = \mathfrak{a} \oplus \mathfrak{p}.$$

Note that by the construction \mathfrak{p} is a subspace of $\mathfrak{l}(X)$. Also, let

$$(6.24) \quad \mathfrak{z} = \mathfrak{a} \cap \mathfrak{l}(X).$$

Then

$$(6.25) \quad \mathfrak{l}(X) = \mathfrak{z} \oplus \mathfrak{p}, \quad \mathfrak{a} = \varphi_{\delta_{\text{mod}}}(\mathfrak{sl}_2) \oplus \mathfrak{z}.$$

Let us describe the introduced subspaces \mathfrak{a} , \mathfrak{z} , and \mathfrak{p} in more detail. For this fix, as before, a skew Young tableaux $\text{Tab}(\delta_{\text{mod}})$ of δ_{mod} . Assume that this tableaux consists of \tilde{k} rows (recall that $\tilde{k} = \text{rank } D - 1$) and let \tilde{L}_i , $1 \leq i \leq \tilde{k}$, be the subspaces of X spanned by the vectors appearing in the boxes of the i th row of $\text{Tab}(\delta_{\text{mod}})$. We have the following decomposition:

$$(6.26) \quad X = \bigoplus_{i=1}^{\tilde{k}} \tilde{L}_i,$$

$$(6.27) \quad \mathfrak{gl}(X) = \bigoplus_{i=1}^{\tilde{k}} \mathfrak{gl}(\tilde{L}_i) \oplus \bigoplus_{i \neq j} \text{Hom}(\tilde{L}_i, \tilde{L}_j).$$

Denote by π_1 and π_2 the projections from $\mathfrak{gl}(X)$ to $\bigoplus_{i=1}^{\tilde{k}} \mathfrak{gl}(\tilde{L}_i)$ and to $\bigoplus_{i \neq j} \text{Hom}(\tilde{L}_i, \tilde{L}_j)$, respectively, with respect to decomposition (6.27). Then by constructions

$$(6.28) \quad \mathfrak{z} = \pi_1(\mathfrak{l}(X)), \quad \mathfrak{p} = \pi_2(\mathfrak{l}(X)).$$

Now prove that if the skew Young diagram of δ_{mod} satisfies condition (1) of Theorem 6.2 and the algebra $\mathfrak{r}(\mathbf{u}^F(\delta_{\text{mod}}))$ contains an element of rank 1, then \mathfrak{a} must contain an element of rank 1. Indeed, if $\mathfrak{r}(\mathbf{u}^F(\delta_{\text{mod}}))$ contains an element of rank 1 with nonzero \mathfrak{p} component with respect to the decomposition 6.23 then, using the second relation of (6.28) and the fact that the space \mathfrak{p} is an invariant subspace of the projection from $\mathfrak{gl}(X)$ to $\mathfrak{gl}(\tilde{L}_i, \tilde{L}_j)$ with respect to the decomposition (6.27), the subspace \mathfrak{p} contains a rank 1 element, which contradicts Proposition 6.1.

Finally, assume that there exists an element $A \in \mathfrak{a}$ of rank 1. Since each subspace \tilde{L}_i is an invariant subspace of A then there exist $i_0, 1 \leq i_0 \leq \tilde{k}$ such that $A|_{\tilde{L}_{i_0}}$ has rank 1. Assume that A splits as $A_1 + A_2$ with respect to the second decomposition in (6.25) so that $A_1 \in \varphi_{\delta_{\text{mod}}}(\mathfrak{sl}_2)$ and $A_2 \in \mathfrak{z}$. On one hand, since \tilde{L}_{i_0} is an irreducible \mathfrak{sl}_2 -module, the spectrum of the endomorphism $A_1|_{\tilde{L}_{i_0}}$ constitute an arithmetic progression centered at (symmetric with respect to) the origin. On the other hand, $A_2|_{\tilde{L}_{i_0}}$ is either a multiple of the identity by the formula of type (6.22), if \tilde{L}_{i_0} is indexed by \mathbb{Z} (or, equivalently, \tilde{L}_{i_0} is a (Lagrangian) subspace of the domain of the indecomposable symbol of type $\mathbb{R}\delta_{s,l}$) or equal to zero by item c) of the proof of Proposition 6.1, if \tilde{L}_{i_0} is indexed by $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ (or, equivalently, \tilde{L}_{i_0} is the domain of the indecomposable symbol of type $\mathbb{R}\tau_m$). Therefore $A|_{\tilde{L}_{i_0}}$ may have rank 1 if and only if $\dim \tilde{L}_{i_0} \leq 2$.

If $\dim \tilde{L}_{i_0} = 1$ then the Young diagram of δ_{mod} contains two rows with one box each, which contradicts condition (1) of Theorem 6.2. If $\dim \tilde{L}_{i_0} = 2$, then there are two possibilities:

- (1) \tilde{L}_{i_0} is indexed by \mathbb{Z} . Then $\text{Diag}(\delta_{\text{mod}})$ contains 2 rows with 2 boxes which contradicts either Remark 4.1 or condition (1) of Theorem 6.2.
- (2) \tilde{L}_{i_0} is indexed by $\frac{1}{2}\mathbb{Z}_{\text{odd}}$. Let us prove that in this case $\text{Diag}(\delta_{\text{mod}})$ must consist of one row with two boxes (in contradiction to condition (2) of Theorem 6.2). Otherwise, the condition $A|_{\tilde{L}_i} = 0$ for all $i \neq i_0, 1 \leq i \leq \tilde{k}$ holds and it is not void. This implies that $A_1|_{\tilde{L}_i} = 0$ for all $i \neq i_0, 1 \leq i \leq \tilde{k}$, which yields that $A_1 = 0$. Besides, $A_2|_{\tilde{L}_i} = 0$ for all $i \neq i_0, 1 \leq i \leq \tilde{k}$ and by item c) of the part 1 of the current proof $A_2|_{\tilde{L}_{i_0}} = 0$. Therefore $A_2 = 0$ and hence $A = 0$, contradicting the assumption that A has rank 1.

This completes the proof of “if” part of Theorem 6.2. □

7. REDUCTION OF CALCULATION OF ALGEBRA $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ TO STANDARD PROLONGATION OF \mathfrak{p}

Next natural task is to describe the algebra $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ as explicit as possible.

First of all consider the case of rank 2 distributions, which is important as a building block for the case of distributions of arbitrary even rank.

Theorem 7.1. *A $(2, n)$ -distribution of maximal class has a modified Jacobi symbol δ_{mod} isomorphic to $\mathbb{R}\tau_{\frac{n-2}{2}}$, i.e. its Young diagram consists of one rows with $2(n-3)$ boxes, the algebra $\mathfrak{u}^F(\delta_{\text{mod}})$ is equal to an embedding of an irreducible representation of \mathfrak{gl}_2 into $\mathfrak{csp}(\eta^{-1})$ and the algebra $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ satisfies the following:*

- (1) [42] *If $n = 5$, then $\mathfrak{u}(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ is isomorphic to the split real form of the exceptional Lie algebra G_2 (as also expected from the classical Cartan work [9]);*

- (2) [15, 16] *If $n > 5$, then the first prolongation $\mathfrak{u}^1(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ of the pair $(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ vanishes, i.e. $\mathfrak{u}(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ is isomorphic to the semidirect sum of the above mentioned embedding of an irreducible representation of \mathfrak{gl}_2 into $\mathfrak{csp}(\eta^{-1})$ with η .*

It turns out that for more general Jacobi symbols with sufficiently long rows of their Young diagrams the subspace \mathfrak{p} introduced in the previous section (see the decomposition (6.23)) plays the crucial role in the description of the algebra $\mathfrak{u}(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$. First, note that since $\eta^{-1} (\cong X)$ generates η , for every $i \geq 0$ any element of i th Tanaka algebraic prolongation $\mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ of the pair $(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ is uniquely determined by its restriction to X .

Let $\phi \in \mathfrak{u}^1(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$. Then by definition for any $v_1, v_2 \in X \cong \eta^{-1}$ we have

$$(7.1) \quad \phi(v_1)v_2 - \phi(v_2)v_1 = \phi([v_1, v_2]).$$

Fix a symplectic form σ , representing the conformal symplectic structure on X . Then there exists the unique $y \in \eta^{-2}$ such that $[v_1, v_2] = \sigma(v_1, v_2)y$ for any v_1, v_2 in X . Let $v = \phi(y)$. Note that $v \in X$. Using this and relation (7.1), we obtain another characterization of $\mathfrak{u}^1(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$: $\phi \in \text{Hom}(X, \mathfrak{u}^F(\delta_{\text{mod}}) \oplus \text{Hom}(\eta^{-2}, X))$ belongs to $\mathfrak{u}^1(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ if and only if the restriction of ϕ to X satisfies the following condition: there exists $v \in X$ such that for any $v_1, v_2 \in X$

$$(7.2) \quad \phi(v_1)v_2 - \phi(v_2)v_1 = \sigma(v_1, v_2)v.$$

On the other hand, by definition $\phi \in \text{Hom}(X, \mathfrak{u}^F(\delta_{\text{mod}}))$ belongs to the standard prolongation $\mathfrak{u}^F(\delta_{\text{mod}})^{(1)}$ if and only if for any $v_1, v_2 \in X$

$$(7.3) \quad \phi(v_1)v_2 - \phi(v_2)v_1 = 0.$$

Relations (7.2) and (7.3) imply that the standard prolongation $(\mathfrak{u}^F(\delta_{\text{mod}}))^{(1)}$ can be considered as the subspace of the space $\mathfrak{u}^1(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ corresponding to $v = 0$ in (7.2). The same statement with the same proof is true for the higher order prolongations, i.e.

$$(7.4) \quad (\mathfrak{u}^F(\delta_{\text{mod}}))^{(i)} \subset \mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}})), \quad i > 0.$$

Theorem 7.2. *Assume that the skew Young diagram of the modified Jacobi symbols δ_{mod} satisfies both conditions (1) and (2) of Theorem 6.2 and in addition the following two conditions hold:*

- (3) *The number of boxes in every row indexed by \mathbb{Z} is at least 4;*

- (4) *The number of boxes in the row indexed by $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ (if it exists) is at least 6.*

Then for any $i > 0$ the i th Tanaka algebraic prolongation $\mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ of the pair $(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ coincides with the standard i th prolongation $\mathfrak{p}^{(i)}$ of the subspace \mathfrak{p} of $\mathfrak{gl}(X)$ defined in (6.28).

Remark 7.1. *Note that condition (3) in Theorem 7.2 is equivalent to the fact that in the decomposition of δ_{mod} into the indecomposable symbols all components of type $\mathbb{R}\delta_{s,l}$ satisfy $l \geq 3$ and condition (4) is equivalent to the fact that in the same decomposition the component $\mathbb{R}\tau_m$, if it*

exists, satisfies $m > \frac{3}{2}$. Also, conditions (1) and (3) are equivalent to condition (1) and the fact that in the decomposition of δ_{mod} into the indecomposable symbols there is no symbol of the type $\mathbb{R}\delta_{1,2}$, i.e. the indecomposable symbol with rectangular Young diagram of size 2×3 .

Before proving Theorem 7.2, note that since $\mathfrak{p} \subset \mathfrak{l}(X)$, one has that $\mathfrak{p}^{(i)} \subset (\mathfrak{l}(X))^{(i)}$. On the other hand, from (7.4) it follows that $(\mathfrak{l}(X))^{(i)} \subset \mathfrak{u}^i(\eta, \mathfrak{l}(X)) \subset \mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$, which implies the following

Corollary 7.1. *Under assumptions of Theorem 7.2 for any $i > 0$ the i th Tanaka algebraic prolongation $\mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ of the pair $(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ coincides with the standard i th prolongation $(\mathfrak{l}(X))^{(i)}$ of the algebra $\mathfrak{l}(X)$ of the infinitesimal symmetries of the flat curve $\mathcal{F}_{\delta_{\text{mod}}}$ with flag symbol δ_{mod} considered as a parametrized curve.*

Proof of Theorem 7.2 First, let us prove our theorem for $i = 1$, i.e. for the first prolongation.

Our theorem will be proved if we will show that for such ϕ we have that $\phi(v_1) \in \mathfrak{p}$ for any $v_1 \in X$ and that $v = 0$.

We can always assume that in the diagram $\text{Diag}(\delta_{\text{mod}})$ the Young subdiagram of the indecomposable component of type $\mathbb{R}\tau_m$, if exists, is situated at the very bottom. Fix again a skew Young tableaux $\text{Tab}(\delta_{\text{mod}})$ and let $r = \lfloor \frac{1}{2}(\text{rank } D - 1) \rfloor$. For any i such that $1 \leq i \leq r$ let E_i be the span of the vectors located in the boxes of the $(2i - 1)$ st row of $\text{Tab}(\delta_{\text{mod}})$ and F_i be the span of the vectors located in the boxes of the $2i$ th rows of $\text{Tab}(\delta_{\text{mod}})$. Finally, if $\text{rank } D$ is even, let \mathcal{L} be the span of the vectors located in the last box of $\text{Tab}(\delta_{\text{mod}})$.

Let us fix a basis in the subspace \mathfrak{z} defined by (6.24). For any i , $1 \leq i \leq r$, let $Z_i \in \mathfrak{csp}(X)$ such that $Z_i|_{E_i} = \text{Id}|_{E_i}$, $Z_i|_{F_i} = -\text{Id}|_{F_i}$, and Z_i equal to zero on all vectors of $\text{Tab}(\delta_{\text{mod}})$ that are not located in the boxes of the $(2i - 1)$ st and $2i$ th row of $\text{Tab}(\delta_{\text{mod}})$. Then using Remark 6.4 and item c) of the proof of Proposition 6.1 one can show that

$$(7.5) \quad \mathfrak{z} = \text{span}\{Z_i\}_{i=1}^r.$$

Now assume that $v_1, v_2 \in E_i$. Let $\overline{E_i}$ be the span of all vectors located in the boxes of $\text{Tab}(\delta_{\text{mod}})$ which are not in the $(2i - 1)$ st row and let Π_i be the projection from X to E_i parallel to $\overline{E_i}$. Let $\Pi_i \circ \mathfrak{u}^F(\delta_{\text{mod}}) = \{\Pi_i \circ A : A \in \mathfrak{u}^F(\delta_{\text{mod}})\}$. Consider the subspace $(\Pi_i \circ \mathfrak{u}^F(\delta_{\text{mod}}))|_{E_i}$ of $\mathfrak{gl}(E_i)$ such that it consists of the restrictions of elements of $\Pi_i \circ \mathfrak{u}^F(\delta_{\text{mod}})$ to E_i . Then from the second identity of 6.25 and Remark 6.4 it follows that the space $(\Pi_i \circ \mathfrak{u}^F(\delta_{\text{mod}}))|_{E_i}$ is 4-dimensional and it is generated by the restrictions of elements of $\varphi_{\delta_{\text{mod}}}$ to E_i and $\text{Id}|_{E_i}$, i.e. it is equal to the image of the irreducible embedding of $\mathfrak{gl}_2(\mathbb{R})$ into $\mathfrak{gl}(E_i)$.

As the restriction of the symplectic form to E_i vanishes, we get $\phi(v_1)v_2 = \phi(v_2)v_1$. Considering this equation mod $\overline{E_i}$ we get exactly the equation for the standard prolongation of $(\Pi_i \circ \mathfrak{u}^F(\delta_{\text{mod}}))|_{E_i} \cong \mathfrak{gl}_2(\mathbb{R})$. According to the result of Kobayashi–Nagano [27], the first prolongation of the irreducible embedding of $\mathfrak{gl}_2(\mathbb{R})$ is non-zero only if the dimension of the representation space does not exceed 3, which contradicts assumption (3) of our theorem. Thus, we see that $\phi(v_1)v_2 = 0 \pmod{\overline{E_i}}$ for all $v_1, v_2 \in E_i$. In particular, this implies that

$$(7.6) \quad \phi(E_i) \in \mathfrak{p} + \mathbb{R}(Z_i + \text{Id}) + \sum_{j \neq i} Z_j.$$

Absolutely analogous analysis of the cases $v_1, v_2 \in F_i$ implies that

$$(7.7) \quad \phi(F_i) \in \mathfrak{p} + \mathbb{R}(Z_i - \text{Id}) + \sum_{j \neq i} Z_j.$$

In the case when $v_1, v_2 \in \mathcal{L}$, using similar arguments, one can get from condition (4) of the theorem under consideration and item (2) of Theorem 7.1 that

$$(7.8) \quad \phi(\mathcal{L}) \in \mathfrak{p} + \sum_{j=1}^r Z_j$$

From relations (7.6)-(7.7) it follows that if $x \in X$ decomposes as $x = \sum_{j=1}^r (x_{E_j} + x_{F_j}) + x_{\mathcal{L}}$, where $x_{E_j} \in E_j$, $x_{F_j} \in F_j$, and $x_{\mathcal{L}} \in \mathcal{L}$, then there exist functionals $\alpha_i \in X^*$, $1 \leq i \leq r$, such that

$$(7.9) \quad \phi(x) = \phi'(x) + \sum_{i=1}^r (\alpha_i(x_{E_i})(Z_i - \text{Id}) + \alpha_i(x_{F_i})(Z_i + \text{Id})) + \sum_{i=1}^r \alpha_i \left(\sum_{j \neq i} (x_{E_j} + x_{F_j}) + x_{\mathcal{L}} \right) Z_i,$$

where ϕ' takes values in \mathfrak{p} . Now assume that $v_1 = v_{E_i} \in E_i$ and $v_2 = v_{F_i} \in F_i$. Then by (7.2) we get:

$$(7.10) \quad \phi(v_{E_i})v_{F_i} - \phi(v_{F_i})v_{E_i} = \sigma(v_{E_i}, v_{F_i})v.$$

Also let $\overline{E_i \oplus F_i} = \overline{E_i} \cap \overline{F_i}$. Consider equation (7.10) modulo $\overline{F_i}$.

- (1) *Assume that the subdiagram of $\text{Diag}(\delta_{\text{mod}})$ consisting of the $(2i-1)$ st and $2i$ th row is not rectangular. Then by (6.21), the second line there, the restriction of an element of \mathfrak{p} on E_i takes values in $\overline{E_i \oplus F_i}$. Taking into account this fact, the fact that an element of \mathfrak{p} sends F_i to $\overline{F_i}$, and relation (7.9), we get from (7.10) that*

$$-2\alpha_i(v_{E_i})v_{F_i} = \sigma(v_{E_i}, v_{F_i})v \pmod{\overline{F_i}}, \quad \text{for any } v_{E_i} \in E_i, v_{F_i} \in F_i.$$

As the dimensions of E_i and F_i are at least 4 by assumption (3) of the theorem, for any $v_{E_i} \in E_i$ we can find a non-zero vector $v_{F_i} \in F_i$ such that $\sigma(v_{E_i}, v_{F_i}) = 0$. Hence, we see that

$$(7.11) \quad \alpha_i|_{E_i} = 0.$$

In particular, we also see that $v \in \overline{F_i}$.

- (2) *Now assume that the subdiagram of $\text{Diag}(\delta_{\text{mod}})$ consisting of the $(2i-1)$ st and $2i$ th row is rectangular. The difference with the previous case is that the restriction of an element of \mathfrak{p} on E_i takes values in $\overline{E_i}$ but not in $\overline{E_i \oplus F_i}$ in general, because the considered case corresponds to the first line of (6.21). Assume that the considered rectangular diagram has size $2 \times (2s+1)$, the boxes of the upper row of the corresponding subtableaux are filled by vectors $e_s, e_{s-1}, \dots, e_{-s}$ from the left to the right, and the boxes of the lower row of the corresponding subtableaux are filled by vectors $f_s, f_{s-1}, \dots, f_{-s}$ from the left to the right. Recall that by constructions*

$$(7.12) \quad \sigma(e_j, e_k) = \sigma(f_j, f_k) = 0, \quad \sigma(e_j, f_{-j}) = (-1)^{s-j}$$

for all integers j, k such that $-s \leq j, k \leq s$. Then from the first line of (6.21) it follows that there exists $\beta_i \in F_i^*$ such that

$$(7.13) \quad \forall -s \leq k \leq s, v_{F_i} \in F_i, \quad \phi'(v_{F_i})e_k = \beta_i(v_{F_i})f_k \pmod{\overline{E_i \oplus F_i}}.$$

Taking into account this fact, the fact that an element of \mathfrak{p} sends F_i to $\overline{F_i}$, and the relation (7.9), we get after substitution of $v_{E_i} = e_k$ into (7.10) that:

$$(7.14) \quad -\beta_i(v_{F_i})f_k - 2\alpha_i(e_k)v_{F_i} = \sigma(e_k, v_{F_i})v \pmod{\overline{F_i}}, \quad \text{for any } v_{F_i} \in F_i, -s \leq k \leq s.$$

Taking $v_{F_i} = f_j$ with $j \neq -k$ and using (7.12) we obtain

$$(7.15) \quad -\beta_i(f_j)f_k - 2\alpha_i(e_k)f_j = 0 \pmod{\overline{F_i}}.$$

Since $s > 1$ by assumption (3) of Theorem 7.2, we can always take $j \neq -k$ such that in addition $j \neq k$. Using this j in (7.15) we get that $\alpha_i(e_k) = 0$. As the latter relation is valid for arbitrary k such that $-s \leq k \leq s$, the relation (7.11) holds in the considered case

as well. Moreover, combining (7.11) and (7.15) one also gets that $\beta_i = 0$, which together with (7.13) implies that

$$(7.16) \quad \text{Im}(\phi'(v_{F_i})|_{E_i}) = 0 \pmod{\overline{E_i \oplus F_i}}.$$

Let us now fix an arbitrary $v_{E_i} \in E_i$. From (7.9) and (7.11) we have $\phi(v_{E_i}) = \phi'(v_{E_i}) \pmod{\overline{E_i \oplus F_i}}$. Let us prove that $\phi'(v_{E_i}) = 0 \pmod{\overline{E_i \oplus F_i}}$. Indeed, by (7.9), (7.10), and (7.16), when the subdiagram of $\text{Diag}(\delta_{\text{mod}})$ consisting of the $(2i - 1)$ st and $2i$ th row is rectangular, we have

$$(7.17) \quad \phi'(v_{E_i})v_{F_i} = \sigma(v_{E_i}, v_{F_i})v + 2\alpha_i(v_{F_i})v_{E_i} \pmod{\overline{E_i \oplus F_i}}, \quad \text{for any } v_{F_i} \in F_i.$$

The splitting

$$(7.18) \quad X = \begin{cases} \left(\bigoplus_{i=1}^r (E_i \oplus F_i) \right) & \text{if } \text{rank } D = 2r + 1, \\ \left(\bigoplus_{i=1}^r (E_i \oplus F_i) \right) \oplus \mathcal{L} & \text{if } \text{rank } D = 2r + 2 \end{cases}$$

induces the natural splitting of $\mathfrak{gl}(X)$ as in (6.6). Let \mathfrak{P}_i be the projection from $\mathfrak{gl}(X)$ to $\mathfrak{gl}(F_i, E_i)$ with respect to the latter splitting. Equation (7.17) and the fact that $\phi'(v_{E_i})v_{F_i}$ takes values in $\overline{F_i}$ imply that the rank of the linear map $\mathfrak{P}_i(\phi'(v_{E_i}))$ is not greater than 2. On one hand, $\mathfrak{P}_i(\phi'(v_{E_i}))$ must belong to $\mathfrak{s}(E_i) \subset S^2(E_i)$ considered as a subspace of $\text{Hom}(F_i, E_i)$ as in item b) of the proof of Proposition 6.1.

On the other hand, from the condition (1) of Theorem 6.2 (which is also assumed in the formulation of the present theorem and which is equivalent to conditions (1) or (3) of Theorem 6.1 depending on $\text{rank } D$) and decomposition (6.20) it follows that the subspace $\mathfrak{s}(E_i)$ does not contain nonzero elements of rank not greater than 2. Below (Corollary to Lemma 8.6) we will give the proof of this using the description of the latter decomposition via geometry of rational normal curve. This and relation (7.17) imply that vectors v and v_{E_i} are collinear $\pmod{\overline{E_i \oplus F_i}}$. As v_{E_i} can be arbitrary and the dimension of the subspace E_i is at least 4, this implies that

$$(7.19) \quad v = 0 \pmod{\overline{E_i \oplus F_i}},$$

$$(7.20) \quad \alpha_i|_{F_i} = 0.$$

Since relation (7.19), holds for an arbitrary $i, 1 \leq i \leq r$, we have

$$(7.21) \quad v \in \mathcal{L}.$$

Now assume that $v_1, v_2 \in \mathcal{L}$ and let $\overline{\mathcal{L}}$ denotes the span of all vectors in $\text{Tab}(\delta_{\text{mod}})$ located in the boxes which are not in the last row of $\text{Tab}(\delta_{\text{mod}})$. Consider equation (7.2) modulo $\overline{\mathcal{L}}$. Using the definition of the space \mathfrak{p} , the fact that $Z_i|_{\mathcal{L}} = 0$, and the relation (7.9), one gets from (7.1) that

$$(7.22) \quad \sigma(v_1, v_2)v \in \overline{\mathcal{L}}.$$

Since $\sigma|_{\mathcal{L}} \neq 0$ (actually $\sigma|_{\mathcal{L}}$ is a symplectic form on \mathcal{L}), we get from (7.22) that $v \in \overline{\mathcal{L}}$, which together with (7.21) implies that

$$(7.23) \quad v = 0.$$

Further, let $v_1 = v_{E_{i_1}} \in E_{i_1}$ and $v_2 = v_{E_{i_2}} \in E_{i_2}$, where $i_1 \neq i_2$. Then from (7.1) and the fact that $\sigma(v_{E_{i_1}}, v_{E_{i_2}}) = 0$ it follows that $\phi(v_{E_{i_1}})v_{E_{i_2}} = \phi(v_{E_{i_2}})v_{E_{i_1}}$. Considering the latter equation modulo $\overline{E_{i_1}}$ and using (7.9) one gets that

$$(7.24) \quad \phi'(v_{E_{i_1}})v_{E_{i_2}} = \alpha_{i_1}(v_{E_{i_2}})v_{E_{i_1}} \pmod{\overline{E_{i_1}}}$$

As before, the splitting (7.18) induces the natural splitting of $\mathfrak{gl}(X)$ as in (6.6). Let \mathfrak{P}_{i_1, i_2} be the projection from $\mathfrak{gl}(X)$ to $\mathfrak{gl}(E_{i_2}, E_{i_1})$ with respect to the latter splitting. Equation (7.24) implies that the rank of the linear map $\mathfrak{P}_{i_1, i_2}(\phi'(v_{E_{i_1}}))$ is not greater than 1.

On one hand, $\mathfrak{P}_{i_1, i_2}(\phi'(v_{E_{i_1}}))$ must belong to $\mathfrak{n}(E_{i_2}, E_{i_1})$. On the other hand, from our assumptions it follows that $\mathfrak{n}(E_{i_2}, E_{i_1})$ does not contain elements of rank 1. This implies that $\mathfrak{P}_{i_1, i_2}(\phi'(v_{E_{i_1}})) = 0$ and thus by (7.24) $\alpha_{i_1}(v_{E_{i_2}})v_{E_{i_1}} = 0$. As $v_{E_{i_1}} \in E_{i_1}$ can be arbitrary this implies that

$$(7.25) \quad \alpha_{i_1}|_{E_{i_2}} = 0, \quad i_1 \neq i_2.$$

In completely analogous way (taking $v_1 \in E_{i_1}$ and $v_2 \in F_{i_2}$ or $v_1 \in E_i$ and $v_2 \in \mathcal{L}$) one can get that $\alpha_{i_1}|_{F_{i_2}} = 0$, $i_1 \neq i_2$ and $\alpha_i|_{\mathcal{L}} = 0$. Combining this with relations (7.11), (7.20), and (7.25) we get that

$$(7.26) \quad \alpha_i = 0, \quad 1 \leq i \leq r.$$

Therefore, by (7.9) $\phi = \phi'$, i.e. ϕ takes values in \mathfrak{p} . This together with (7.23) completes the proof of the Theorem 7.2 in the case of the first prolongation.

Now we are going to establish the induction step for proving our theorem for prolongations of higher order. For this first let us consider the following more abstract situation:

Let Y be a vector space endowed with a skew-symmetric (possibly degenerate) form ω . Similarly to symplectic case (i.e. when the form is degenerate) consider the graded nilpotent Lie algebra $\tilde{\eta} = \tilde{\eta}^{-1} \oplus \tilde{\eta}^{-2}$, where $\tilde{\eta}^{-1} = Y$, $\dim \tilde{\eta}^{-2} = 1$, $\tilde{\eta}^{-2}$ belongs to the center of $\tilde{\eta}$, and if a vector z generates $\tilde{\eta}^{-2}$ then $[y_1, y_2] = \omega(y_1, y_2)z$ for any $y_1, y_2 \in \tilde{\eta}^{-1}$.

Let W be a subspace in $\mathfrak{r}(\mathfrak{g}^0(\eta))$. As in Remark 1.1, W can be considered as a subspace of $\mathfrak{gl}(Y)$. Let $W^{(1)}$ be the standard prolongation of W and $W^{(1\text{sp})}$ be the first prolongation of the pair (η, W) (in our previous notation $W^{(1\text{sp})} = \mathfrak{u}^1(\eta, W)$). Note that both $W^{(1)}$ and $W^{(1\text{sp})}$ can be considered as subspaces of $\text{Hom}(Y, W)$ that in turn can be naturally identified with the subspace of $\mathfrak{gl}(W \times Y)$ as follows: that assigns to a map $\Phi \in \text{Hom}(Y, W)$ the map $\tilde{\Phi} \in \mathfrak{gl}(Y')$ as follows:

$$(7.27) \quad \tilde{\Phi}(w, y) := (\Phi(y), 0).$$

Further, assume that the space Y decomposes as follows

$$(7.28) \quad Y = \bigoplus_{i=1}^{\tilde{k}} Y_i,$$

where subspaces Y_i satisfy the following properties:

- (Y1) For any integer j such that $1 \leq j \leq \lfloor \frac{\tilde{k}}{2} \rfloor$ there exist vectors $v_1 \in Y_{2j-1}$ and $v_2 \in Y_{2j}$ such that $\omega(v_1, v_2)v \neq 0$;
- (Y2) In the case of odd \tilde{k} the restriction of the form ω to the subspace $Y_{\tilde{k}}$ is nonzero identically zero.

As before, splitting (7.28) of Y induces the splitting $\mathfrak{gl}(Y) = \bigoplus_{i_1, i_2=1}^{\tilde{k}} \text{Hom}(Y_{i_1}, Y_{i_2})$. Let $\pi_{i_1, i_2} : \mathfrak{gl}(Y) \rightarrow \text{Hom}(Y_{i_1}, Y_{i_2})$ the canonical projection with respect to this splitting.

Lemma 7.1. *With all notations as above, assume that the space W satisfies the following two properties:*

- (W1) $W \subset \bigoplus_{i_1 \neq i_2} \text{Hom}(Y_{i_1}, Y_{i_2})$;

(W2) For every $i_1, i_2 \in \{1, \dots, \tilde{k}\}$ which are distinct (i.e. $i_1 \neq i_2$) the subspace $\pi_{i_1, i_2}(W)$ of $\text{Hom}(Y_{i_1}, Y_{i_2})$ does not contain elements of rank 1.

Then the following two statements hold:

- (1) $W^{(1\text{sp})} = W^{(1)}$
- (2) Consider the space

$$Y' = \left(\bigoplus_{i_1 \neq i_2} \pi_{i_1, i_2}(W) \right)$$

endowed with the symplectic form ω' which is the pullback of the form ω with respect to the canonical projection from Y' to Y (note that $W \times Y \subset Y'$ by condition (W1)). Take the following splitting of Y' :

$$(7.29) \quad Y' = \bigoplus_{i=1}^{\tilde{k}} Y'_i,$$

where

$$(7.30) \quad Y'_i = \left(\bigoplus_{\{j: j \neq i, 1 \leq j \leq \tilde{k}\}} \pi_{j, i}(W) \right) \times Y_i.$$

Then the space Y' satisfies conditions (Y1) and (Y2) and the space $W' = W^{(1\text{sp})} = W^{(1)}$, considered as the subspace of $\mathfrak{gl}(Y')$ (under the natural identification as in (7.27)), satisfies conditions (W1) and (W2) with respect to the splitting (7.29).

Proof. Take $\Phi \in W^{(1\text{sp})}$. Similarly to (7.2), there exists a vector $v \in Y$ such that

$$(7.31) \quad \Phi(v_1)v_2 - \Phi(v_2)v_1 = \omega(v_1, v_2)v, \quad \forall v_1, v_2 \in Y$$

The first part of the lemma will be proved if we will show that $v = 0$.

Similarly to the previous considerations, let $\overline{Y}_i = \bigoplus_{j \neq i} Y_j$ and $\overline{Y}_{i_1} \oplus \overline{Y}_{i_2} = \bigoplus_{j \notin \{i_1, i_2\}} Y_j = \overline{Y}_{i_1} \cap \overline{Y}_{i_2}$.

Fix integer j such that $1 \leq j \leq \left\lfloor \frac{\tilde{k}}{2} \right\rfloor$ and take $v_1 \in Y_{2j-1}$, $v_2 \in Y_{2j}$. By property (W1) $\Phi(v_2)v_1 \in \overline{Y}_{2j-1}$. Therefore, from (7.31) it follows that $\Phi(v_1)v_2 = \omega(v_1, v_2)v \pmod{\overline{Y}_{2j-1}}$. Hence, the linear map $\Phi(v_1)v_2$ has rank not greater than 1. From this and assumption (W2) it follows that $\pi_{2j, 2j-1}(\Phi(v_1)) = 0$, i.e. $\omega(v_1, v_2)v \in \overline{Y}_{2j-1}$ for any $v_1 \in Y_{2j-1}$, $v_2 \in Y_{2j}$. By assumptions (Y1) there exists $v_1 \in Y_{2j-1}$ and $v_2 \in Y_{2j}$ such that $\omega(v_1, v_2)v \neq 0$, which implies that

$$(7.32) \quad v \in \overline{Y}_{2j-1}.$$

By the same arguments, exchanging the role of v_1 and v_2 , one gets that

$$(7.33) \quad v \in \overline{Y}_{2j}.$$

As $j \in \{1, \dots, \left\lfloor \frac{\tilde{k}}{2} \right\rfloor\}$ is arbitrary, we get from (7.32) and (7.33) that $v = 0$ if \tilde{k} is even or

$$(7.34) \quad v \in Y_{\tilde{k}} \text{ if } \tilde{k} \text{ is odd.}$$

In the latter case take v_1 and v_2 in $Y_{\tilde{k}}$. Then by assumption (W1) $\Phi(v_1)v_2 - \Phi(v_2)v_1 \in \overline{Y}_{\tilde{k}}$. Hence, using (7.31), we get that $\omega(v_1, v_2)v \in \overline{Y}_{\tilde{k}}$. By assumption (W2) there exist vectors v_1 and

v_2 in $Y_{\tilde{k}}$ such that $\omega(v_1, v_2) \neq 0$, which implies that $v \in \overline{Y_{\tilde{k}}}$. Combining this with (7.34) we get that $v = 0$ for odd \tilde{k} as well. This proves the first part of the lemma.

Now let us prove the second part of the lemma. First it is clear from definition of the form ω' that the splitting (7.29) satisfies conditions (Y1) and (Y2).

Let us prove that the space W' satisfies condition (W1) with respect to the splitting (7.29). Let $\pi'_{i_1, i_2} : \mathfrak{gl}(Y') \mapsto \text{Hom}(Y'_{i_1}, Y'_{i_2})$ be the canonical projection with respect to the splitting of $\mathfrak{gl}(Y')$ induced by the splitting (7.29) of the space Y' . We actually have to prove that $\pi'_{i, i}(\Psi) = 0$ for any $\Psi \in W'$ and any integer i , $1 \leq i \leq \tilde{k}$.

If $\pi_i : Y \mapsto Y_i$ denote the canonical projection with respect to the splitting (7.28), then directly from definition (7.30) of subspaces Y'_i for any $\Psi \in \text{Hom}(Y, W) \subset \mathfrak{gl}(Y')$ we have the following relation:

$$(7.35) \quad \pi'_{i_1, i_2}(\Psi)(y) = \pi_{i_2} \circ (\Psi(\pi_{i_1} y)), \quad \forall y \in Y.$$

In the righthand side of the last equation \circ means the composition of two linear maps: $\Psi(\pi_{i_1} y) \in W \subset \mathfrak{gl}(Y)$ and $\pi_{i_2} \in \text{Hom}(Y, Y_{i_2})$.

If $\Psi \in W'$ ($= W^{(1)}$), then $\Psi(y)z = \Psi(z)y$ for any $y, z \in Y$. Therefore, by (7.35)

$$(\pi_{i, i}\Psi)(y)z = \pi_i(\Psi(\pi_i y)z) = \pi_i(\Psi(z)\pi_i y).$$

Since $\Psi(z) \in W$ and W satisfies condition (W1), then $\Psi(z)\pi_i y \in \overline{Y_i}$, which implies that $\pi_i(\Psi(z)\pi_i y) = 0$. So, $\pi_{i, i}\Psi = 0$, hence the space W' satisfies condition (W1).

Finally, let us prove that the space W' satisfies condition (W1) with respect to the splitting (7.29). Assume by contradiction that there exists $\Psi \in W'$ and indices $i_1 \neq i_2$, $1 \leq i_1, i_2 \leq \tilde{k}$ such that the map $\pi'_{i_1, i_2}(\Psi)$ has rank 1. It means that there exist $w_0 \in \mathfrak{gl}(Y)$ and $\alpha \in Y_{i_1}^*$, $\alpha \neq 0$ such that

$$(7.36) \quad \pi'_{i_1, i_2}(\Psi)(y) = \alpha(\pi_{i_1} y)w_0.$$

Take $y_1, y_2 \in Y_{i_1}$. Then from (7.35) it follows that

$$(7.37) \quad \pi'_{i_1, i_2}(\Psi)(y_1)y_2 = \pi_{i_2}\Psi(y_1)y_2 = \alpha(y_1)w_0(y_2).$$

In the same way,

$$\pi'_{i_1, i_2}(\Psi)(y_2)y_1 = \alpha(y_2)w_0(y_1).$$

Since $\Psi \in W' = W^{(1)}$,

$$\pi'_{i_1, i_2}(\Psi)(y_1)y_2 = \pi_{i_2}\Psi(y_1)y_2 = \pi_{i_2}\Psi(y_2)y_1 = \pi'_{i_1, i_2}(\Psi)(y_2)y_1.$$

Therefore, $\alpha(y_1)w_0(y_2) = \alpha(y_2)w_0(y_1)$

Since by our assumptions $\alpha \neq 0$, there exists $y_2 \in Y_{i_1}$ such that $\alpha(y_2) \neq 0$, which implies that

$$(7.38) \quad w_0(y) = \frac{\alpha(y)}{\alpha(y_2)}w_0(y_2), \quad \forall y \in Y_{i_1}$$

So, the map w_0 has rank 1. Finally, (7.37) implies that $w_0(y_2) \in Y_{i_2}$, which yields that $w_0 \in \pi_{i_1, i_2}(W)$. This means that the space $\pi_{i_1, i_2}(W)$ contains elements of rank 1, contradicting assumption (W2) on W . \square

Now let us explain how to use Lemma 7.1 as an induction step in proving of our Theorem 7.2 for higher order prolongations. Assume as before that $\eta = \eta^{-1} \oplus \eta^{-2}$ be the Heisenberg algebra, the algebra $\mathfrak{g}^0 \subset \mathfrak{osp}(\eta^{-1})$ and for a given $i \in \mathbb{N}$

$$Y = \eta^{-1} \oplus \mathfrak{g}^0 \oplus \bigoplus_{j=1}^{i-1} \mathfrak{u}^j(\eta, \mathfrak{g}^0).$$

Let ω be the skew-symmetric form on Y , which is the pullback of a symplectic form on η^{-1} (defined up to a multiplication by a constant). Take $W = \mathfrak{u}^i(\eta, \mathfrak{g}^0)$. Note that W can be considered as a subspace of $\text{Hom}(\eta^{-1}, \mathfrak{u}^{i-1}(\eta, \mathfrak{g}^0))$, which in turn can be identified with a subspace of $\mathfrak{gl}(Y)$. Directly from the definitions of prolongations one can show that under this identification

$$(7.39) \quad W^{(1\text{sp})} \cong \mathfrak{u}^{i+1}(\eta, \mathfrak{g}^0).$$

Here again $\mathfrak{u}^{i+1}(\eta, \mathfrak{g}^0)$ is identified with a subspace in $\mathfrak{gl}(Y \oplus \mathfrak{u}^i(\eta, \mathfrak{g}^0))$.

To finish the proof of Theorem 7.2 we proceed by induction. We proved previously that $\mathfrak{u}^1(\eta, \mathfrak{g}^0) = \mathfrak{p}^1$. Also, the algebra \mathfrak{p} satisfies conditions of Lemma 7.1. Hence, $\mathfrak{p}^{(1\text{sp})} = \mathfrak{p}^{(1)}$. Moreover, using the fact that $(\mathfrak{p}^{(i)})^{(1)} = \mathfrak{p}^{(i+1)}$ and applying the same lemma inductively we get that $\mathfrak{p}^{(i+1)} = (\mathfrak{p}^{(i)})^{(1\text{sp})}$. Assuming by induction that $\mathfrak{u}^i(\eta, \mathfrak{g}^0) = \mathfrak{p}^{(i)}$ for some $i \in \mathbb{N}$ and using (7.39), we get that

$$\mathfrak{u}^{i+1}(\eta, \mathfrak{g}^0) = (\mathfrak{u}^i(\eta, \mathfrak{g}^0))^{(1\text{sp})} = (\mathfrak{p}^{(i)})^{(1\text{sp})} = \mathfrak{p}^{(i+1)},$$

which completes the induction step and therefore the proof of the theorem. \square

8. DESCRIPTION OF ALGEBRA $\mathfrak{u}(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ VIA GEOMETRY OF CERTAIN PROJECTIVE VARIETIES

In this section we make an attempt given a flag symbol satisfying conditions of Theorem 7.2 to describe the algebra $\mathfrak{u}(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ in the more explicitly way using the fact that if a subspace in $\mathfrak{sp}(X)$ can be identified with a space of quadratic polynomials vanishing on a given projective variety \mathcal{E} in $\mathbb{P}X$, then the standard prolongations of this subspace can be identified with the space of certain polynomials vanishing on the secant variety of \mathcal{E} of certain order. For general Jacobi symbols in this way we are able only to show that $\mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ is a subspace in a space of this kind, obtaining an upper bound for the dimension of $\mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$. However, for a large class of Jacobi symbols the spaces $\mathfrak{u}^i(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ can be identified with certain spaces of polynomials of this kind (and not only with certain subspaces of them). In this way we get the description of the prolongation algebra $\mathfrak{u}(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ for all Jacobi symbols appearing in rank 3 distributions, for which this algebra is finite dimensional, and also for most of such Jacobi symbols appearing in rank 4 distributions (see Remark 8.3 below). In particular, in the case of rank 3 distributions with Jacobi symbol having nonrectangular skew Young diagram this description is given in terms of tangential developable and secant varieties of a rational normal curve.

To begin with, to any $A \in \text{Hom}(X, Y)$ one can naturally assign a bilinear form \tilde{A} on $X \times Y^*$ as follows: $\tilde{A}(v, p) := p(Av)$, $v \in X, p \in Y^*$. Given a subspace W in $\text{Hom}(X, Y)$ denote by \tilde{W} the corresponding subspace in the space of bilinear forms on $X \times Y^*$. Let r and s are dimensions of X and Y respectively and $x = (x_1, \dots, x_r)$ and $y = (y_1, \dots, y_s)$ are coordinates on X and Y^* w.r.t. some bases in those vector spaces. Then any bilinear form is identified with a polynomial in x and y , which is linear in x and linear in y . The following lemma is well known and immediately follows from definitions:

Lemma 8.1. *The i th prolongation of a subspace W of $\text{Hom}(X, Y)$ can be naturally identified with all polynomials $F(x, y)$ of degree $i + 1$ in x and linear in y such that all their partial derivatives in x of order i (i.e. polynomials $\frac{\partial^n F}{\partial x_1^{\alpha_1} \dots \partial x_r^{\alpha_r}}$ for all multi-indices $\alpha = (\alpha_1, \dots, \alpha_r)$ with $|\alpha| = i$) belong to the space \widetilde{W} .*

Now assume that the space X is endowed with a symplectic form σ . The form σ defines the identification between X and X^* given by the map $v \mapsto \sigma(v, \cdot)$, $x \in X$. Under this identification given an endomorphism A on X the form \widetilde{A} is a bilinear form on $X \times X$. An endomorphism A of X belongs to $\mathfrak{sp}(X)$ if and only if the corresponding bilinear form \widetilde{A} is symmetric. In this case \widetilde{A} is completely determined by the corresponding quadratic form

$$(8.1) \quad v \mapsto \widetilde{A}(v, v) = \sigma(v, Av).$$

Therefore a subspace W of $\mathfrak{sp}(X)$ can be identified with a subspace of quadratic forms on X which by analogy with the previous considerations will be denoted by \widetilde{W} . Then in this situation Lemma 8.1 can be reformulated as follows:

Lemma 8.2. *The k th prolongation of a subspace W of $\mathfrak{sp}(X)$ can be naturally identified with the space of all polynomials of degree $k + 2$ in X such that all their partial derivatives of order k belong to the corresponding subspace \widetilde{W} in the space of quadratic forms.*

Given a projective variety \mathcal{E} denote by $\mathcal{I}(\mathcal{E})$ the ideal of homogeneous polynomials vanishing on \mathcal{E} . We shall also denote by $\mathcal{I}_k(\mathcal{E})$ the subspace of all polynomials of degree k in $\mathcal{I}(\mathcal{E})$. Define the k th secant variety $\mathcal{S}^k \mathcal{E}$ of \mathcal{E} as the algebraic closure of the union of k -planes in the projective space passing through $k + 1$ points from \mathcal{E} . By definition we set $\mathcal{S}^0(\mathcal{E}) = \mathcal{E}$.

Lemma 8.3. *Assume that a subspace W of $\mathfrak{sp}(X)$ satisfies the following property: there exists a projective variety \mathcal{E} in X such that the corresponding subspace \widetilde{W} belongs to $\mathcal{I}_2(\mathcal{E})$. Then the k th standard prolongation of W can be identified with a subspace of $\mathcal{I}_{k+2}(\mathcal{S}^k \mathcal{E})$. Moreover, if $\widetilde{W} = \mathcal{I}_2(\mathcal{E})$, then $W^{(k)} \cong \mathcal{I}_{k+2}(\mathcal{S}^k \mathcal{E})$.*

Proof. By Lemma 8.2 the k th prolongation of W can be identified with polynomials F of degree $k + 2$ such that all their partial derivatives of order k belong to $\mathcal{I}_2(\mathcal{E})$. It is clear that any such polynomial and all its partial derivatives of degree $\leq k$ lie in $\mathcal{I}(\mathcal{E})$.

Let us prove that any such polynomial F vanishes identically at $\mathcal{S}^k(\mathcal{E})$. It is sufficient to prove that it vanishes at any secant k -plane of \mathcal{E} . Indeed, let $p_0, \dots, p_k \in \mathcal{E}$ be the set of $k + 1$ points and let

$$W: \mathbb{P}^k \rightarrow \mathbb{P}^{r-1}, \quad [y_0 : y_1 : \dots : y_k] \mapsto y_0 p_0 + y_1 p_1 + \dots + y_k p_k;$$

be the embedding of the corresponding secant k -plane, where $r = \dim X$. Then $F \circ W$ is a polynomial of degree $k + 2$ on \mathbb{P}^k that vanishes at basis points $q_i \in \mathbb{P}^k$, $i = 0, \dots, k$, and, in addition, all its derivatives of degree $\leq k$ also satisfy this property. This immediately implies that $F \circ W = 0$. Hence, F vanishes identically at $\mathcal{S}^k(\mathcal{E})$, i.e. it belongs to $\mathcal{I}_{k+2}(\mathcal{S}^k \mathcal{E})$.

Finally, if $\widetilde{W} = \mathcal{I}_2(\mathcal{E})$ by make the arguments in the reversed order one gets that if a polynomials F of degree $k + 2$ vanishes on $\mathcal{S}^k \mathcal{E}$ then all its derivatives of order k vanishes on \mathcal{E} and hence F belongs to $W^{(k)}$. □

The question now is what varieties can be taken as \mathcal{E} for subspaces \mathfrak{p} or $\mathfrak{l}(X)$? Throughout the rest of the section we assume that the symbol δ_{mod} satisfies all assumptions of Theorem (7.2).

As in the beginning of the proof of Theorem (7.2) assume that in the diagram $\text{Diag}(\delta_{\text{mod}})$ the Young subdiagram of the indecomposable component of type $\mathbb{R}\tau_m$, if exists, is situated at the very bottom. Fix again a skew Young tableaux $\text{Tab}(\delta_{\text{mod}})$ and let $r = \lceil \frac{1}{2}(\text{rank } D - 1) \rceil$. For any i such that $1 \leq i \leq r$ let E_i be the span of the vectors located in the boxes of the $(2i - 1)$ st row of $\text{Tab}(\delta_{\text{mod}})$ and F_i be the span of the vectors located in the boxes of the $2i$ th rows of $\text{Tab}(\delta_{\text{mod}})$. Finally, if $\text{rank } D$ is even, let \mathcal{L} be the span of the vectors located in the last box of $\text{Tab}(\delta_{\text{mod}})$.

Let j_0 be the minimal index in I (with respect to the natural order) such that the subspace X_{j_0} is isotropic. Note that in fact j_0 is the minimal positive index in I , i.e. $j_0 = 1$ if $I = \mathbb{Z}$ and $j_0 = \frac{1}{2}$ if $I = \frac{1}{2}\mathbb{Z}_{\text{odd}}$ or $I = \frac{1}{2}\mathbb{Z}$. Let \mathcal{A} be a subset $\{1, \dots, r\}$ of all i such that the subdiagram of $\text{Diag}(\delta_{\text{mod}})$ consisting of the $(2i - 1)$ st and $2i$ th row is not rectangular. Assume that the restriction of the symbol δ_{mod} to $E_i \oplus F_i$ is isomorphic to the symbol $\mathbb{R}\delta_{s,l}$ and fix the skew Young tableaux for this restriction as in (4.3). Given $i \in \mathcal{A}$ let \mathcal{L}_i be the linear span of X_{j_0} and the vector e_0 from the skew Young tableaux (4.3) corresponding to of the symbol δ_{mod} to $E_i \oplus F_i$. If $i \notin \mathcal{A}$ we set $\mathcal{L}_i = X_{j_0}$.

Let $\mathcal{E}_i(\delta_{\text{mod}})$ be the algebraic closure of the curve $x \mapsto e^x \mathcal{L}_i(\delta_{\text{mod}})$, $x \in \delta_{\text{mod}}$. $\mathcal{E}_i(\delta_{\text{mod}})$ can be considered as algebraic varieties in the projective space $\mathbb{P}X$ (by taking the closure of the union of all lines belonging to spaces $e^x \mathcal{L}_i(\delta_{\text{mod}})$ for all $x \in \delta_{\text{mod}}$).

Lemma 8.4. *If $A \in \mathfrak{p}$, then the corresponding quadratic form \tilde{A} vanishes on $\mathcal{E}_i(\delta_{\text{mod}})$ for every i such that $1 \leq i \leq r$, if $\text{rank } D$ is odd, and $1 \leq i \leq r + 1$, if $\text{rank } D$ is even. In other words,*

$$(8.2) \quad \tilde{\mathfrak{p}} \subset \mathcal{I}_2(\mathcal{E}_i(\delta_{\text{mod}})).$$

Proof. If $A \in \mathfrak{p}$ then (6.4) implies that $A(e^x(X_j)) \subset e^x(X_j) \forall x \in \delta_{\text{mod}}$. Since by the definition of j_0 all subspaces $e^x(X_{j_0})$ are isotropic, from this and (8.1) it follows that the quadratic form \tilde{A} vanishes on $e^x(X_{j_0})$. This completes the proof in case $i \notin \mathcal{A}$.

Now assume that $i \in \mathcal{A}$. Since $e^t x \mathcal{L}_i \subset (e^x(X_{j_0}))^\perp$, then $\tilde{A}(v_1, v_2) = 0$ for any $v_1 \in e^x(X_{j_0})$ and $v_2 \in e^t x \mathcal{L}_i$. Finally, in case by (6.21), the second line there, Ae_0 belongs to $\overline{E_i \oplus F_i} \subset E_i^\perp$, where e_0 is as in (4.3), considered as the skew Young tableaux for the restriction of the symbol δ_{mod} to $E_i \oplus F_i$. This implies that the quadratic form \tilde{A} vanishes on $e^x \mathcal{L}_i$ and completes the proof of the lemma. \square

As a direct consequence of Corollary 7.1 and Lemmas 8.3 and 8.4, we get the following

Theorem 8.1. *Assume that the flag symbol δ_{mod} satisfies all assumptions of Theorem 7.2. Then for every i such that $1 \leq i \leq r$, if $\text{rank } D$ is odd, and $1 \leq i \leq r + 1$, if $\text{rank } D$ is even and for any positive integer k the k th Tanaka algebraic prolongation $\mathfrak{u}^k(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ of the pair $(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ can be identified with a subspace of the space $\mathcal{I}_{k+2}(S^k \mathcal{E}_i(\delta_{\text{mod}}))$ of all polynomials of degree $k + 2$ on X that vanishes on the k th secant variety of the variety $\mathcal{E}_i(\delta_{\text{mod}})$.*

Note that under assumptions of this theorem on the flag symbol δ_{mod} certain osculating subspace of the curve $x \mapsto e^x X_{j_0}$, $x \in \delta_{\text{mod}}$, is equal to the whole ambient vector space X at any point. Besides, by construction $X_{j_0} \subseteq \mathcal{L}_i$ for any i . Therefore, $S^K \mathcal{E}_i(\delta_{\text{mod}}) = \mathbb{P}X$ for sufficiently large K and by Theorem 8.1 the spaces $\mathfrak{u}^K(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ vanishes for such K . This gives more constructive proof of the finiteness type result for distributions with Jacobi symbols satisfying assumptions of Theorem 8.1 without referring to the Spencer criterium. In addition, this gives some estimate from above on the size of the spaces $\mathfrak{u}^k(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$.

The space $\mathcal{I}_2(\mathcal{E}_i(\delta_{\text{mod}}))$ is usually larger than the space \mathfrak{p} , so Theorem 8.1 gives only an estimation from above for the size of $\mathbf{u}^k(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$. If the quadratic form \tilde{A} belongs to $\mathcal{I}_2(\mathcal{E}_i(\delta_{\text{mod}}))$, then by polarization the corresponding bilinear form \tilde{A} satisfies $\tilde{A}(v_1, v_2) = 0$ for every $v_1, v_2 \in e^x(\mathcal{L}_i)$ and $x \in \delta_{\text{mod}}$. Hence, if $A \in \mathfrak{sp}(X)$ is related to the form \tilde{A} by (8.1), then $\sigma(v_1, Av_2) = 0$ for every $v_1, v_2 \in e^x(\mathcal{L}_i)$, which is equivalent to the following inclusion

$$(8.3) \quad A(e^x \mathcal{L}_i) \subset \left(e^x \mathcal{L}_i \right)^\angle, \quad \forall x \in \delta_{\text{mod}}$$

If $i \in \mathcal{A}$, then $\left(e^x(\mathcal{L}_i) \right)^\angle$ is equal to the linear span of all element of nonnegative degree in $\overline{E_i \oplus F_i}$ and vector e_1 , while if $i \notin \mathcal{A}$ then $\left(e^x(\mathcal{L}_i) \right)^\angle$ is equal to the linear span of element of nonnegative degrees in X . So, condition (8.3) does not ensure that any $A \in \mathcal{I}_2(\mathcal{E}_i(\delta_{\text{mod}}))$ is concentrated in nonnegative degrees. It is guarantied only in the case $r = 0$ (corresponding to rank 2 distributions of maximal class), because in this case the subspaces X_{j_0} is Lagrangian, but this case is not interesting since here all subspaces involved such as \mathfrak{p} , $\mathfrak{l}(X)$, and $\mathcal{I}_2(\mathcal{E}_i(\delta_{\text{mod}}))$ are equal to zero, as follows from above arguments and part c) of the proof of Lemma 6.1.

Now we give a class of flag symbols δ_{mod} for which the spaces $\mathbf{u}^i(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ can be identified with spaces of polynomials of this kind (and not only with subspaces of them). Continuing the numeration of assumptions Theorems 6.2 and 7.2 assume that

- (5) In a decomposition of the symbol δ_{mod} into indecomposable components there is no indecomposable symbols of the type $\mathbb{R}\delta_{s,2s}$, i.e. symbols with rectangular Young diagram consisting of two rows. Equivalently, the introduced above set \mathcal{A} coincides with $\{1, \dots, r\}$.
- (6) For any two indecomposable symbols $\mathbb{R}\delta_{s_1, l_2}$ and $\mathbb{R}\delta_{s_2, l_2}$ in the decomposition of the symbol δ_{mod} into indecomposable components we have

$$(8.4) \quad s_1 + s_2 > \max\{l_1, l_2\}.$$

Remark 8.1. Note that assumption (6) formally implies assumption (5) if one takes the same indecomposable component as $\mathbb{R}\delta_{s_1, l_2}$ and $\mathbb{R}\delta_{s_2, l_2}$. However, we would like to emphasis assumption (5) as a separate one.

The key point of assumption (5) is that under this assumption

$$(8.5) \quad \mathfrak{s}(F_i) = 0, \quad \forall 1 \leq i \leq r,$$

where, as in the part b) of the proof of Lemma 6.1, $\mathfrak{s}(F_i)$ is the maximal \mathfrak{sl}_2 -submodule of $S^2(F_{s,l})$ concentrated in the non-negative degree part of $S^2(F_{s,l})$, considered as the subspace of $\text{Hom}(E_i, F_i)$ under the identification of E_i^* with F_i via the symplectic form (see also (6.21), second line).

The key point of assumption (6) is that

$$(8.6) \quad \mathfrak{n}(E_{i_1}, F_{i_2}) = 0, \quad \forall 1 \leq i_1 \neq i_2 \leq r,$$

where as in the part b) of the proof of Lemma 6.1, $\mathfrak{n}(E_{i_1}, F_{i_2})$ is the maximal \mathfrak{sl}_2 -submodule of $\text{Hom}(E_{i_1}, F_{i_2})$ concentrated in the non-negative degree part of $\text{Hom}(E_{i_1}, F_{i_2})$. Identity (8.6) can be obtain easily from (6.13).

Further, let \mathcal{L} be equal to the linear span of all spaces \mathcal{L}_i with $1 \leq i \leq r$, if rank D is odd, and $1 \leq i \leq r + 1$, if rank D is even.

Remark 8.2. Note that under assumption (5) the space \mathcal{L} is Lagrangian.

Now let $\mathcal{E}(\delta_{\text{mod}})$ be the algebraic closure of the curve $x \mapsto e^x \mathfrak{L}(\delta_{\text{mod}})$, $x \in \delta_{\text{mod}}$. Again $\mathcal{E}(\delta_{\text{mod}})$ can be considered as algebraic varieties in the projective space $\mathbb{P}X$ (by taking the closure of the union of all lines belonging to spaces $e^x \mathfrak{L}(\delta_{\text{mod}})$ for all $x \in \delta_{\text{mod}}$).

Lemma 8.5. *Assume that Jacobi symbols satisfies conditions of Theorem 6.2 and in addition satisfies assumptions (5) and (6) above. If $A \in \mathfrak{p}$, then the corresponding quadratic form \tilde{A} vanishes on $\mathcal{E}(\delta_{\text{mod}})$. In other words,*

$$(8.7) \quad \tilde{\mathfrak{p}} \subset \mathcal{I}_2(\mathcal{E}(\delta_{\text{mod}})).$$

Proof. We proceed exactly as in the proof of Lemma 8.4. The only difference is in the final step: here we need to show that $\sigma(v, Av) = 0$ for any $v \in \mathfrak{L}$ which has degree 0. Indeed, from (8.5) and (8.6) it follows that $Av \in \mathfrak{L} \cap \bigoplus_{i=1}^r E_i$. Then by Remark 8.2 $\sigma(Av, v) = 0$. \square

Further, using the same arguments as ones leading to relation (8.3) we get that $\tilde{A} \in \mathcal{I}_2(\mathcal{E}(\delta_{\text{mod}}))$ if and only if

$$(8.8) \quad A(e^x \mathfrak{L}) \subset (e^x \mathfrak{L})^\perp = e^x \mathfrak{L}, \quad \forall x \in \delta_{\text{mod}}.$$

The last equality follows again from the fact that \mathfrak{L} is Lagrangian. The next important observation is that if the symbol δ_{mod} satisfies assumption (1) of Theorem 6.2 then the whole flat curve $\mathcal{F}_{\delta_{\text{mod}}}$ can be uniquely recovered from the curve $x \mapsto e^x \mathfrak{L}$, $x \in \delta_{\text{mod}}$ by osculation and taking the skew-orthogonal complement. This together with (8.8) and Remark 6.2 implies that $A \in \mathfrak{l}(X)$. In other words,

$$(8.9) \quad \mathcal{I}_2(\mathcal{E}(\delta_{\text{mod}})) \subset \mathfrak{l}(X).$$

Combining inclusions (8.7) and (8.9) with Theorem 7.2, Corollary 7.1, and the last statement of Lemma 8.3, we get the following

Theorem 8.2. *Assume that the flag symbol δ_{mod} satisfies all assumptions of Theorem 7.2 and in addition assumptions (5) and (6) above. Then for any positive integer k the k th Tanaka algebraic prolongation $\mathfrak{u}^k(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ of the pair $(\eta, \mathfrak{u}^F(\delta_{\text{mod}}))$ can be identified with the space $\mathcal{I}_{k+2}(S^k \mathcal{E}(\delta_{\text{mod}}))$ of all polynomials of degree $k+2$ on X that vanishes on the k th secant variety of the variety $\mathcal{E}(\delta_{\text{mod}})$.*

Remark 8.3. *Note that if $r \leq 1$, i.e. in the case of Jacobi symbols of distributions of rank 2, 3, or 4 of maximal class condition 6 is void (if one considers distinct indecomposable symbols in the decomposition of δ_{mod}) and therefore can be omitted in Theorem 8.2. Consider each of these ranks separately:*

- (1) *In the case of rank 2 distributions we already have Theorem 7.1, so Theorem 8.2 does not add anything.*
- (2) *In the case of rank 3 distributions by the first sentence of this remark and Corollary 6.2 all assumptions in Theorem 8.2 can be replaced by the assumption that the symplectically flat distribution is of maximal class, has 6-dimensional square, and the corresponding skew Young diagram is not rectangular.*
- (3) *In the case of rank 4 all assumptions in all assumptions in Theorem 8.2 can be replaced by the assumption that the symplectically flat distribution is of maximal class, the skew Young diagram of Z -graded component of its symbol is nonrectangular and has at least*

3 columns with 2 boxes and the skew diagram of $\frac{1}{2}\mathbb{Z}_{\text{odd}}$ -graded component is a row with at least 6 boxes. Note that the minimal dimension of the ambient space M , when such situation happens is 11 and it corresponds to the Jacobi symbol $\mathbb{R}\left(\delta_{3,2} \oplus \tau_{\frac{5}{2}}\right)$.

Now, following the same line, we show that in the case of Jacobi symbols of rank 3 distributions of maximal class with 6-dimensional square and non-rectangular skew Young diagram one can obtain more simple description of spaces $\mathbf{u}^k(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ in terms of a space of certain polynomials vanishing on some projective variety in a projective space of smaller dimension.

In the considered case $r = 1$, $X = E_1 \oplus F_1$, and $\delta = \delta_{\text{mod}} = \mathbb{R}\delta_{s,l}$ for some positive integer s and $s < l < 2s$. We also fix the skew Young tableaux as in 4.3. Recall that F_1 is an invariant subspace of δ . Let \mathfrak{C} be the (regular) rational normal curve in $\mathcal{P}F_1$, which is the closure of the orbit of the line represented by the vector f_{l-s} under the one-parametric group e^δ . The j th tangential developable $\mathcal{T}^j\mathfrak{C}$ is by definition (the algebraic closure of) the union of \mathfrak{C}^j of the j osculating subspaces $\mathfrak{C}^{(i)}$ of the curve \mathfrak{C} . Obviously $\mathcal{T}^0\mathfrak{C} = \mathfrak{C}$. Note that in our case $i_0 = 1$. Also by constructions we have the following relation

$$(8.10) \quad \mathcal{T}^{l-s-1}\mathfrak{C} = \mathcal{F}_\delta^1 \cap F_1$$

Lemma 8.6. *In the case $\delta_{\text{mod}} = \mathbb{R}\delta_{s,l}$ with $s < l < 2s$ the subspace \mathfrak{p} can be identified with the space $\mathcal{I}_2(\mathcal{T}^{l-s-1}\mathfrak{C})$ of all quadratic forms on F_1 that vanishes on the $(l-s-1)$ st tangential developable $\mathcal{T}^{l-s-1}\mathfrak{C}$ of the rational normal curve \mathfrak{C} .*

Proof. Following part b) of the proof of Lemma 6.1, we consider $S^2(E_1)$ as a subspace of $\text{Hom}(F_1, E_1)$ after identification of F_1^* with E_1 via the symplectic form σ and let $\mathfrak{s}(E_1)$ be the maximal \mathfrak{sl}_2 -submodule of $S^2(E_1)$ concentrated in the non-negative degree part of $S^2(E_1)$. From the same part b) of the proof of Lemma 6.1 the subspace \mathfrak{p} coincides with the subspace $\mathfrak{s}(E_1)$. Then, using the same arguments as in the proof of Lemma 8.2 and the relation (8.10), we get that $\tilde{\mathfrak{p}} \subset \mathcal{I}_2(\mathcal{T}^{l-s-1}\mathfrak{C})$.

On the other hand, by the same arguments as in the proof of inclusion (8.3) we get that the space $\mathcal{I}_2(\mathcal{T}^{l-s-1}\mathfrak{C})$ can be identified with the maximal \mathfrak{sl}_2 -submodule of $S^2(E_1)$ concentrated in the degrees greater or equal to -1 . However, from decomposition (6.19) of $S^2(E_1)$ into irreducible \mathfrak{sl}_2 modules it follows that the part of this decomposition concentrated in nonnegative degree is equal to the part of this decomposition concentrated in degree greater or equal than -1 . This completes the proof of the lemma. \square

As a direct consequence of Theorem 7.2, the last sentence of Lemma 8.3 and Lemma 8.6, we get the following

Theorem 8.3. *Assume that the flag symbol $\delta_{\text{mod}} = \mathbb{R}\delta_{s,l}$ with $s < l < 2s$. Then the k th Tanaka algebraic prolongation $\mathbf{u}^k(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ of the pair $(\eta, \mathbf{u}^F(\delta_{\text{mod}}))$ can be identified with the space $\mathcal{I}_{k+2}(S^k(\mathcal{T}^{l-s-1}\mathfrak{C}))$ of all polynomials of degree $k+2$ on $F_{s,l}$ that vanishes on the k th secant variety of the $l-s-1$ st tangential developable $\mathcal{T}^{l-s-1}\mathfrak{C}$ of the regular rational normal curve \mathfrak{C} on $\mathbb{P}F_{s,l}$.*

Remark 8.4. *In order to prove Theorem 8.3 we can also proceed as follows: prove that $\mathfrak{p} \subset \mathcal{I}_2(\mathcal{T}^{l-s-1}\mathfrak{C})$, as in the first paragraph of the proof of Lemma 8.6. Then observe that the space $\mathcal{I}_2(\mathcal{T}^{l-s-1}\mathfrak{C})$ can be naturally embedded into the space $\mathcal{I}_2(\mathcal{E}(\delta_{\text{mod}}))$ by extension of a quadratic form from F_1 to X such that E_1 is the kernel of this extension. Then from (8.9) we have that*

$\mathcal{I}_2(\mathcal{T}^{l-s-1}\mathfrak{C}) \subset \mathfrak{l}(X)$ and Theorem 8.3 follows from Theorem 7.2, Corollary 7.1, and the last statement of Lemma 8.3.

Let us consider the case $l = s + 1$ in more detail. First note that $\dim \mathbb{P}F_{s,s+1} = s + 1$. Let (x_1, \dots, x_{s+2}) are coordinates in $F_{s,s+1}$ with respect to the basis $(f_1, f_0, \dots, f_{-s})$. We have a well-known description of k -secant varieties of the rational normal curve \mathfrak{C} :

Lemma 8.7 ([24]). *The ideal $\mathcal{I}(\mathcal{S}^k\mathfrak{C})$ is generated (as an ideal) by $I_{k+2}(\mathcal{S}^k\mathfrak{C})$. The space $I_{k+2}(\mathcal{S}^k\mathfrak{C})$, in its turn, is generated (as a linear space) by all rank $k + 2$ minors of the matrix:*

$$\begin{pmatrix} x_1 & x_2 & x_3 & \dots & x_{s+2-\alpha} \\ x_2 & x_3 & x_4 & \dots & x_{s+3-\alpha} \\ \vdots & & & & \\ x_{\alpha+1} & x_{\alpha+2} & x_{\alpha+3} & \dots & x_{s+2} \end{pmatrix},$$

where α is an arbitrary integer between $k + 2$ and $s - k - 1$.

In particular, we see that in the considered case the space $\mathfrak{p}^{(k)}$, $k \geq 0$ coincides with the space $I_{k+2}(\mathcal{S}^k\mathfrak{C})$ and is described explicitly by Lemma 8.7.

Finally, in order to complete the case of rank 3 distributions of maximal class with 6-dimensional square consider the case of Jacobi symbol with rectangular Young diagram:

Theorem 8.4. *If a $(3, n)$ - distribution of maximal class with 6-dimensional square has Jacobi symbol with rectangular diagram, then the following statements hold:*

- (1) *The dimension n of the ambient manifold M is even and the (modified) Jacobi symbol is isomorphic to $\mathbb{R}\delta_{s,2s}$ with $s = \frac{n-4}{2}$.*
- (2) *The algebra $\mathfrak{u}^F(\delta_{mod})$ is isomorphic to the direct product $\mathfrak{sl}_2 \times \mathfrak{gl}_2$ acting on $X \cong E_1 \otimes \mathbb{R}^2$ componentwise such that \mathfrak{sl}_2 acts irreducibly on E_1 and the action of \mathfrak{gl}_2 on \mathbb{R}^2 is standard.*
- (3) [42] *If $n = 6$, then $\mathfrak{u}(\eta, \mathfrak{u}^F(\mathbb{R}\delta_{1,2}))$ is isomorphic to the algebra $\mathfrak{so}(4, 3)$ (as also expected from [4, 5]);*
- (4) *If n is even and greater than 6, then the first prolongation $\mathfrak{u}^1(\eta, \mathfrak{u}^F(\mathbb{R}\delta_{\frac{n-4}{2}, n-4}))$ of the pair $(\eta, \mathfrak{u}^F(\mathbb{R}\delta_{\frac{n-4}{2}, n-4}))$ vanishes, i.e. $\mathfrak{u}(\eta, \mathfrak{u}^F(\mathbb{R}\delta_{\frac{n-4}{2}, n-4}))$ is isomorphic to the semidirect sum of $\mathfrak{sl}_2 \times \mathfrak{gl}_2$ and $X \cong E_1 \otimes \mathbb{R}^2$ (with the action of $\mathfrak{sl}_2 \times \mathfrak{gl}_2$ on $E_1 \otimes \mathbb{R}^2$ described above).*

Proof. For item (1) as was already discussed in section 4, paragraph before Remark 4.2 a $(3, n)$ -distribution of maximal class has Jacobi symbol isomorphic to $\mathbb{R}\delta_{s, n-4}$ with nonnegative integer s satisfying $\frac{n-4}{2} \leq s \leq n-4$. This symbol has rectangular diagram if $s = \frac{n-4}{2}$, which implies that $n = 2s + 4$, i.e. n is even.

Item (2) follows from part b) of the proof of Lemma 6.1 applied for the case of rectangular Young diagram: take equations (6.22) and (6.21), first line there) and plug then into formulas (6.3) and (6.2).

For item (3) we refer to [42], where all cases of semisimple universal algebraic prolongations are considered, including our case here.

Let us prove item (4). The flag symbol in the considered situation satisfies all assumptions of Theorem 7.2: we are in the case of \mathbb{Z} grading and each row of the corresponding Young diagram consist of at least 5 boxes. Therefore to prove item (4) it is enough to show that

$$(8.11) \quad \mathfrak{p}^{(1)} = 0.$$

Note that from part b) of the proof of Lemma 6.1 it follows that \mathfrak{p} is 2-dimensional and, if the Young tableaux of our symbol is as in (4.3) then \mathfrak{p} is spanned by endomorphisms p_1 and p_2 of X defined as follows:

$$p_1(f_i) = e_i, p_2(e_i) = f_i, \quad \forall -s \leq i \leq s; p_1|_{E_1} = 0, p_2|_{F_1} = 0.$$

In the terminology used in item (2) of the Theorem p_1 and p_2 are the endomorphisms induced by the action of $\left(0, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}\right)$ and $\left(0, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}\right)$ (considered as element of $\mathfrak{sl}_2 \times \mathfrak{gl}_2$) on $X \cong E_1 \otimes \mathbb{R}^2$. Keeping this in mind, the relation (8.11) can be checked directly from the definition of the first standard prolongation and this check is left to the reader. \square

REFERENCES

- [1] A.A. Agrachev, *Feedback-invariant optimal control theory - II. Jacobi Curves for Singular Extremals*, J. Dynamical and Control Systems, **4**(1998), No. 4 , 583-604.
- [2] A.A. Agrachev, R.V. Gamkrelidze, *Feedback-invariant optimal control theory - I. Regular extremals*, J. Dynamical and Control Systems, **3**(1997), No. 3, 343-389.
- [3] D.V. Alekseevsky, A. Spiro, *Prolongations of Tanaka structures and regular CR structures*, Selected topics in Cauchy-Riemann geometry, 1-37, Quad. Mat., **9**, Dept. Math., Seconda Univ. Napoli, Caserta, 2001.
- [4] R. Bryant, *Conformal geometry and 3-plane fields on 6-manifolds*, Proceedings of the RIMS symposium "Developments of Cartan geometry and related mathematical problems" (24-27 October 2005).
- [5] R. Bryant, *Some aspects of the local and global theory of Pfaffian systems*, PhD thesis, University of North Carolina, 1979.
- [6] R.L. Bryant, S.S. Chern, R. B. Gardner, H.L. Goldschmidt, P. A. Griffiths, *Exterior Differential Systems*, Mathematical Sciences Research Institute Publications, vol. 18, Springer-Verlag.
- [7] A. Čap and H. Schichl, *Parabolic geometries and canonical Cartan connection*, Hokkaido Math. J. 29 (2000), 453505.
- [8] A. Čap and J. Slovák, *Parabolic Geometries I: Background and general theory*, Mathematical Surveys and Monographs, vol. 154, American Mathematical Society, Providence, RI, 2009
- [9] E. Cartan, *Les systemes de Pfaff a cinq variables et les equations aux derivees partielles du second ordre*, Oeuvres completes, Partie II, vol.2, Paris, Gautier-Villars, 1953, 927-1010.
- [10] Y. Chitour, F. Jean, E. Trélat, *Genericity results for singular curves*, J. Differential Geom. Volume 73, Number 1 (2006), 45-73.
- [11] B. Doubrov, *Contact trivialization of ordinary differential equations*, Differential Geometry and Its Applications, Proc. Conf., Opava (Czech Republic), 2001, Silestian University, Opava, pp. 73–84.
- [12] B. Doubrov, *Generalized Wilczynski invariants for non-linear ordinary differential equations*, In: Symmetries and Overdetermined Systems of Partial Differential Equations, IMA Volume 144, Springer, NY, 2008, pp. 25–40.
- [13] B. Doubrov, *Projective parametrization of homogeneous curves*, Arch. Math. (Brno) 41 (2005), no. 1, 129–133.
- [14] B. M. Doubrov, B. P. Komrakov, *Classification of homogeneous submanifolds in homogeneous spaces*, Lobachevskii J. Math., 1999, Volume 3, Pages 19-38.
- [15] B. Doubrov, I. Zelenko, *A canonical frame for nonholonomic rank two distributions of maximal class*, C.R. Acad. Sci. Paris, Ser. I, Vol. 342, Issue 8 (15 April 2006), 589-594, (see also arxiv math. DG/0504319)
- [16] B. Doubrov, I. Zelenko, *On local geometry of nonholonomic rank 2 distributions*, Journal of London Mathematical Society, (2) 80 (2009), no. 3, 545–566.
- [17] B. Doubrov, I. Zelenko, *Equivalence of variational problems of higher order*, Differential Geometry and its Applications , Volume 29, Issue 2, March 2011, 255-270.
- [18] B. Doubrov, I. Zelenko, *On local geometry of rank 3 distributions with 6-dimensional square*, submitted, arXiv:0807.3267v1[math. DG],40 pages.
- [19] B. Doubrov, I. Zelenko, *Geometry of curves in parabolic homogeneous spaces*, arXiv:1110.0226 v1 [math. DG], 20 pages, submitted.
- [20] B. Doubrov, I. Zelenko, *On geometry of curves of flags of constant type*, Cent. Eur. J. Math., 2012, 10(5), 1836-1871.
- [21] B. Doubrov, I. Zelenko *Prolongation of quasi-principal frame bundles and geometry of flag structures on manifolds*, preprint, arXiv:1210.7334 v2 [math.DG], 49 pages.
- [22] W. Fulton, J. Harris, *Representation theory: a first course*, Springer–Verlag, NY, 1991.

- [23] V. Guillemin, D. Quillen, S. Sternberg, *The classification of the irreducible complex algebras of infinite type*, J. Anal. Math., **18** (1967), pp. 107112.
- [24] J. Harris, *Algebraic geometry: a first course*, Springer-Verlag, NY, 1997.
- [25] N. Jacobson, *Lie algebras*, Intersci. Tracts in Pure and Appl. Math., 10, New York-London: John Wiley and Sons, 1962.
- [26] S. Kobayashi, *Transformation Groups in Differential Geometry*, Classics in Mathematics, Springer, 1995, 182 pages.
- [27] S. Kobayashi, T. Nagano, *On filtered Lie algebras and geometric structures III*, J. Math. Mech., v. 14, 1965, pp. 679–706.
- [28] B. Kruglikov, *Finite-dimensionality in Tanaka theory*, Ann. Inst. H. Poincaré Anal. Non Linéaire 28, no. 1, 7590 (2011).
- [29] B. Kruglikov, *Symmetries of filtered structures via the filtered Lie equations*, accepted to Journal of Geometry and Physics, arXiv:1401.6599v2 [math.DG].
- [30] O. Kuzmich, *Graded nilpotent Lie algebras in low dimensions*, Lobachevskii Journal of Mathematics, Vol.3, 1999, pp.147-184.
- [31] A. Ottazzi, B. Warhurst, *Algebraic prolongation and rigidity of Carnot groups*, Monatsh Math (2011), **162**: pp. 179-195.
- [32] N. Tanaka, *On differential systems, graded Lie algebras and pseudo-groups*, J. Math. Kyoto. Univ., **10** (1970), pp. 1–82.
- [33] N. Tanaka, *On the equivalence problems associated with simple graded Lie algebras*, Hokkaido Math. J.,**6**(1979), pp 23-84.
- [34] R. Montgomery, *A Tour of Subriemannian Geometries, Their Geodesics and Applications*, Mathematical Surveys and Monographs, Volume 91, (2002) American Mathematical Society.
- [35] T. Morimoto, *Geometric structures on filtered manifolds*, Hokkaido Math. J.,**22**(1993), pp. 263-347.
- [36] Pontryagin, Boltyanskii, Gamkrelidze and Mischenko, *The Mathematical Theory of Optimal Processes*. New York, Interscience Publishers, 1962.
- [37] D.C. Spencer, *Overdetermined systems of linear partial differential equations*, Bull. AMS, **75** (1969), pp. 179239.
- [38] S. Sternberg, *Lectures on differential geometry*, Prentice Hall, Englewood Cliffs, N.J., 1964.
- [39] N. Tanaka, *On nondegenerate real hypersurfaces, graded Lie algebras and Cartan connections*, Japan. J. Math. (N.S.) 2 (1976), no. 1, 131190.
- [40] N. Tanaka, *On affine symmetric spaces and the automorphism groups of product manifolds*, Hokkaido Math. J. 14 (1985) 277–351.
- [41] É. B. Vinberg, *Classification of homogeneous nilpotent elements of a semisimple graded Lie algebra*, Sel. Math. Sov., v. 6, 1987, 15–35.
- [42] K. Yamaguchi, *Differential Systems Associated with Simple Graded Lie Algebras*, Adv. Studies in Pure Math.,**22**(1993), pp.413-494.
- [43] I. Zelenko, *Nonregular abnormal extremals of 2-distribution: existence, second variation and rigidity*, J. Dynamical and Control systems , 5(1999), No. 3, 347-383.
- [44] I. Zelenko, *On Tanaka’s prolongation procedure for filtered structures of constant type* , Symmetry, Integrability and Geometry: Methods and Applications (SIGMA), Special Issue "Elie Cartan and Differential Geometry", v. 5, 2009, doi:10.3842/SIGMA.2009.094, 0906.0560 v3 [math.DG], 21 pages

BELARUSSIAN STATE UNIVERSITY, NEZAVISIMOSTI AVE. 4, MINSK 220030, BELARUS; E-MAIL: DOUBROV@ISLC.ORG

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA; E-MAIL: ZELENKO@MATH.TAMU.EDU