

Section 10.9 Applications of Taylor polynomials

Suppose that

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

Consider

$$T_n(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k$$

is the n th-degree Taylor polynomial of f at a .

We can use a Taylor polynomial T_n to approximate f . But how good an approximation is? To answer this question we need to look at

$$|R_n| = |f(x) - T_n(x)|$$

(a) If the series happen to be an alternating series, then

$$|R_n| \leq \frac{|f^{(n+1)}(a)|}{(n+1)!} |x-a|^{n+1}$$

(b) In other cases we can use **Taylor's Inequality**, which says if $|f^{(n+1)}(x)| \leq M$, then

$$|R_n| \leq \frac{M}{(n+1)!} |x-a|^{n+1}$$

Example 1.

(a) Approximate $f(x) = \sqrt{x}$ by a Taylor polynomial of degree 3 at $a = 1$.

(b) How accurate is this approximation if $0.9 \leq x \leq 1.1$?

Example 2. In Einstein's theory of special relativity the mass of an object moving with velocity v is

$$m = \frac{m_0}{\sqrt{1 - v^2/c^2}}$$

where m_0 is the mass of the object when at rest and c is the speed of the light ($c \approx 300 \times 10^6$ m/s). The kinetic energy of the object is the difference between its total energy and its energy at rest:

$$K = mc^2 - m_0c^2$$

(a) Show that when v is very small compared with c , this expression for K agrees with classical Newtonian physics: $K = \frac{1}{2}m_0v^2$.

(b) Use Taylor's Inequality to estimate the difference in these expression for K when $|v| \leq 100$ m/s.