

Section 10.5 Power series

A **power series** is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n + \dots$$

Constants c_n are called the **coefficients** of the series. For each fixed x , the series $\sum_{n=0}^{\infty} c_n x^n$ is a series of constants that we can test for convergence or divergence. A power series may converge for some values of x and diverge for other values of x . The sum of the series is a function

$$f(x) = c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n + \dots$$

whose domain is the set of all x for which the series converges.

More generally, a series of the form

$$\sum_{n=0}^{\infty} c_n (x - a)^n$$

is called a **power series centered at a** or a **power series about a** .

A power series is convergent if $|x - a| < R$, where

$$R = \lim_{n \rightarrow \infty} \left| \frac{c_n}{c_{n+1}} \right|$$

or

$$R = \lim_{n \rightarrow \infty} \frac{1}{\sqrt[n]{|c_n|}}$$

R is called the **radius of convergence**.

- If $R = 0$, then the series converges only at one point $x = a$.
- If $R = \infty$, then the series converges for all x .
- If $R \neq 0$ and $R < \infty$, then the series converges if $a - R < x < a + R$. Also, test the series for convergence at $x = a - R$ and $x = a + R$.

The **interval of convergence** of a power series is the interval that consists of all values of x for which the series is convergent.

Example. Find the radius of convergence and interval of convergence for each of the following series

1. $\sum_{n=0}^{\infty} x^n$

2. $\sum_{n=0}^{\infty} \frac{x^n}{(n+2)!}$

$$3. \sum_{n=0}^{\infty} \frac{n!(x+1)^n}{\sqrt{n+3}}$$

$$4. \sum_{n=1}^{\infty} \frac{(-1)^n x^n}{\sqrt[3]{n}}$$

$$5. \sum_{n=1}^{\infty} \frac{(x-4)^n}{n5^n}$$

$$6. \sum_{n=0}^{\infty} \frac{n^2(x+1)^{2n}}{10^n}$$