

Math 251. WEEK in REVIEW 8. Fall 2013

- Find the gradient vector field of $f(x, y, z) = x \ln(y^4 + z^2)$
- Let C be the line segment starting at $(0, 1, 1)$ and ending at $(3, 1, 4)$. Find the mass of a thin wire in the shape of C with the density $\rho(x, y) = x + y$.
- Find the mass of a thin wire in the shape of C with the density $\rho(x, y, z) = 7y^2z$ if C is given by $\vec{r}(t) = \langle \frac{2}{3}t^3, t, t^2 \rangle$, $0 \leq t \leq 1$.
- Find the line integral of the vector field $\vec{F}(x, y, z) = \langle -yz^2, xz^2, z^3 \rangle$ around the circle $\vec{r}(t) = \langle 2 \cos t, 2 \sin t, 8 \rangle$.
- Find the work done by the force field $\vec{F}(x, y, z) = \langle z, x, y \rangle$ in moving a particle from the point $(3, 0, 0)$ to the point $(0, \pi/2, 3)$
 - along the straight line
 - along the helix $x = 3 \cos t$, $y = t$, $z = 3 \sin t$
- Find a scalar function $f(x, y, z)$ such that $\nabla f = \langle 2xy + z, x^2 - 2y, x \rangle$ and $f(1, 2, 0) = 3$.
- Let $\vec{F}(x, y) = \langle x^3y^4, x^4y^3 \rangle$. Compute $\int_C \vec{F} \cdot d\vec{r}$ where
 - C is any simple closed path.
 - C is any path from the point $M(0, 0)$ to the point $N(1, 2)$.
- Let $\vec{F}(x, y) = \langle x + y^2, 2xy + y^2 \rangle$.
 - Show that \vec{F} is conservative vector field.
 - Compute $\int_C \vec{F} \cdot d\vec{r}$ where C is any path from $(-1, 0)$ to $(2, 2)$.
- Let $\vec{F}(x, y) = \langle 2x + y^2 + 3x^2y, 2xy + x^3 + 3y^2 \rangle$.
 - Show that \vec{F} is conservative vector field.
 - Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where C is the arc of the curve $y = x \sin x$ from $(0, 0)$ to $(\pi, 0)$.
- Compute the integral $I = \oint_C (\cos x^4 + xy) dx + (y^2e^y + x^2) dy$, where C is the triangular curve consisting of the line segments from $(0, 0)$ to $(1, 0)$, from $(1, 0)$ to $(1, 3)$, and from $(1, 3)$ to $(0, 0)$.
- Compute the integral along the given positively oriented curve C :

$$\int_C (y^2 - \arctan x) dx + (3x + \sin y) dy,$$

where C is the boundary of the region enclosed by the parabola $y = x^2$ and the line $y = 4$.

12. Compute the integral

$$\int_C (12 - x^2y - y^3 + \tan x) dx + (xy^2 + x^3 - e^y) dy$$

where C is positively oriented boundary of the region enclosed by the circle $x^2 + y^2 = 4$. Sketch the curve C indicating the positive direction.

13. Given the line integral $I = \int_C 4x^2y dx - (2 + x) dy$ where C consists of the line segment from $(0, 0)$ to $(2, -2)$, the line segment from $(2, -2)$ to $(2, 4)$, and the part of the parabola $y = x^2$ from $(2, 4)$ to $(0, 0)$. Use Green's theorem to **evaluate** the given integral and **sketch** the curve C indicating the *positive direction*.