

GROUPOID NORMALIZERS OF TENSOR PRODUCTS

Junsheng Fang
jfang@cisunix.unh.edu

Roger R. Smith*
rsmith@math.tamu.edu

Stuart A. White
s.white@maths.gla.ac.uk

Alan D. Wiggins
alan.d.wiggins@vanderbilt.edu

September 15, 2008

Abstract

We consider an inclusion $B \subseteq M$ of finite von Neumann algebras satisfying $B' \cap M \subseteq B$. A partial isometry $v \in M$ is called a groupoid normalizer if vBv^* , $v^*Bv \subseteq B$. Given two such inclusions $B_i \subseteq M_i$, $i = 1, 2$, we find approximations to the groupoid normalizers of $B_1 \overline{\otimes} B_2$ in $M_1 \overline{\otimes} M_2$, from which we deduce that the von Neumann algebra generated by the groupoid normalizers of the tensor product is equal to the tensor product of the von Neumann algebras generated by the groupoid normalizers. Examples are given to show that this can fail without the hypothesis $B'_i \cap M_i \subseteq B_i$, $i = 1, 2$. We also prove a parallel result where the groupoid normalizers are replaced by the intertwiners, those partial isometries $v \in M$ satisfying $vBv^* \subseteq B$ and $v^*v, vv^* \in B$.

1 Introduction

The focus of this paper is an inclusion $B \subseteq M$ of finite von Neumann algebras. Such inclusions have a rich diverse history, first being studied by Dixmier [3] in the context of maximal abelian subalgebras (masas) of II_1 factors. These inclusions provided the basic building blocks for the theory of subfactors developed by Jones in [10] and today they are a key component in the study of structural properties of II_1 factors using the deformation-rigidity techniques introduced by Popa in [14].

In [3], Dixmier introduced a classification of masas in II_1 factors using *normalizers*, defining $\mathcal{N}_M(B) = \{u \text{ a unitary in } M : uBu^* = B\}$. A masa $B \subset M$ is *Cartan* or *regular* if these normalizers generate M and *singular* if $\mathcal{N}_M(B) \subset B$. Feldman and Moore demonstrated the importance of Cartan masas, and hence normalizers, in the study of II_1 factors, showing that inclusions of Cartan masas arise from measurable equivalence relations and that, up to orbit equivalence, these relations determine the resulting inclusion, [7, 8].

Given two inclusions $B_i \subset M_i$ of masas in II_1 factors, it is immediate that an elementary tensor $u_1 \otimes u_2$ of unitaries $u_i \in M_i$ normalizes the tensor product inclusion $B = B_1 \overline{\otimes} B_2 \subset$

*Partially supported by a grant from the National Science Foundation

$M = M_1 \overline{\otimes} M_2$ if and only if each u_i normalizes B_i . As a simple consequence, the tensor product of Cartan masas is again Cartan. More generally, the operation of passing to the von Neumann algebra generated by the normalizers was shown to commute with the tensor product operation for masas inside II_1 factors, in the sense that the equality

$$\mathcal{N}_{M_1}(B_1)'' \overline{\otimes} \mathcal{N}_{M_2}(B_2)'' = \mathcal{N}_{M_1 \overline{\otimes} M_2}(B_1 \overline{\otimes} B_2)'', \quad (1.1)$$

holds. This was proved when both masas are singular in [19] and the general case was established by Chifan in [1]. Since the containment from left to right in (1.1) is immediate, the problem in both cases is to eliminate the possibility that some unexpected unitary in the tensor product normalizes $B_1 \overline{\otimes} B_2$. This difficulty was overcome in [19] and [1] by employing techniques of Popa [15] to analyse the basic construction algebra $\langle M, e_B \rangle$ of Jones [10]. Beyond the masa setting, (1.1) holds when each B_i satisfies $B'_i \cap M_i = \mathbb{C}1$, the defining property of irreducible subfactors. When each B_i has finite Jones index in M_i , the identity (1.1) can be deduced from results of [13]. The infinite index case was established in [20], where every normalizing unitary of such a tensor product of irreducible subfactors was shown to be of the form $w(v_1 \otimes v_2)$, where w is a unitary in $B_1 \overline{\otimes} B_2$ and each $v_i \in \mathcal{N}_{M_i}(B_i)$. Some other situations where (1.1) holds are discussed in [6].

For general inclusions $B_i \subseteq M_i$ of finite von Neumann algebras, the commutation identity (1.1) can fail. Indeed, taking each M_i to be a copy of the 3×3 matrices and each $B_i \cong \mathbb{C} \oplus \mathbb{M}_2(\mathbb{C})$, one obtains inclusions with $\mathcal{N}_{M_i}(B_i) \subset B_i$, yet there are non-trivial normalizers of $B_1 \otimes B_2$ inside $M_1 \otimes M_2$. This is due to the presence of partial isometries v in $M_i \setminus B_i$ with $vB_iv^* \subseteq B_i$ and $v^*B_iv \subseteq B_i$, as the non-trivial unitary normalizers of $B_1 \otimes B_2$ can all be written in the form $\sum_j x_j(v_{1,j} \otimes v_{2,j})$, where x_j lie in $B_1 \otimes B_2$ and the $v_{i,j}$ are partial isometries with $v_{i,j}B_iv_{i,j}^* \subseteq B$ and $v_{i,j}^*B_iv_{i,j} \subseteq B_i$. Defining the *groupoid normalizers* of a unital inclusion $B \subset M$ to be the set $\mathcal{GN}_M(B) = \{v \text{ a partial isometry in } M : vBv^* \subseteq B, v^*Bv \subseteq B\}$, the example discussed above satisfies the commutation identity

$$\mathcal{GN}_{M_1}(B_1)'' \overline{\otimes} \mathcal{GN}_{M_2}(B_2)'' = \mathcal{GN}_{M_1 \overline{\otimes} M_2}(B_1 \overline{\otimes} B_2)''. \quad (1.2)$$

In this paper we examine groupoid normalizers of tensor product algebras, establishing (Corollary 5.6) the identity (1.2) whenever $B_i \subseteq M_i$ are inclusions of finite von Neumann algebras with separable preduals satisfying $B'_i \cap M_i \subseteq B_i$ for each i . In [4] Dye shows that every groupoid normalizer v of a masa B in M is of the form $v = ue$ for some projection $e = v^*v \in B$ and some unitary normalizer u of B in M , see also [18, Lemma 6.2.3]. The same result holds by a direct computation when B is an irreducible subfactor of M , so that in these two cases $\mathcal{N}_M(B)'' = \mathcal{GN}_M(B)''$ and (1.2) directly generalizes (1.1) established in [1] and [20] respectively.

The following example shows why the hypothesis $B'_i \cap M_i \subseteq B_i$ (which is satisfied by both masas and irreducible subfactors) is necessary in this result.

Example 1.1. Consider the subalgebra

$$B = \left\{ \begin{pmatrix} \alpha & 0 & 0 \\ 0 & \alpha & 0 \\ 0 & 0 & \beta \end{pmatrix} : \alpha, \beta \in \mathbb{C} \right\} \subseteq \mathbb{M}_3,$$

and note that $B' \cap \mathbb{M}_3$ strictly contains B . A direct computation shows that $\mathcal{GN}(B)'' = \mathbb{M}_2 \oplus \mathbb{C}$, and so $\mathcal{GN}_{\mathbb{M}_3}(B)'' \otimes \mathcal{GN}_{\mathbb{M}_3}(B)'' \cong \mathbb{M}_4 \oplus \mathbb{M}_2 \oplus \mathbb{M}_2 \oplus \mathbb{C}$. However, $B \otimes B$ is isomorphic to $\mathbb{C}I_4 \oplus \mathbb{C}I_2 \oplus \mathbb{C}I_2 \oplus \mathbb{C}$ inside \mathbb{M}_9 , and $\mathcal{GN}_{\mathbb{M}_3 \otimes \mathbb{M}_3}(B \otimes B)''$ is $\mathbb{M}_4 \oplus \mathbb{M}_4 \oplus \mathbb{C}$.

A new feature of [20] was the notion of *one-sided normalizers* of an irreducible inclusion $B \subset M$ of II_1 factors, namely those unitaries $u \in M$ with $uBu^* \not\subseteq B$. These cannot arise for finite index inclusions by index considerations, or in the case when $B \subset M$ is a masa. To establish (1.1) for irreducible subfactors, it was necessary to first establish the general form of a one-sided normalizer of a tensor product of irreducible subfactors and then deduce the normalizer result from this. The same procedure is necessary here, so we introduce the notion of an *intertwiner* to study groupoid normalizers in a one-sided situation.

Definition 1.2. Given an inclusion $B \subseteq M$ of von Neumann algebras satisfying $B' \cap M \subseteq B$, define the collection $\mathcal{GN}_M^{(1)}(B)$ of *intertwiners* of B in M by

$$\mathcal{GN}_M^{(1)}(B) = \{v \text{ a partial isometry in } M : vBv^* \subseteq B, v^*v \in B\}.$$

We will write $\mathcal{GN}^{(1)}(B)$ for $\mathcal{GN}_M^{(1)}(B)$ when there is no confusion about the underlying algebra M . We use the superscript ⁽¹⁾ to indicate that our intertwiners are *one-sided*, namely that although $vBv^* \subseteq B$, we are not guaranteed to have a containment $v^*v \subseteq B$. Note that $v \in \mathcal{GN}_M(B)$ if, and only if, both v and v^* lies in $\mathcal{GN}_M^{(1)}(B)$. Note too that while the groupoid normalizers form a groupoid, the intertwiners do not. Finally, the terminology intertwiner comes from the fact that, under the hypothesis $B' \cap M \subseteq B$, these are exactly the partial isometries that witness the embeddability of a corner of B into itself inside M in the sense of Popa's intertwining procedure for subalgebras from [14, 15].

We obtain a similar commutation result to (1.2) for intertwiners. In fact our main theorem, stated below, obtains more as it gives approximate forms for intertwiners and groupoid normalizers of tensor products.

Theorem 1.3. *Let $B_i \subset M_i$ be inclusions of finite von Neumann algebras with separable preduals and with fixed faithful normal traces τ_i on M_i . For $v \in \mathcal{GN}_{M_1 \overline{\otimes} M_2}^{(1)}(B_1 \overline{\otimes} B_2)$ and $\varepsilon > 0$, there exist $k \in \mathbb{N}$ and operators $x_1, \dots, x_k \in B_1 \overline{\otimes} B_2$, intertwiners $w_{1,1}, \dots, w_{1,k}$ of B_1 in M_1 and intertwiners $w_{2,1}, \dots, w_{2,k}$ of B_2 in M_2 such that*

$$\left\| v - \sum_{j=1}^k x_j (w_{1,j} \otimes w_{2,j}) \right\|_2 < \varepsilon, \quad (1.3)$$

where the $\|\cdot\|_2$ -norm arises from the trace $\tau_1 \otimes \tau_2$ on $M_1 \overline{\otimes} M_2$. If in addition v is a groupoid normalizer, then each $w_{i,j}$ can be taken to be a groupoid normalizer rather than just an intertwiner.

The intertwiner form of Theorem 1.3 is established as Theorem 4.7 and additional analysis in Section 5 enables us to deduce the groupoid normalizer form of Theorem 1.3 as Theorem 5.5. For the remainder of the introduction we give a summary of the main steps used to establish these results and where they can be found in the paper.

Given inclusions $B_i \subseteq M_i$ of finite von Neumann algebras with $B'_i \cap M_i \subseteq B_i$, write $B \subset M$ for the tensor product inclusion $B_1 \overline{\otimes} B_2 \subseteq M_1 \overline{\otimes} M_2$. Let $v \in \mathcal{GN}_M^{(1)}(B)$. Then

the element v^*e_Bv is a projection in the basic construction algebra $\langle M, e_B \rangle$, the properties of which are recalled in Section 2. Section 3 discusses the properties of these projections in the basic construction arising from intertwiners. In particular, we show that the projection v^*e_Bv is central in the cutdown $(B' \cap \langle M, e_B \rangle)v^*v$ (Lemma 3.2) and construct an explicit projection $P_v \in Z(B' \cap \langle M, e_B \rangle)$ with $P_v v^*v = v^*e_Bv$. We need to construct this projection explicitly rather than appeal to general theory, as its properties (established in Lemma 3.8) are crucial subsequently.

Since the basic construction factorizes as a tensor product $\langle M, e_B \rangle \cong \langle M_1, e_{B_1} \rangle \overline{\otimes} \langle M_2, e_{B_2} \rangle$, Tomita's commutation theorem gives

$$Z(B' \cap \langle M, e_B \rangle) \cong Z(B'_1 \cap \langle M_1, e_{B_1} \rangle) \overline{\otimes} Z(B'_2 \cap \langle M_2, e_{B_2} \rangle). \quad (1.4)$$

For each $i = 1, 2$, let Q_i denote the supremum of all projections in $Z(B'_i \cap \langle M_i, e_{B_i} \rangle)$ of the form $\sum_j w_{i,j}^* e_{B_i} w_{i,j}$, where the $w_{i,j}$ lie in $\mathcal{GN}_{M_i}^{(1)}(B_i)$ and satisfy $w_{i,j} w_{i,k}^* = 0$ when $j \neq k$. If we can show that

$$P_v \leq Q_1 \otimes Q_2, \quad (1.5)$$

then it will follow that we can approximate P_v in $L^2(\langle M, e_B \rangle)$ by projections of the form $\sum_j (w_{1,j} \otimes w_{2,j})^* e_B (w_{1,j} \otimes w_{2,j})$ for intertwiners $w_{i,j} \in \mathcal{GN}_{M_i}^{(1)}(B_i)$. To do this, we use the fact that projections in the tensor product (1.4) of *abelian* von Neumann algebras can be approximated by sums of elementary tensors of projections, and so it is crucial that the original projection v^*e_Bv be *central* in $(B' \cap \langle M, e_B \rangle)v^*v$, for which the hypothesis $B' \cap M \subseteq B$ is necessary. Finally, we push the approximation for P_v down to M and obtain the required approximation for v in M (see Theorem 4.7).

Most of Section 4 is taken up with establishing (1.5). We give a technical result (Theorem 4.1), which in particular characterizes when a projection in the basic construction arises from an intertwiner. By applying Theorem 4.1 to P_v and the inclusion

$$Z(B'_1 \cap \langle M_1, e_{B_1} \rangle) \overline{\otimes} B_2 \subseteq Z(B'_1 \cap \langle M_1, e_{B_1} \rangle) \overline{\otimes} M_2,$$

regarded as a direct integral of inclusions of finite von Neumann algebras, we are able to establish $P_v \leq 1 \otimes Q_2$ in Lemma 4.6 and so (1.5) follows by symmetry. It should be noted that the introduction of the projections Q_i is essential in order to make use of measure theory, particularly the uniqueness of product measures on σ -finite spaces, [16, p. 312]. The canonical trace on the basic construction need not be a semifinite weight on $Z(B'_i \cap \langle M_i, e_{B_i} \rangle)$ but does have this property on the compression $Z(B'_i \cap \langle M_i, e_{B_i} \rangle)Q_i$ where it can be treated as a measure (see Lemma 2.6 and the discussion preceding Definition 4.4). The remaining difficulty is to check that the projection P_v satisfies the hypotheses of Theorem 4.1, for which we require certain order properties of the pull-down map on the basic construction. These are described in the next section, in which we also set out our notation, review the properties of the basic construction, and establish some technical lemmas. Finally, the paper ends with Section 5, which handles the additional details required to deduce the groupoid normalizer result (Theorem 5.5) from our earlier work.

Acknowledgment: The work in this paper originated during the *Workshop in Analysis and Probability*, held at Texas A& M University during Summer 2007. It is a pleasure to express our thanks to both the organizers of the workshop and to the NSF for providing financial support to the workshop.

2 Notation and preliminaries

Throughout the paper, all von Neumann algebras are assumed to have separable preduals. The basic object of study in this paper is an inclusion $B \subseteq M$ of finite von Neumann algebras, where M is equipped with a faithful normal trace τ satisfying $\tau(1) = 1$. We always assume that M is standardly represented on the Hilbert space $L^2(M, \tau)$, or simply $L^2(M)$. The letter ξ is reserved for the image of $1 \in M$ in this Hilbert space, and J will denote the isometric conjugate linear operator on $L^2(M)$ defined on $M\xi$ by $J(x\xi) = x^*\xi$, $x \in M$, and extended by continuity to $L^2(M)$ from this dense subspace. Then $L^2(B)$ is a closed subspace of $L^2(M)$, and e_B denotes the projection of $L^2(M)$ onto $L^2(B)$, called the Jones projection. The von Neumann algebra generated by M and e_B is called the basic construction and is denoted by $\langle M, e_B \rangle$, [2, 10]. Let \mathbb{E}_B denote the unique trace preserving conditional expectation of M onto B . In the next proposition we collect together standard properties of e_B, \mathbb{E}_B and $\langle M, e_B \rangle$ from [10, 13, 9, 18].

Proposition 2.1. (i) $e_B(x\xi) = \mathbb{E}_B(x)\xi$, $x \in M$.

(ii) $e_Bxe_B = \mathbb{E}_B(x)e_B = e_B\mathbb{E}_B(x)$, $x \in M$.

(iii) $M \cap \{e_B\}' = B$.

(iv) $\langle M, e_B \rangle' = JBJ$, $Z(\langle M, e_B \rangle) = JZ(B)J$.

(v) e_B has central support 1 in $\langle M, e_B \rangle$.

(vi) $\text{Span}\{xe_{By} : x, y \in M\}$ generates a $*$ -strongly dense subalgebra, denoted Me_BM , of $\langle M, e_B \rangle$.

(vii) $x \mapsto e_Bx$ and $x \mapsto xe_B$ are injective maps for $x \in M$.

(viii) Me_B and e_BM are $*$ -strongly dense in $\langle M, e_B \rangle e_B$ and $e_B \langle M, e_B \rangle$ respectively.

(ix) $e_B \langle M, e_B \rangle e_B = Be_B = e_BB$.

(x) $(Me_BM) \langle M, e_B \rangle (Me_BM) \subseteq Me_BM$.

(xi) There is a unique faithful normal semifinite trace Tr on $\langle M, e_B \rangle$ satisfying

$$\text{Tr}(xe_{By}) = \tau(xy), \quad x, y \in M. \quad (2.1)$$

This trace is given by the formula

$$\text{Tr}(t) = \sum_{i=1}^{\infty} \langle tJv_i^*\xi, Jv_i^*\xi \rangle, \quad t \in \langle M, e_B \rangle^+, \quad (2.2)$$

where the v_i 's are partial isometries in $\langle M, e_B \rangle$ satisfying $\sum_{i=1}^{\infty} v_i^*e_Bv_i = 1$.

(xii) The algebra Me_BM is $\|\cdot\|_{2, \text{Tr}}$ -dense in $L^2(\langle M, e_B \rangle, \text{Tr})$ and $\|\cdot\|_{1, \text{Tr}}$ -dense in $L^1(\langle M, e_B \rangle, \text{Tr})$.

(xiii) Given inclusions $B_i \subset M_i$ of finite von Neumann algebras for $i = 1, 2$, the basic construction $\langle M_1 \overline{\otimes} M_2, e_{B_1} \overline{\otimes} e_{B_2} \rangle$ is isomorphic to $\langle M_1, e_{B_1} \rangle \overline{\otimes} \langle M_2, e_{B_2} \rangle$. Under this isomorphism, the canonical trace Tr on $\langle M_1 \overline{\otimes} M_2, e_{B_1} \overline{\otimes} e_{B_2} \rangle$ is given by $\text{Tr}_1 \otimes \text{Tr}_2$, where Tr_i is the canonical trace on $\langle M_i, e_{B_i} \rangle$.

(xiv) There is a well defined map $\Psi : Me_B M \rightarrow M$, given by

$$\Psi(xe_B y) = xy, \quad x, y \in M. \quad (2.3)$$

This is the pull down map of [13], where it was shown to extend to a contraction from $L^1(\langle M, e_B \rangle, \text{Tr})$ to $L^1(M, \tau)$.

Using Part (xii) of the previous proposition, the equation

$$\text{Tr}((xe_B y)z) = \tau((xy)z) = \tau(\Psi(xe_B y)z), \quad x, y, z \in M, \quad (2.4)$$

shows that Ψ is the pre-adjoint of the identity embedding $M \hookrightarrow \langle M, e_B \rangle$ and is, in particular, positive. The basic properties of Ψ are set out in [13], but we will need more detailed information on this map than is currently available in the literature. We devote much of this section to obtaining further properties of Ψ , the main objective being to apply them in Lemma 4.5.

In the next three lemmas, the inclusion $B \subset M$ is always of arbitrary finite von Neumann algebras with a fixed faithful normalized normal trace τ on M , inducing the trace Tr on $\langle M, e_B \rangle$.

Lemma 2.2. *Let $x \in L^1(\langle M, e_B \rangle)^+ \cap \langle M, e_B \rangle$. If $\Psi(x) \in L^1(M) \cap M$, then $\Psi(x) \geq x$.*

Proof. It suffices to show that

$$\langle \Psi(x)y\xi, y\xi \rangle \geq \langle xy\xi, y\xi \rangle, \quad y \in M. \quad (2.5)$$

The maximality argument, preceding [18, Lemma 4.3.4], to establish part (xi) of Proposition 2.1 can be easily modified to incorporate the requirement that $v_1 = 1$. Thus there are vectors $\xi_i = Jv_i^* \xi \in L^2(M)$ so that (2.2) becomes

$$\text{Tr}(t) = \sum_{i=1}^{\infty} \langle t\xi_i, \xi_i \rangle, \quad t \in \langle M, e_B \rangle^+, \quad (2.6)$$

where $\xi_1 = \xi$. Now, for $y \in M$, we may use the M -modularity of Ψ to write

$$\langle \Psi(x)y\xi, y\xi \rangle = \langle \Psi(y^*xy)\xi, \xi \rangle = \tau(\Psi(y^*xy)) = \text{Tr}(y^*xy). \quad (2.7)$$

It follows from (2.6) and (2.7) that

$$\langle \Psi(x)y\xi, y\xi \rangle = \langle xy\xi, y\xi \rangle + \sum_{i=2}^{\infty} \langle xy\xi_i, y\xi_i \rangle \geq \langle xy\xi, y\xi \rangle, \quad y \in M, \quad (2.8)$$

establishing that $\Psi(x) \geq x$. □

We now extend this result to tensor products. Let N be a semifinite von Neumann algebra with a specified faithful normal semifinite trace TR . In [5], Effros and Ruan identified the predual of a tensor product of von Neumann algebras X and Y by $(X \overline{\otimes} Y)_* = X_* \otimes_{op} Y_*$, the operator space projective tensor product of the preduals. In the presence of traces, this identifies $L^1(X \overline{\otimes} Y)$ with $L^1(X) \otimes_{op} L^1(Y)$, so $I \otimes \Psi$ is well defined, positive, and bounded from $L^1(N \overline{\otimes} \langle M, e_B \rangle, \text{TR} \otimes \text{Tr})$ to $L^1(N \overline{\otimes} M, \text{TR} \otimes \tau)$, being the pre-adjoint of the identity embedding $N \overline{\otimes} M \hookrightarrow N \overline{\otimes} \langle M, e_B \rangle$. Following [21, Chapter IX], we will always assume that N is faithfully represented on $L^2(N, \text{TR})$, for which $\text{span}\{y \in N : \text{TR}(y^*y) < \infty\}$ is a dense subspace.

Lemma 2.3. *Let $x \in L^1(N \overline{\otimes} \langle M, e_B \rangle)^+ \cap (N \overline{\otimes} \langle M, e_B \rangle)$. If $(I \otimes \Psi)(x) \in L^1(N \overline{\otimes} M) \cap (N \overline{\otimes} M)$, then $(I \otimes \Psi)(x) \geq x$.*

Proof. Suppose that the result is not true. Then we may find a finite projection $p \in N$, elements $y_i \in pNp$ and $z_i \in M$, $1 \leq i \leq k$, so that

$$\left\langle (x - (I \otimes \Psi)(x)) \sum_{i=1}^k y_i \otimes z_i \xi, \sum_{i=1}^k y_i \otimes z_i \xi \right\rangle > 0, \quad (2.9)$$

since such sums $\sum_{i=1}^k y_i \otimes z_i \xi$ are dense in $L^2(N, \text{TR}) \otimes_2 L^2(M, \tau)$. Then the inequality

$$(I \otimes \Psi)((p \otimes 1)x(p \otimes 1)) \geq (p \otimes 1)x(p \otimes 1) \quad (2.10)$$

fails. The element on the left of (2.10) is $(p \otimes 1)((I \otimes \Psi)(x))(p \otimes 1)$, and so is bounded by hypothesis. The restriction of $I \otimes \Psi$ to $L^1(pNp \overline{\otimes} \langle M, e_B \rangle)$ is the pull down map for the inclusion $pNp \overline{\otimes} B \subseteq pNp \overline{\otimes} M$ of finite von Neumann algebras with basic construction $pNp \overline{\otimes} \langle M, e_B \rangle$. The failure of (2.10) then contradicts Lemma 2.2 applied to this inclusion, establishing that $(1 \otimes \Psi)(x) \geq x$. \square

The next lemma completes our investigation of the order properties of pull down maps.

Lemma 2.4. *If $x \in L^1(N \overline{\otimes} \langle M, e_B \rangle)^+$ is unbounded, then so also is $(1 \otimes \Psi)(x)$.*

Proof. Suppose that $(1 \otimes \Psi)(x)$ is bounded. Following [21, Section IX.2], we may regard x as a self-adjoint positive densely defined operator on $L^2(N \overline{\otimes} \langle M, e_B \rangle)$. For $n \geq 1$, let $p_n \in N$ be the spectral projection of x for the interval $[0, n]$. Then $p_n x \leq x$, so $(I \otimes \Psi)(p_n x) \leq (I \otimes \Psi)(x)$, since $I \otimes \Psi$ is the pre-adjoint of a positive map. In particular, $I \otimes \Psi(p_n x)$ is bounded. By Lemma 2.3 applied to $p_n x$,

$$(I \otimes \Psi)(x) \geq (I \otimes \Psi)(p_n x) \geq p_n x. \quad (2.11)$$

Since $n \geq 1$ was arbitrary, we conclude from (2.11) that x is bounded, a contradiction which completes the proof. \square

We note for future reference that these results are equally valid for pull down maps of the form $\Psi \otimes I$, due to symmetry. These lemmas will be used in Section 4 to derive an important inequality. The next lemma formulates exactly what will be needed.

Lemma 2.5. *Let $B_i \subseteq M_i$, $i = 1, 2$, be inclusions of finite von Neumann algebras with pull down maps Ψ_i . Let $B \subseteq M$ be the inclusion $B_1 \overline{\otimes} B_2 \subseteq M_1 \overline{\otimes} M_2$. If $x \in L^1(\langle M, e_B \rangle)^+ \cap \langle M, e_B \rangle$ is such that $(\Psi_1 \otimes \Psi_2)(x) \in L^1(M) \cap M$ and the inequality*

$$\|(\Psi_1 \otimes \Psi_2)(x)\| \leq 1 \tag{2.12}$$

is satisfied, then $(I \otimes \Psi_2)(x) \in L^1(\langle M_1, e_{B_1} \rangle \overline{\otimes} M_2)^+ \cap (\langle M_1, e_{B_1} \rangle \overline{\otimes} M_2)$ and $\|(I \otimes \Psi_2)(x)\| \leq 1$.

Proof. Using the isomorphism of Proposition 2.1 (xiii), $\Psi_1 \otimes \Psi_2$ is the pull down map for $\langle M, e_B \rangle$. Since $(\Psi_1 \otimes I)((I \otimes \Psi_2)(x)) = (\Psi_1 \otimes \Psi_2)(x)$ is a bounded operator by hypothesis, it follows from Lemma 2.4 that $(I \otimes \Psi_2)(x)$ is also bounded in $\langle M_1, e_{B_1} \rangle \overline{\otimes} M_2$. Thus the three operators x , $(I \otimes \Psi_2)(x)$ and $(\Psi_1 \otimes \Psi_2)(x)$ are all bounded, and so we may apply Lemma 2.3 twice to the pull down maps $I \otimes \Psi_2$ and $\Psi_1 \otimes I$ to obtain

$$(\Psi_1 \otimes \Psi_2)(x) \geq (I \otimes \Psi_2)(x) \geq x. \tag{2.13}$$

The result then follows from (2.13) and the hypothesis (2.12). \square

In the proof of Lemma 4.6, we will need the following fact regarding inclusions of finite von Neumann algebras $B \subseteq M$ with $B' \cap M \subseteq B$. Here, and elsewhere in the paper, we consider inclusions induced by cut-downs. Recall that if $Q \subseteq N$ is an inclusion of von Neumann algebras and q is a projection in Q , then

$$(Q' \cap N)q = (qQq)' \cap (qNq), \quad Z((Q' \cap N)q) = Z((qQq)' \cap (qNq)), \tag{2.14}$$

see, for example, [18, Section 5.4].

Lemma 2.6. *Let $B \subseteq M$ be a containment of finite von Neumann algebras such that $B' \cap M \subseteq B$. If $p \in M$ is a nonzero projection, then there exists a nonzero projection $q \in B$ which is equivalent to a subprojection of p .*

Observe that if M is a finite factor, then Lemma 2.6 is immediate. Our proof of Lemma 2.6 is classical, proceeding by analysing the center-valued trace on P . Alternatively one can establish the lemma by taking a direct integral over the center. Since we have been unable to find this fact in the literature we give the details for completeness.

Proof of Lemma 2.6. Let Δ denote the center-valued trace on M . We will make use of two properties of Δ from [12, Theorem 8.4.3]. The first is that $p_1 \preceq p_2$ if and only if $\Delta(p_1) \leq \Delta(p_2)$, and the second is that $p_1 \sim p_2$ if and only if $\Delta(p_1) = \Delta(p_2)$.

The hypothesis $B' \cap M \subseteq B$ implies that $B' \cap M = Z(B)$ and, in particular, that $Z(M) \subseteq Z(B)$. For some sufficiently small $c > 0$, the spectral projection z of $\Delta(p)$ for the interval $[c, 1]$ is nonzero, and $\Delta(pz) \geq cz$. Since $Bz \subseteq Mz$ also satisfies the relative commutant hypothesis, it suffices to prove the result under the additional restriction $\Delta(p) \geq c1$ for some constant $c > 0$.

Let $n \geq c^{-1}$ be any integer. Suppose that it is possible to find a nonzero projection $q \in B$ and an orthogonal set $\{q, p_2, \dots, p_n\}$ of equivalent projections in M . The sum of these projections has central trace equal to $n\Delta(q)$ and is also bounded by 1, so that $\Delta(q) \leq n^{-1}1 \leq$

c1. But then $q \lesssim p$ and we are done. Thus we may assume that there is an absolute bound on the length of any such set, and we may then choose one, $\{q_1, p_2, \dots, p_n\}$, of maximal length. By cutting by the central support of q_1 , we may assume that this central support is 1.

Now consider the inclusion $q_1 B q_1 \subseteq q_1 M q_1$, and note that

$$(q_1 B q_1)' \cap q_1 M q_1 = q_1 (B' \cap M) = q_1 Z(B) = Z(q_1 B q_1) \quad (2.15)$$

from (2.14). Let f_1 and f_2 be non-zero orthogonal projections in $q_1 B q_1$ and $q_1 M q_1$ respectively. By the comparison theory of projections, there exists a projection $z \in Z(q_1 M q_1) \subseteq Z(q_1 B q_1)$ so that

$$z f_1 \lesssim z f_2, \quad (1 - z) f_2 \lesssim (1 - z) f_1, \quad (2.16)$$

the equivalence being taken in $q_1 M q_1$. Now $z f_1 \in q_1 B q_1$ and is equivalent to a subprojection p_0 of $z f_2 \leq q_1$. Then the pair $z f_1, p_0$ is equivalent to orthogonal pairs below each p_i , $2 \leq i \leq n$, which will contradict the maximal length of $\{q_1, p_2, \dots, p_n\}$ unless $z f_1 = 0$. Similarly $(1 - z) f_2 = 0$. Thus f_1 and f_2 have orthogonal central supports in $q_1 B q_1$ and so [18, Lemma 5.5.3] shows that $q_1 B q_1$ is abelian. Equation (2.15) then shows that $q_1 B q_1$ is a masa in $q_1 M q_1$, and so another application of [18, Lemma 5.5.3] shows that $q_1 M q_1$ is also abelian. Thus $q_1 B q_1 = q_1 M q_1$.

Now the projection $1 - q_1 - p_2 \cdots - p_n$ must be 0, otherwise it would have a non-zero subprojection equivalent to a nonzero projection $\tilde{q}_1 \in q_1 M q_1 = q_1 B q_1$, since q_1 has central support 1, and \tilde{q}_1 would lie in a set of $n + 1$ equivalent orthogonal projections. Thus q_1, p_2, \dots, p_n are abelian projections in M with sum 1, so M is isomorphic to $L^\infty(\Omega) \otimes \mathbb{M}_n$ for some measure space Ω . Identify p and q_1 with measurable \mathbb{M}_n -valued functions. Since q_1 is abelian, the rank of $q_1(\omega)$ is 1 almost everywhere, and the rank of $p(\omega)$ is at least 1 almost everywhere since $\Delta(p) \geq c1$. Then q_1 is equivalent to a subprojection of p since $\Delta(q_1) \leq \Delta(p)$. This completes the proof. \square

We conclude this section with a brief explanation of an averaging technique in finite von Neumann algebras which we will use subsequently. It has its origins in [2], but is also used extensively in [14]. If $\eta \in L^2(M)$ and \mathcal{U} is a group of unitaries in M then the vector can be averaged over \mathcal{U} . This is normally associated with amenable groups, but can be made to work in this setting without this assumption. Form the $\|\cdot\|_2$ -norm closure \overline{K} of

$$K = \text{conv}\{u\eta u^* : u \in \mathcal{U}\}.$$

There is a unique vector $\tilde{\eta} \in \overline{K}$ of minimal norm, and uniqueness of $\tilde{\eta}$ implies that $u\tilde{\eta}u^* = \tilde{\eta}$ for all $u \in \mathcal{U}$. We refer to $\tilde{\eta}$ as the result of averaging η over \mathcal{U} , and many variations of this are possible. We give an example of this technique by establishing a technical result which will be needed in the proof of Theorem 4.7.

Recall (Proposition 2.1, (xii)) that $M e_B M$ is $\|\cdot\|_{2, \text{Tr}}$ -dense in $L^2(\langle M, e_B \rangle, \text{Tr})$. Consider a $*$ -subalgebra A which is strongly dense in M . If $x, y \in M$, then fix sequences $\{x_n\}_{n=1}^\infty, \{y_n\}_{n=1}^\infty$ from A converging strongly to x and y , respectively. Then

$$\begin{aligned} \|(x - x_n) e_B\|_{2, \text{Tr}}^2 &= \text{Tr}(e_B (x - x_n)^* (x - x_n) e_B) \\ &= \tau((x - x_n)^* (x - x_n)) = \langle (x - x_n) \xi, (x - x_n) \xi \rangle, \end{aligned} \quad (2.17)$$

so $x_n e_B \rightarrow x e_B$ in $\|\cdot\|_{2, \text{Tr}}$ -norm. Thus $x_n e_B y \rightarrow x e_B y$ so, given $\varepsilon > 0$, we may choose n_0 so large that $\|x_{n_0} e_B y - x e_B y\|_{2, \text{Tr}} < \varepsilon/2$. The same argument on the right allows us to choose n_1 so large that $\|x_{n_0} e_B y - x_{n_0} e_B y_{n_1}\|_{2, \text{Tr}} < \varepsilon/2$, whereupon $\|x e_B y - x_{n_0} e_B y_{n_1}\|_{2, \text{Tr}} < \varepsilon$. The conclusion reached is that the algebra $A e_B A = \{\sum_{i=1}^n x_i e_N y_i : x_i, y_i \in A\}$ is $\|\cdot\|_{2, \text{Tr}}$ -norm dense in $L^2(\langle M, e_B \rangle, \text{Tr})$. In the next lemma, we will use this when M is a tensor product $M_1 \overline{\otimes} M_2$ where we take A to be the algebraic tensor product $M_1 \otimes M_2$.

Lemma 2.7. *Let B_1, B_2 be von Neumann subalgebras of finite von Neumann algebras M_1, M_2 and let $B = B_1 \overline{\otimes} B_2$, $M = M_1 \overline{\otimes} M_2$. Then*

$$L^2(Z(B' \cap \langle M, e_B \rangle), \text{Tr}) = L^2(Z(B'_1 \cap \langle M_1, e_{B_1} \rangle), \text{Tr}_1) \otimes_2 L^2(Z(B'_2 \cap \langle M_2, e_{B_2} \rangle), \text{Tr}_2).$$

Note that, although Tr_i is a semifinite trace on M_i , it need not be semifinite on $Z(B'_i \cap \langle M_i, e_{B_i} \rangle)$. This is why the lemma cannot be obtained immediately from the uniqueness of product measures on σ -finite measure spaces.

Proof of Lemma 2.7. If $z_i \in Z(B'_i \cap \langle M_i, e_{B_i} \rangle)$, $i = 1, 2$, then $z_1 \otimes z_2 \in Z(B' \cap \langle M, e_B \rangle)$ and $\|z_1 \otimes z_2\|_{2, \text{Tr}} = \|z_1\|_{2, \text{Tr}_1} \|z_2\|_{2, \text{Tr}_2}$. This shows the containment from right to left.

Suppose that $z \in Z(B' \cap \langle M, e_B \rangle)$ with $\text{Tr}(z^* z) < \infty$. Then z lies in $L^2(\langle M, e_B \rangle, \text{Tr})$ so can be approximated in $\|\cdot\|_{2, \text{Tr}}$ -norm by sums of the form $\sum_{i=1}^k x_i e_B y_i$ with $x_i, y_i \in M_1 \overline{\otimes} M_2$. The preceding remarks then allow us to assume that x_i and y_i lie in the algebraic tensor product $M_1 \otimes M_2$. Thus, given $\varepsilon > 0$, we may find elements $a_i, c_i \in M_1$, $b_i, d_i \in M_2$ so that

$$\left\| z - \sum_{i=1}^n (a_i \otimes b_i)(e_{B_1} \otimes e_{B_2})(c_i \otimes d_i) \right\|_{2, \text{Tr}} \leq \varepsilon. \quad (2.18)$$

This may be rewritten as

$$\left\| z - \sum_{i=1}^n (a_i e_{B_1} c_i) \otimes (b_i e_{B_2} d_i) \right\|_{2, \text{Tr}} \leq \varepsilon, \quad (2.19)$$

and then as

$$\left\| z - \sum_{i=1}^n f_i \otimes g_i \right\|_{2, \text{Tr}} \leq \varepsilon, \quad (2.20)$$

where $f_i \in M_1 e_{B_1} M_1$ and $g_i \in M_2 e_{B_2} M_2$. We may further suppose that the set $\{g_1, \dots, g_n\}$ is linearly independent.

For $j = 1, 2$, let N_j be the von Neumann algebra generated by B_j and $B'_j \cap \langle M_j, e_{B_j} \rangle$, and note that z commutes with $N_1 \overline{\otimes} N_2$. Let

$$K = \overline{\text{conv}}^w \left\{ \sum_{i=1}^n u f_i u^* \otimes g_i : u \in \mathcal{U}(N_1) \right\}, \quad K_i = \overline{\text{conv}}^w \{ u f_i u^* : u \in \mathcal{U}(N_1) \}$$

for $1 \leq i \leq n$. Then $K \subseteq \sum_{i=1}^n K_i \otimes g_i$. By [18, Lemma 9.2.1] K and each K_i are closed in their respective $\|\cdot\|_{2, \text{Tr}}$ -norms. If $k \in K$ is the element of minimal $\|\cdot\|_{2, \text{Tr}}$ -norm in K then it may

be written as $k = \sum_{i=1}^n k_i \otimes g_i$ with $k_i \in K_i$. Since k is invariant for the action of $\mathcal{U}(N_1 \otimes 1)$, we see that

$$\sum_{i=1}^n (uk_i u^* - k_i) \otimes g_i = 0, \quad u \in \mathcal{U}(N_1). \quad (2.21)$$

The linear independence of the g_i 's allows us to conclude that $uk_i u^* = k_i$ for $1 \leq i \leq n$ and $u \in \mathcal{U}(N_1)$. Thus $k_i \in N'_1 \cap \langle M_1, e_{B_1} \rangle = Z(B'_1 \cap \langle M_1, e_{B_1} \rangle)$. The inequality (2.20) is preserved by averaging in this manner over $\mathcal{U}(N_1 \otimes 1)$ so, replacing each f_i by k_i if necessary, we may assume that $f_i \in Z(B'_1 \cap \langle M_1, e_{B_1} \rangle)$ for $1 \leq i \leq n$. Now repeat this argument on the right, averaging over $\mathcal{U}(1 \otimes N_2)$, to replace the g_i 's by elements of $Z(B'_2 \cap \langle M_2, e_{B_2} \rangle)$. With these changes, (2.20) now approximates z by a sum from

$$L^2(Z(B'_1 \cap \langle M_1, e_{B_1} \rangle), \text{Tr}_1) \otimes_2 L^2(Z(B'_2 \cap \langle M_2, e_{B_2} \rangle), \text{Tr}_2)$$

which proves the containment from left to right and establishes equality. \square

3 Projections in the basic construction

In this section, we relate intertwiners of a subalgebra to certain projections in the basic construction. We consider a finite von Neumann algebra M and a von Neumann subalgebra B whose unit will always coincide with that of M . For the most part, we will be interested in the condition $B' \cap M \subseteq B$ (equivalent to $B' \cap M = Z(B)$), but we will make this requirement explicit when it is needed.

Lemma 3.1. *Let B be a von Neumann subalgebra of a finite von Neumann algebra M and let $v \in \mathcal{GN}^{(1)}(B)$.*

1. *Then $v^* e_B v$ is a projection in $(B' \cap \langle M, e_B \rangle) v^* v$.*
2. *Suppose q is a projection in B . Then $v^* e_B v$ lies in $(B' \cap \langle M, e_B \rangle) q$ if, and only if, $v^* v \in Z(B) q = Z(q B q)$.*

Proof. 1. The element $v^* e_B v$ is positive in $\langle M, e_B \rangle$. Since $vv^* \in B$ and so commutes with e_B , the following calculation establishes that $v^* e_B v$ is a projection:

$$(v^* e_B v)^2 = v^* e_B v v^* e_B v = v^* v v^* e_B v = v^* e_B v. \quad (3.1)$$

For an arbitrary $b \in v^* v B v^* v$,

$$(v^* e_B v) b = v^* e_B v b v^* v = v^* v b v^* e_B v = v^* v b v^* (v v^*) e_B v = b (v^* e_B v), \quad (3.2)$$

where the second equality uses $v b v^* \in B$ to commute this element with e_B . Thus (3.2) establishes that $v^* e_B v \in (v^* v B v^* v)' \cap v^* v \langle M, e_B \rangle v^* v$ which is $(B' \cap \langle M, e_B \rangle) v^* v$ (see (2.14)).

2. Suppose now that $v^* v \in Z(B) q = Z(q B q)$. It is immediate that $Z(q B q) \subseteq Z((B' \cap \langle M, e_B \rangle) q)$, so we have a decomposition

$$(B' \cap \langle M, e_B \rangle) q = (B' \cap \langle M, e_B \rangle) v^* v \oplus (B' \cap \langle M, e_B \rangle) (q - v^* v). \quad (3.3)$$

We have already shown that v^*e_Bv is in the first summand of (3.3) so must lie in $(B' \cap \langle M, e_B \rangle)q$.

Conversely, the hypothesis on v^*e_Bv implies that $v^*e_Bv = v^*e_Bvq = qv^*e_Bv$, so the pull down map gives $v^*v = v^*vq = qv^*v$, showing that $v^*v \in qBq$. For each $b \in B$, $v^*e_Bvqbq = qbqv^*e_Bv$. Applying the pull down map gives $v^*vqbq = qbqv^*v$ and hence $v^*v \in Z(qBq)$. \square

We now strengthen this lemma under the additional hypothesis that $B' \cap M \subseteq B$. Recall from part (iv) of Proposition 2.1 that $JZ(B)J = Z(\langle M, e_B \rangle)$. When $B \subset M$ is a finite index inclusion of irreducible subfactors, Lemma 3.2 is contained in [13, Proposition 1.7 (2) and Proposition 1.9]. The proof follows the extension to infinite index inclusions of irreducible subfactors in [20, Lemma 3.3].

Lemma 3.2. *Let B be a von Neumann subalgebra of a finite von Neumann algebra M and suppose that $B' \cap M \subseteq B$. Let $v \in \mathcal{GN}^{(1)}(B)$. Then the projection v^*e_Bv is central in $(B' \cap \langle M, e_B \rangle)v^*v$.*

Proof. Define two projections $p, q \in B$ by $p = v^*v$ and $q = vv^*$. Now consider an arbitrary $x \in (B' \cap \langle M, e_B \rangle)p = (pBp)' \cap p\langle M, e_B \rangle p$. Then, for each $b \in B$,

$$(v xv^*)(v bv^*) = v x p b p v^* = v p b p x v^* = (v bv^*)(v xv^*), \quad (3.4)$$

showing that $v xv^* \in (vBv^*)' \cap (q\langle M, e_B \rangle q)$. We next prove that qe_B is central in $(vBv^*)' \cap (q\langle M, e_B \rangle q)$. It lies in this algebra by the previous calculation and Lemma 3.1, as $qe_B = v(v^*e_Bv)v^*$.

Take $t \in (vBv^*)' \cap (q\langle M, e_B \rangle q)$ to be self-adjoint and let η be $t\xi \in L^2(M)$. Now take a sequence $\{x_n\}_{n=1}^\infty$ from M converging in $\|\cdot\|_2$ -norm to $t\xi$. Since $t = qt = tq$, we may assume that the sequence $\{x_n\}_{n=1}^\infty$ lies in qMq , otherwise replace it by $\{qx_nq\}_{n=1}^\infty$.

For each u in the unitary group $\mathcal{U}(pBp)$, t commutes with $vu v^*$ and so

$$\begin{aligned} Jvuv^*Jvuv^*\eta &= Jvuv^*Jvuv^*t\xi = Jvuv^*Jtvuv^*\xi \\ &= tJvuv^*Jvuv^*\xi = tvuv^*vu^*v^*\xi \\ &= tq\xi = t\xi = \eta, \end{aligned} \quad (3.5)$$

where the third equality holds because $vu v^* \in B$ so that $Jvuv^*J \in (\langle M, e_B \rangle)'$. For each $n \geq 1$ and each $u \in \mathcal{U}(pBp)$,

$$\begin{aligned} \|Jvuv^*Jvuv^*x_n\xi - \eta\|_2 &= \|Jvuv^*Jvuv^*(x_n\xi - \eta)\|_2 \\ &\leq \|x_n\xi - \eta\|_2, \end{aligned} \quad (3.6)$$

from (3.5). If we let y_n be the element of qMq obtained by averaging x_n over the unitary group $v\mathcal{U}(pBp)v^* \subseteq qBq$, then (3.6) gives

$$\|y_n\xi - \eta\|_2 \leq \|x_n\xi - \eta\|_2, \quad n \geq 1, \quad (3.7)$$

while $y_n \in (vBv^*)' \cap qMq$. Since

$$\begin{aligned} (vBv^*)' \cap qMq &= (vBv^*)' \cap (vMv^*) = v(B' \cap M)v^* \\ &= vZ(B)v^* \subseteq qBq, \end{aligned} \quad (3.8)$$

we see that $y_n \in qBq$ for $n \geq 1$. From (3.7) it follows that $\eta \in L^2(qBq)$. For $b \in B$,

$$\begin{aligned} tqb\xi &= tJb^*qJ\xi = Jb^*qJt\xi \\ &= \lim_{n \rightarrow \infty} Jb^*qJy_n\xi = \lim_{n \rightarrow \infty} y_nqb\xi, \end{aligned} \quad (3.9)$$

showing that $tqb\xi \in L^2(qB)$. Thus $L^2(qB)$ is an invariant subspace for t . The projection onto it is qe_B , so $tqe_B = qe_Btqe_B$. Since t is self-adjoint, we obtain that qe_B commutes with $(vBv^*)' \cap (q\langle M, e_B \rangle q)$, establishing centrality.

It was established in equation (3.4) that $v xv^* \in (vBv^*)' \cap (q\langle M, e_B \rangle q)$ whenever $x \in (B' \cap \langle M, e_B \rangle)p$, and so each such $v xv^*$ commutes with qe_B . Thus

$$\begin{aligned} v^*e_B v x &= v^*q e_B v x v^* v = v^* v x v^* q e_B v \\ &= x v^* e_B v \end{aligned} \quad (3.10)$$

for $x \in (B' \cap \langle M, e_B \rangle)p$, showing that $v^*e_B v$ is central in $(B' \cap \langle M, e_B \rangle)p$. \square

Since the centrality of the projections $v^*e_B v$ in $(B' \cap \langle M, e_B \rangle)v^*v$ is crucial to our subsequent arguments, the following corollary highlights why the hypothesis $B' \cap M \subseteq B$ is essential.

Corollary 3.3. *Let B be a von Neumann subalgebra of a finite von Neumann algebra M . Then e_B is central in $B' \cap \langle M, e_B \rangle$ if and only if $B' \cap M \subseteq B$.*

Proof. If $B' \cap M \subseteq B$, then centrality of e_B is a special case of Lemma 3.2 (i) with $v = 1$. Conversely, suppose that e_B is central in $B' \cap \langle M, e_B \rangle$ and consider $x \in B' \cap M$. Then x commutes with e_B so $x \in B$ by Proposition 2.1 (iii). Thus $B' \cap M \subseteq B$. \square

Lemma 3.4. *Let B be a von Neumann subalgebra of a finite von Neumann algebra M , suppose that $B' \cap M \subseteq B$, and let $v \in \mathcal{GN}^{(1)}(B)$. Then any subprojection of $v^*e_B v$ in $(B' \cap \langle M, e_B \rangle)v^*v$ has the form $pv^*e_B v$ where p is a central projection in B .*

Proof. By Lemma 3.1 (i), $v^*e_B v \in (B' \cap \langle M, e_B \rangle)v^*v$. Suppose that a projection $q \in (B' \cap \langle M, e_B \rangle)v^*v$ lies below $v^*e_B v$. Then $(vqv^*)^2 = vqv^*vqv^* = vqv^*$, so vqv^* is a projection. The relation

$$(vqv^*)(v bv^*) = vq(v^*v bv^*v)v^* = v(v^*v bv^*v)qv^* = (v bv^*)(vqv^*), \quad b \in B, \quad (3.11)$$

shows that $vqv^* \in (vBv^*)' \cap vv^*\langle M, e_B \rangle vv^*$. Moreover, $vqv^* \leq vv^*e_B vv^* = e_B vv^*$, so there exists a projection $f \in vv^*Bvv^*$ such that $vqv^* = fe_B$. For $b \in B$,

$$f v b v^* e_B = f e_B v b v^* = v q v^* v b v^* = v b v^* v q v^* = v b v^* f e_B, \quad (3.12)$$

and so $f v b v^* = v b v^* f$. Thus

$$f \in (vBv^*)' \cap vv^*Bvv^*. \quad (3.13)$$

If $b_0 \in B$ is such that $vv^*b_0vv^*$ commutes with vBv^* , then

$$vv^*b_0vv^*v b v^* = v b v^*vv^*b_0vv^*, \quad b \in B. \quad (3.14)$$

Multiply on the left by v^* and on the right by v to obtain

$$v^*b_0vv^*vbv^*v = v^*vbv^*vv^*b_0v, \quad b \in B. \quad (3.15)$$

Thus

$$v^*b_0v \in (v^*vBv^*v)' \cap v^*vMv^*v = v^*v(B' \cap M)v^*v = v^*vZ(B)v^*v. \quad (3.16)$$

Consequently, $vv^*b_0vv^* \in vZ(B)v^*$. It follows that $(vBv^*)' \cap vv^*Bvv^* \subseteq vZ(B)v^*$, and so there is a central projection $p \in Z(B)$ so that $f = vpv^*$. We now have

$$q = v^*fe_Bv = v^*vpv^*e_Bv = pv^*e_Bv, \quad (3.17)$$

as required. \square

In Section 4, we wish to use the projections v^*e_Bv to investigate an intertwiner v of a tensor product $B = B_1 \overline{\otimes} B_2 \subset M_1 \overline{\otimes} M_2 = M$, where each $B'_i \cap M_i \subseteq B_i$. In conjunction with Proposition 2.1 (xiii), Tomita's commutation theorem gives

$$(B' \cap \langle M, e_B \rangle) \cong (B'_1 \cap \langle M_1, e_{B_1} \rangle) \overline{\otimes} (B'_2 \cap \langle M_2, e_{B_2} \rangle). \quad (3.18)$$

By Lemma 3.2, such an intertwiner gives rise to a central projection v^*e_Bv in $(B' \cap \langle M, e_B \rangle)v^*v$. Unfortunately, in general the projection v^*v will not factorise as an elementary tensor of projections $b_1 \otimes b_2$, with $b_i \in B_i$, and so the algebra $(B' \cap \langle M, e_B \rangle)v^*v$ will not decompose as a tensor product. This prevents us from applying tensor product techniques to the projection v^*e_Bv directly. However, standard von Neumann algebra theory (see for example [11]) gives a central projection $P \in B' \cap \langle M, e_B \rangle$ such that $v^*e_Bv = Pv^*v$. Since we need to ensure that the projection P fully reflects the properties of v , we cannot just appeal to the general theory to obtain P so we give an explicit construction in Definition 3.5 below. The subsequent lemmas set out the properties of P that we require later.

Definition 3.5. Let $B \subset M$ be an inclusion of finite von Neumann algebras with $B' \cap M \subseteq B$ and let $v \in \mathcal{GN}_M^{(1)}(B)$. Let $z \in Z(B)$ be the central support of v^*v . Define p_0 to be v^*v , and let $\{p_0, p_1, \dots\}$ be a family of nonzero pairwise orthogonal projections in B which is maximal with respect to the requirements that $p_n \leq z$ and each p_n is equivalent in B to a subprojection in B of p_0 . Since two projections in a von Neumann algebra with non-orthogonal central supports have equivalent non-zero subprojections, maximality gives $\sum_{n \geq 0} p_n = z$. For $n \geq 1$, choose partial isometries $w_n \in B$ so that $w_n^*w_n = q_n \leq p_0$ and $w_nw_n^* = p_n$. Then define $v_n = vw_n^* \in \mathcal{GN}^{(1)}(B)$. Lemma 3.1 shows that $v_n^*e_Bv_n \in (B' \cap \langle M, e_B \rangle)v_n^*v_n$ and this space is $(B' \cap \langle M, e_B \rangle)p_n$ since $v_n^*v_n = w_nv^*vw_n^* = p_n$. In particular, $\{v_n^*e_Bv_n\}_{n \geq 0}$ is a set of pairwise orthogonal projections so we may define a projection $P_v = \sum_{n \geq 0} v_n^*e_Bv_n$ in $\langle M, e_B \rangle$.

Lemma 3.6. *With the notation of Definition 3.5, the projection P_v is central in $B' \cap \langle M, e_B \rangle$ and satisfies $P_vv^*v = v^*e_Bv$.*

Proof. This projection is $P_v = \sum_{n \geq 0} w_nv^*e_Bvw_n^*$. By Lemma 3.2 there exists $t \in Z(B' \cap \langle M, e_B \rangle)$ so that $v^*e_Bv = tv^*v$, and so P_v becomes

$$P_v = \sum_{n \geq 0} w_nv^*vw_n^* = \sum_{n \geq 0} tw_nv^*vw_n^* = \sum_{n \geq 0} tw_np_0w_n^* = tz. \quad (3.19)$$

Thus $P_v \in B' \cap \langle M, e_B \rangle$ and $P_v v^* v = t v^* v = v^* e_B v$ since z is the central support of $v^* v$. Since $z \in Z(B) \subset Z(B' \cap \langle M, e_B \rangle)$, it follows that $P_v = tz$ also lies in $Z(B' \cap \langle M, e_B \rangle)$ \square

Remark 3.7. The proof of Lemma 3.6 shows that P_v is the minimal projection in $B' \cap \langle M, e_B \rangle$ with $P_v v^* v = v^* e_B v$. This gives a canonical description of P_v which is independent of the choices made in Definition 3.5. The explicit formulation of the definition is useful in transferring properties from $v^* e_B v$ to P_v .

We now identify the subprojections of P_v . This will be accomplished by the next lemma, which considers a wider class of projections needed subsequently. Let $\{v_i\}_{i=1}^{\infty}$ be a sequence from $\mathcal{GN}^{(1)}(B)$ satisfying $v_i v_j^* = 0$ for $i \neq j$, let $p \in B$ be the projection $\sum_{i=1}^{\infty} v_i^* v_i$ and let $P \in \langle M, e_B \rangle$ be the projection $\sum_{i=1}^{\infty} v_i^* e_B v_i$. In particular, the projection P_v of Definition 3.5 is of this form. Let $N(P)$ denote the von Neumann algebra

$$N(P) = \{x \in pMp : xP = Px\} \subseteq M. \quad (3.20)$$

Lemma 3.8. Let $P = \sum_{i=1}^{\infty} v_i^* e_B v_i$ be as above, and let $p = \sum_{i=1}^{\infty} v_i^* v_i \in B$.

- (i) If $x \in pMp$ satisfies $xP = 0$, then $x = 0$;
- (ii) The map $x \rightarrow xP$ is a $*$ -isomorphism of $N(P)$ into $\langle M, e_B \rangle$;
- (iii) A projection $Q \in \langle M, e_B \rangle$ satisfies $Q \leq P$ if and only if there exists a projection $f \in N(P)$ such that $Q = fP$. Moreover, if P and Q lie in $B' \cap \langle M, e_B \rangle$, then $f \in Z(B)$ and Q has the same form as P .

Proof. (i) Suppose that $x \in pMp$ and $xP = 0$. Then

$$x \sum_{i=1}^{\infty} v_i^* e_B v_i = 0. \quad (3.21)$$

Multiply on the right in (3.21) by $v_k^* v_k$ to obtain $x v_k^* e_B v_k = 0$ for $k \geq 1$. The pull down map gives $x v_k^* v_k = 0$. Summing over k shows that $xp = 0$ and the result follows since $x = xp$.

(ii) Since $P \in N(P)'$, the map $x \mapsto xP$ is a $*$ -homomorphism on $N(P)$. It has trivial kernel, by (i), so is a $*$ -isomorphism.

(iii) If $f \in N(P)$, then it is clear that fP is a projection below P , since f commutes with P . Conversely, consider a projection $Q \leq P$ with $Q \in \langle M, e_B \rangle$. The introduction of partial sums below is to circumvent some questions of convergence.

Define $P_k = \sum_{i=1}^k v_i^* e_B v_i$. Then $\lim_{k \rightarrow \infty} P_k = P$ strongly, so $P_k Q P_k$ converges strongly to $P Q P = Q$. For $m, n \geq 1$, let $b_{m,n} \in B$ be the element such that $b_{m,n} e_B = e_B v_m Q v_n^* e_B$. Then

$$\begin{aligned} P_k Q P_k &= \sum_{m,n=1}^k v_m^* e_B v_m Q v_n^* e_B v_n = \sum_{m,n=1}^k v_m^* b_{m,n} e_B v_n \\ &= \sum_{m,n=1}^k v_m^* b_{m,n} e_B v_n v_n^* v_n = \sum_{m,n=1}^k v_m^* b_{m,n} v_n v_n^* e_B v_n. \end{aligned} \quad (3.22)$$

Now define $x_k \in pW^*(\mathcal{GN}^{(1)}(B))p$ by $x_k = \sum_{m,n=1}^k v_m^* b_{m,n} v_n$. The relations $v_i v_j^* = 0$ for $i \neq j$ allow us to verify that $x_k P = P_k Q P_k$, and consequently

$$x_k P = P x_k^*, \quad k \geq 1, \quad (3.23)$$

since $P_k Q P_k$ is self-adjoint. Thus

$$\sum_{i=1}^{\infty} x_k v_i^* e_B v_i = \sum_{i=1}^{\infty} v_i^* e_B v_i x_k^*. \quad (3.24)$$

The sums in (3.24) converge in $\|\cdot\|_1$ -norm, so we may apply the pull down map to obtain $x_k p = p x_k^*$. Since $x_k = x_k p$, we conclude that x_k is self-adjoint. Thus, from (3.23), x_k commutes with P , and so lies in $N(P)$. From above, $x_k P \geq 0$ and $\|x_k P\| \leq 1$, so $x_k \geq 0$ and $\|x_k\| \leq 1$ by (ii). Let f be a σ -weak accumulation point of the sequence $\{x_k\}_{k=1}^{\infty}$. Then $f \geq 0$, $\|f\| \leq 1$, and $f \in N(P)$. Since $x_k P = P_k Q P_k$, we conclude that $f P = Q$. It now follows from (ii) that f is a projection in $N(P)$.

If $P \in B' \cap \langle M, e_B \rangle$, then the pull down map gives $bp = pb$ for $b \in B$, so $p \in Z(B)$. If also $Q \in B' \cap \langle M, e_B \rangle$, then $Q = fP$ and $f \leq p$. If $b \in Bp$ then commutation with Q gives $(bf - fb)P = 0$, so $f \in B' \cap M = Z(B)$, by (i). Finally $Q = fP = \sum_i (v_i f)^* e_B (v_i f)$, so is of the same form as P . \square

4 Intertwiners of tensor products

In this section we will prove one of our main results, the equality of $W^*(\mathcal{GN}^{(1)}(B_1)) \overline{\otimes} W^*(\mathcal{GN}^{(1)}(B_2))$ and $W^*(\mathcal{GN}^{(1)}(B_1 \overline{\otimes} B_2))$, where $B_i \subseteq M_i$, $i = 1, 2$, are inclusions of finite von Neumann algebras satisfying $B_i' \cap M_i \subseteq B_i$. The key theorem for achieving this is the following one, which enables us to detect those central projections in corners of the relative commutant of the basic construction which arise from intertwiners. It is inspired by [1, Proposition 2.7], although is not a direct generalization of that result. For comparison, [1, Proposition 2.7] shows that, in the case of a masa A , a projection $P \in A' \cap \langle M, e_A \rangle$ which is subequivalent to e_A dominates an operator $v^* e_B v$ for some $v \in \mathcal{GN}(A)$. Example 4.3 below will show that such a result will not hold in general without additional hypotheses.

Theorem 4.1. *Let A be an abelian von Neumann algebra with a fixed faithful normal semifinite weight Φ . Let B be a von Neumann subalgebra of a finite von Neumann algebra M with a faithful normal trace τ satisfying $B' \cap M \subseteq B$. Fix a projection $q \in A \overline{\otimes} B$ and suppose that $P \in (A \overline{\otimes} (B' \cap \langle M, e_B \rangle))q$ is a nonzero projection such that $P \lesssim (1 \otimes e_B)$ in $A \overline{\otimes} \langle M, e_B \rangle$, and satisfies*

$$(\Phi \otimes \text{Tr})(Pr) \leq (\Phi \otimes \tau)(qr) \quad (4.1)$$

for all projections $r \in Z(q(A \overline{\otimes} B)q)$. Then there exists an element $v \in \mathcal{GN}_{A \overline{\otimes} M}^{(1)}(A \overline{\otimes} B)$ such that $P = v^*(1 \otimes e_B)v$.

Before embarking on the proof, let us recall that, for a finite von Neumann algebra M with a faithful normal trace τ , we regard the Hilbert space $L^2(M)$ as the completion of M in the

norm $\|x\|_{2,\tau} = \tau(x^*x)^{1/2}$ and $L^1(M)$ as the completion of M in the norm $\|x\|_{1,\tau} = \tau(|x|)$. The Cauchy-Schwarz inequality gives $\|xy^*\|_{1,\tau} \leq \|x\|_{2,\tau}\|y\|_{2,\tau}$ for $x, y \in M$ and so this inequality allows us to define $\zeta\eta^* \in L^1(M)$ for $\zeta, \eta \in L^2(M)$. In particular, if (y_n) is a sequence in M converging to $\eta \in L^2(M)$, then $y_n^*y_n \rightarrow \eta^*\eta$ in $L^1(M)$.

Recall too that we can regard elements of $L^2(M)$ as unbounded operators on $L^2(M)$ affiliated to M . The only fact we need about these unbounded operators is that if $\eta \in L^2(M)$ satisfies $\eta^*\eta \in M$ (regarded as a subset of $L^1(M)$), then in fact $\eta \in M$. This follows as η has a polar decomposition $v(\eta^*\eta)^{1/2}$, where v is a partial isometry in M and $(\eta^*\eta)^{1/2}$ is an element of $L^2(M)$, which lies in M if $\eta^*\eta$ does.

Proof of Theorem 4.1. The first case that we will consider is where $A = \mathbb{C}$ and Φ is the identity map. Then the hypothesis becomes

$$\text{Tr}(Pr) \leq \tau(qr) \quad (4.2)$$

for all projections $r \in Z(B)q$. Since $P \preceq e_B$, there exists a partial isometry $V \in \langle M, e_B \rangle$ such that $P = V^*V$ and $VV^* \leq e_B$. Define the map $\theta: qBq \rightarrow Be_B$ by

$$\theta(qbq) = VqbqV^* = e_B VqbqV^* e_B, \quad b \in B. \quad (4.3)$$

Then θ is a $*$ -homomorphism since V^*V commutes with qBq , and so there is a $*$ -homomorphism $\phi: qBq \rightarrow B$ so that $\theta(qbq) = \phi(qbq)e_B$ for $qbq \in qBq$. Thus

$$qbqV^* = qbqV^*VV^* = V^*VqbqV^* = V^*e_B\phi(qbq) = V^*\phi(qbq) \quad (4.4)$$

for $qbq \in qBq$. Now define $\eta \in L^2(M)$ by $\eta = JV^*\xi$, and observe that $\eta \neq 0$ since $VJ\eta = VV^*\xi = b_0\xi$, where $VV^* = b_0e_B$ for some $b_0 \in B$. If we apply (4.4) to ξ , then the result is

$$qbqJ\eta = V^*\phi(qbq)\xi = V^*J\phi(qb^*q)J\xi = J\phi(qb^*q)JV^*\xi = J\phi(qb^*q)\eta, \quad qbq \in qBq, \quad (4.5)$$

where we have used $\langle M, e_B \rangle = (JBJ)'$ to commute $J\phi(qb^*q)J$ with V^* . Multiply (4.10) on the left by J and replace b by b^* to obtain

$$\eta qbq = \phi(qbq)\eta, \quad qbq \in qBq. \quad (4.6)$$

Taking $b = 1$, (4.5) becomes

$$JqJ\eta = \phi(q)\eta. \quad (4.7)$$

so

$$JqV^*\xi = \phi(q)\eta. \quad (4.8)$$

Multiply on the left by VJ to obtain

$$\phi(q)\xi = VJ\phi(q)\eta, \quad (4.9)$$

showing that $\phi(q)\eta \neq 0$. From (4.7), $\phi(q)\eta q = \phi(q)\eta \neq 0$, and this allows us to assume in (4.6) that the vector η is nonzero and satisfies $\eta q = \eta$, $\phi(q)\eta = \eta$ by replacing η with $\phi(q)\eta q$ if necessary. For unitaries $u \in qBq$, (4.6) becomes

$$\phi(u^*)\eta u = \phi(u^*)\phi(u)\eta = \phi(q)\eta = \eta. \quad (4.10)$$

Choose a sequence $\{x_n\}_{n=1}^\infty$ from M such that $x_n\xi \rightarrow \eta$ in $\|\cdot\|_{2,\tau}$ -norm. Since $\phi(q)\eta q = \eta$, we may assume that $\phi(q)x_n q = x_n$ for $n \geq 1$. Let y_n be the element of minimal $\|\cdot\|_{2,\tau}$ -norm in $\overline{\text{conv}}^w\{\phi(u^*)x_n u : u \in \mathcal{U}(qbq)\}$. Since

$$\|\phi(u^*)x_n u \xi - \eta\|_{2,\tau} = \|\phi(u^*)(x_n \xi - \eta)u\|_{2,\tau} \leq \|x_n \xi - \eta\|_{2,\tau} \quad (4.11)$$

for all $u \in \mathcal{U}(qBq)$, we see that $\|y_n \xi - \eta\|_{2,\tau} \leq \|x_n \xi - \eta\|_{2,\tau}$, so $y_n \xi \rightarrow \eta$ in $\|\cdot\|_{2,\tau}$ -norm and $\phi(u^*)y_n u = y_n$ for $n \geq 1$ by the choice of y_n . Then $y_n u = \phi(u)y_n$ for $u \in \mathcal{U}(qBq)$, so

$$y_n q b q = \phi(q b q) y_n, \quad n \geq 1, \quad q b q \in q B q. \quad (4.12)$$

Thus

$$y_n^* y_n q b q = y_n^* \phi(q b q) y_n, \quad n \geq 1, \quad q b q \in q B q, \quad (4.13)$$

and this implies that $y_n^* y_n \in (qBq)' \cap qMq = (B' \cap M)q = Z(B)q$ for each $n \geq 1$. The discussion preceding the proof ensures that $y_n^* y_n \rightarrow \eta^* \eta$ in $L^1(M)$, so we see that $\eta^* \eta \in L^1(Z(B)q)$. For each $z \in Z(B)q$,

$$\begin{aligned} |\tau(\eta^* \eta z q)| &= \lim_{n \rightarrow \infty} |\tau(y_n^* y_n q z)| = \lim_{n \rightarrow \infty} |\langle y_n q z \xi, y_n \xi \rangle| \\ &= |\langle J z^* q J \eta, \eta \rangle| = |\langle J z^* q J J V^* \xi, J V^* \xi \rangle| \\ &= |\langle z^* q V^* \xi, V^* \xi \rangle| = |\langle V z^* q V^* \xi, \xi \rangle| \\ &= |\langle \phi(z^* q) \xi, \xi \rangle| = |\tau(\phi(z q))|. \end{aligned} \quad (4.14)$$

Now

$$\begin{aligned} \text{Tr}(P z q) &= \text{Tr}(V^* V z q) = \text{Tr}(V z q V^*) \\ &= \text{Tr}(\phi(z q) e_B) = \tau(\phi(z q)). \end{aligned} \quad (4.15)$$

Thus, from (4.14), (4.15) and the hypothesis (4.2),

$$|\tau(\eta^* \eta r)| = |\text{Tr}(P r)| \leq \tau(r) \quad (4.16)$$

for all projections $r \in Z(B)q$. Since $Z(B)q$ is abelian, simple measure theory allows us to conclude from (4.16) that $\eta^* \eta \in Z(B)q$ (rather than just $L^1(Z(B)q)$) and so $\eta \in M$, by the discussion prior to the start of the proof. Moreover, (4.16) also gives $\|\eta\| \leq 1$, by taking r to be the spectral projection of $\eta^* \eta$ for the interval (c, ∞) where $c > 1$ is arbitrary.

Since $\eta \in L^2(M)$ has been proved to lie in M , we rename this nonzero operator as $x \in M$. From above, $\|x\| \leq 1$ and $x^* x \in Z(B)q$. Since $J V^* \xi = \eta = x \xi = x J \xi$, for $y \in M$ and $b \in B$,

$$\begin{aligned} \langle (V - x) y \xi, b \xi \rangle &= \langle y \xi, V^* J b^* J \xi \rangle - \langle x y \xi, b \xi \rangle \\ &= \langle y \xi, J b^* J V^* \xi \rangle - \langle x y \xi, b \xi \rangle \\ &= \langle y \xi, J b^* x J \xi \rangle - \langle x y \xi, b \xi \rangle \\ &= \langle y \xi, x^* b \xi \rangle - \langle x y \xi, b \xi \rangle = 0, \end{aligned} \quad (4.17)$$

and so $e_B V = e_B x$, implying that $V = e_B x$. Thus $P = V^* V = x^* e_B x$. Since $P^2 = P$,

$$x^* e_B x = x^* e_B x x^* e_B x = x^* \mathbb{E}_B(x x^*) e_B x, \quad (4.18)$$

so the pull down map gives

$$x^*x = x^*\mathbb{E}_B(xx^*)x. \quad (4.19)$$

This equation is $x^*(1 - \mathbb{E}_B(xx^*))x = 0$, so $(1 - \mathbb{E}_B(xx^*))^{1/2}x = 0$. Thus $\mathbb{E}_B(xx^*)x = x$. If we multiply on the right by x^* and apply \mathbb{E}_B , then we conclude that $\mathbb{E}_B(xx^*)$ is a projection. Moreover,

$$\mathbb{E}_B(xx^*)xx^* = xx^* \quad (4.20)$$

so $\mathbb{E}_B(xx^*) \geq xx^*$ since $\|x\| \leq 1$. The trace then gives equality, and so x is a partial isometry with $x^*x, xx^* \in B$. Since $x = xq$ and $x^*e_Bx = P$, which commutes with qBq , we obtain

$$xbx^*e_B = xqbqx^*xx^*e_B = xqbqx^*e_Bxx^* = xx^*e_Bxqbqx^* = e_Bxbx^*, \quad b \in B, \quad (4.21)$$

showing that $xBx^* \subseteq B$. Thus $x \in \mathcal{GN}^{(1)}(B)$ and $P = x^*e_Bx$. This completes the proof when $A = \mathbb{C}$ and Φ is the identity map.

The second case is when A is an arbitrary abelian von Neumann algebra, and Φ is bounded, so we may assume that Φ is a state on A since (4.1) is unaffected by scaling. The result now follows from the first case by replacing the inclusion $B \subseteq M$ by $A \overline{\otimes} B \subseteq A \overline{\otimes} M$. The trace on $A \overline{\otimes} M$ is $\Phi \otimes \tau$ and $e_{A \overline{\otimes} B}$ is $1 \otimes e_B$, so the canonical trace on $\langle A \overline{\otimes} M, e_{A \overline{\otimes} B} \rangle$ is $\Phi \otimes Tr$.

The last case is where Φ is a faithful normal semifinite weight. Then there is a family $\{f_\lambda\}_{\lambda \in \Lambda}$ of orthogonal projections in A with sum 1 such that $\Phi(f_\lambda) < \infty$ for each $\lambda \in \Lambda$. If Φ_λ denotes the restriction of Φ to Af_λ , then Φ_λ is bounded. If we replace P, A, Φ and q by respectively $P(f_\lambda \otimes 1), Af_\lambda, \Phi_\lambda$ and $q(f_\lambda \otimes 1)$, then we are in the second case. Thus there exists, for each $\lambda \in \Lambda$, a partial isometry $v_\lambda \in \mathcal{GN}_{Af_\lambda \overline{\otimes} M}^{(1)}(Af_\lambda \overline{\otimes} B)$ so that $P(f_\lambda \otimes 1) = v_\lambda^*(e_{Af_\lambda \overline{\otimes} B})v_\lambda$. The central support of v_λ lies below $f_\lambda \otimes 1$ so we may define $v \in \mathcal{GN}_{A \overline{\otimes} M}^{(1)}(A \overline{\otimes} B)$ by $v = \sum_{\lambda \in \Lambda} v_\lambda$, and it is routine to check that $P = v^*e_Bv$. \square

Theorem 4.1 characterizes those projections in the basic construction which arise from intertwiners.

Corollary 4.2. *Given an inclusion $B \subseteq M$ of finite von Neumann algebras with $B' \cap M \subseteq B$ and a projection $q \in B$, a projection $P \in (B' \cap \langle M, e_B \rangle)q$ is of the form v^*e_Bv for some intertwiner $v \in \mathcal{GN}^{(1)}(B)$ if and only if $P \lesssim e_B$ in $\langle M, e_B \rangle$ and $\text{Tr}(Pr) \leq \tau(qr)$ for all projections $r \in Z(B)q$. Furthermore in this case the domain projection v^*v must lie in $Z(B)q$.*

Proof. Taking $A = \mathbb{C}$ and Φ to be the identity in theorem 4.1 shows that any projection P satisfying the conditions of the corollary is of the form v^*e_Bv for some intertwiner $v \in \mathcal{GN}^{(1)}(B)$. Lemma 3.1 (ii) then shows that $v^*v \in Z(B)q$. Conversely, given an intertwiner $v \in \mathcal{GN}^{(1)}(B)$, Lemma 3.1 (ii) shows that $v^*e_Bv \in (B' \cap \langle M, e_B \rangle)q$ precisely when $v^*v \in Z(B)q$. The other two conditions of the corollary follow as $v^*e_Bv \sim vv^*e_B \leq e_B$ in $\langle M, e_B \rangle$, and for a projection $r \in Z(B)q$, $\text{Tr}(v^*e_Bvr) = \tau(v^*vr) \leq \tau(qr)$. \square

The tracial hypothesis (4.1) of Theorem 4.1 is an extra ingredient in this theorem as compared with [1, Proposition 2.7]. The following example shows that Theorem 4.1 can fail without this hypothesis.

Example 4.3. Let R be the hyperfinite II_1 factor and fix an outer automorphism θ of period two. Let $M = \mathbb{M}_3 \otimes R$ and let

$$B = \left\{ \begin{pmatrix} a & 0 & 0 \\ 0 & b & c \\ 0 & \theta(c) & \theta(b) \end{pmatrix} : a, b, c \in R \right\} \subseteq M.$$

Note that $B \cong R \oplus (R \rtimes_{\theta} \mathbb{Z}_2)$. It is straightforward to verify that $B' \cap M = Z(B) = \mathbb{C}e_{11} \oplus \mathbb{C}(e_{22} + e_{33})$ where $\{e_{i,j}\}_{i,j=1}^3$ are the matrix units. Let $P \in \langle M, e_B \rangle$ be $(\sqrt{2} e_{12})e_B(\sqrt{2} e_{21})$. Since $\mathbb{E}_B(e_{22}) = (e_{22} + e_{33})/2$, P is a projection, and it is routine to verify that P commutes with B . If there is a nonzero intertwiner $v \in M$ such that $v^*e_Bv \leq P$, then $v = ve_{11}$. Direct calculation shows that v would then have the form we_{11} for some partial isometry $w \in R$, so v^*e_Bv would be $qe_{11}e_B$ for some projection $q \in R$. However, this nonzero projection is orthogonal to P and so cannot lie under it. Thus the conclusion of Theorem 4.1 fails in this case. Note that $\text{Tr}(Pe_{11}) = \tau(2e_{11}) = 2/3$, while $\tau(e_{11}) = 1/3$, so the tracial hypothesis of Theorem 4.1 is not satisfied.

It is worth noting that $B' \cap \langle M, e_B \rangle$ can be explicitly calculated in this case. This algebra is abelian and five dimensional with minimal projections $e_{11}e_B$, $(1 - e_{11})e_B$, $(1 - e_{11})(1 - e_B)$, $(\sqrt{2} e_{12})e_B(\sqrt{2} e_{21})$, and $e_{21}e_Be_{12} + e_{31}e_Be_{13}$. The corresponding B -bimodules in $L^2(M)$ are generated respectively by the vectors e_{11} , $e_{22} + e_{33}$, $e_{22} - e_{33}$, e_{12} , and e_{21} . \square

For the remainder of the section we fix inclusions $B_i \subseteq M_i$ of finite von Neumann algebras satisfying $B'_i \cap M_i \subseteq B_i$ for $i = 1, 2$, and we denote the inclusion $B_1 \overline{\otimes} B_2 \subseteq M_1 \overline{\otimes} M_2$ by $B \subseteq M$. For $i = 1, 2$, let

$$S_i = \sup\{P_j \in Z(B'_i \cap \langle M_i, e_{B_i} \rangle) : \text{Tr}_i(P_j) < \infty\}.$$

Note that S_i acts as the identity on $L^2(Z(B'_i \cap \langle M_i, e_{B_i} \rangle), \text{Tr}_i)$. Given $v \in \mathcal{GN}_M^{(1)}(B)$, the projection P_v of Definition 3.5 satisfies $\text{Tr}(P_v) \leq 1$, and $P_v \in Z(B' \cap \langle M, e_B \rangle)$ by Lemma 3.6. It follows from Lemma 2.7 that $P_v \leq S_1 \otimes S_2$. Although Tr_i restricted to $Z(B'_i \cap \langle M_i, e_{B_i} \rangle)$ might not be semifinite, it does have this property on the abelian von Neumann algebra $A_i = Z(B'_i \cap \langle M_i, e_{B_i} \rangle)S_i$ by the choice of S_i . Moreover, each P_v is an element of $A_1 \overline{\otimes} \langle M_2, e_{B_2} \rangle$. We need two further projections which we define below.

Definition 4.4. For $i = 1, 2$, let \mathcal{P}_i denote the collection of projections $R \in Z(B'_i \cap \langle M_i, e_{B_i} \rangle)$ which are expressible as

$$R = \sum_{n \geq 1} v_n^* e_{B_i} v_n, \quad v_n \in \mathcal{GN}^{(1)}(B_i)$$

where $\{v_n^* v_n\}_{n \geq 1}$ is an orthogonal set of projections in B_i . Such a projection satisfies $\text{Tr}_i(R) \leq 1$. Let Q_i be the supremum of the projections in \mathcal{P}_i , so $Q_i \leq S_i$. \square

Our next objective is to show that each projection P_v arising from an intertwiner lies below $Q_1 \otimes Q_2$. For the next two lemmas, let $v \in \mathcal{GN}_M^{(1)}(B)$ be fixed. We continue to employ the notation P_v for the projection $\sum_n v_n^* e_B v_n \in Z(B' \cap \langle M, e_B \rangle)$ which satisfies $P_v v^* v = v^* e_B v$.

Let A_1 be the abelian von Neumann algebra $Z(B'_1 \cap \langle M_1, e_{B_1} \rangle)S_1$ on which Φ , the restriction of Tr_1 , is semifinite.

Lemma 4.5. *If r is a projection in $A_1 \overline{\otimes} Z(B_2)$, then*

$$(\Phi \otimes \text{Tr}_2)(P_v r) \leq (\Phi \otimes \tau_2)(r). \quad (4.22)$$

Proof. There is a measure space (Ω, Σ, μ) so that A_1 corresponds to $L^\infty(\Omega)$ while Φ is given by integration with respect to the σ -finite measure μ . Then P_v is viewed as a projection-valued function $P_v(\omega)$, with the same representation $r(\omega)$ for r . For $i = 1, 2$, let Ψ_i be the pull down map for $\langle M_i, e_{B_i} \rangle$. Then $(\Psi_1 \otimes \Psi_2)(P_v) = \sum_n v_n^* v_n$ which is a projection, so has norm 1. By Lemma 2.5, $\|(I \otimes \Psi_2)(P_v)\| \leq 1$, and so this element of $A_1 \overline{\otimes} M_2$ can be represented as a function $f(\omega)$ with $\|f(\omega)\| \leq 1$ almost everywhere. It follows that

$$\begin{aligned} (\Phi \otimes \tau_2)(r) &= \int_{\Omega} \tau_2(r(\omega)) d\mu(\omega) \geq \int_{\Omega} \tau_2(r(\omega)f(\omega)) d\mu(\omega) \\ &= (\Phi \otimes \tau_2)((I \otimes \Psi_2)(P_v)r) = (\Phi \otimes \tau_2)((I \otimes \Psi_2)(P_v r)) \\ &= (\Phi \otimes \text{Tr}_2)(P_v r), \end{aligned} \quad (4.23)$$

where the penultimate equality is valid because $r \in A_1 \overline{\otimes} Z(B_2)$ and the M_2 -bimodular property of Ψ_2 applies. \square

Lemma 4.6. *For $v \in \mathcal{GN}^{(1)}(B)$, the associated projection $P_v \in Z(B' \cap \langle M, e_B \rangle)$ satisfies $P_v \leq Q_1 \otimes Q_2$.*

Proof. The remarks preceding Definition 4.4 show that $P_v \leq S_1 \otimes S_2$. We will prove the stronger inequality $P_v \leq S_1 \otimes Q_2$, which is sufficient to establish the result since we will also have $P_v \leq Q_1 \otimes S_2$ by a symmetric argument.

As before, let A_1 denote $Z(B'_1 \cap \langle M_1, e_{B_1} \rangle)S_1$ and let Φ be the restriction of Tr_1 to A_1 . Then, as noted earlier, A_1 is an abelian von Neumann algebra and Φ is a faithful normal semifinite weight on A_1 . Let $\{q_n\}_{n=1}^\infty$ be a maximal family of nonzero orthogonal projections in $Z(B_1) \overline{\otimes} B_2$ so that $P_v q_n = w_n^*(1 \otimes e_{B_2})w_n$ for partial isometries $w_n \in A_1 \overline{\otimes} M_2$ which are intertwiners of $A_1 \overline{\otimes} B_2$. Let $q = \sum_{n=1}^\infty q_n$, defining q to be 0 if no such projections exist. We will first show that $P_v \leq q$, so suppose that $(1 - q)P_v \neq 0$.

The central support of $1 \otimes e_{B_2}$ in $A_1 \overline{\otimes} \langle M_2, e_{B_2} \rangle$ is 1, so there is a nonzero subprojection Q of $(1 - q)P_v$ in $A_1 \overline{\otimes} \langle M_2, e_{B_2} \rangle$ with $Q \lesssim 1 \otimes e_{B_2}$ in this algebra. The projection P_v has the form $P_v = \sum_{n \geq 0} v_n^* e_B v_n$ where $\sum_{n \geq 0} v_n^* v_n$ is the central support $z \in Z(B)$ of $v^* v \in B$. With this notation, (3.20) becomes

$$N(P_v) = \{x \in zMz: xP_v = P_v x\}.$$

By Lemma 3.8 (iii), there is a projection $f \in N(P_v)$ so that $Q = fP_v$. Both Q and P_v commute with $B_1 \otimes 1$, so the relation $(b_1 \otimes 1)Q - Q(b_1 \otimes 1) = 0$ for $b_1 \in B_1$ becomes $((b_1 \otimes 1)f - f(b_1 \otimes 1))P_v = 0$. The element $(b_1 \otimes 1)f - f(b_1 \otimes 1)$ lies in $N(P_v)$ so, by Lemma 3.8 (i), $(b_1 \otimes 1)f = f(b_1 \otimes 1)$ for all $b_1 \in B_1$. This shows that $f \in B'_1 \cap N(P_v)$. Moreover, $f = (1 - q)f$ follows from the equation

$$0 = qQ = qfP_v, \quad (4.24)$$

which implies that $qf = 0$ since $qf \in N(P_v)$. Note that $(1-q)z \neq 0$, otherwise $zf = 0$. Now

$$B'_1 \cap N(P_v) \subseteq B'_1 \cap M = Z(B_1) \overline{\otimes} M_2 \quad (4.25)$$

and

$$\left(Z(B_1) \overline{\otimes} B_2 \right)' \cap \left(Z(B_1) \overline{\otimes} M_2 \right) = Z(B_1) \overline{\otimes} Z(B_2) \subseteq Z(B_1) \overline{\otimes} B_2. \quad (4.26)$$

Thus the inclusion

$$(1-q)z(Z(B_1) \overline{\otimes} B_2)z(1-q) \subseteq (1-q)z(B'_1 \cap N(P_v))z(1-q) \quad (4.27)$$

has the property that the first algebra contains its relative commutant in the second algebra, which is the hypothesis of Lemma 2.6. Thus we may choose a nonzero projection $b \in (1-q)z(Z(B_1) \overline{\otimes} B_2)z(1-q)$ with $b \preceq f$ in $(1-q)z(B'_1 \cap N(P_v))z(1-q)$. Let w be a partial isometry in this algebra with $w^*w = b$ and $ww^* \leq f$, and note that w commutes with P_v by definition of $N(P_v)$. Then

$$bP_v = w^*wP_v = P_vw^*wP_v \sim wP_vw^* = ww^*P_v \leq fP_v \quad (4.28)$$

in $A_1 \overline{\otimes} \langle M_2, e_{B_2} \rangle$. Since $b \leq z$, Lemma 3.8 (i) ensures that $bP_v \neq 0$. Moreover, $bP_v \preceq 1 \otimes e_{B_2}$ in $A_1 \overline{\otimes} \langle M_2, e_{B_2} \rangle$ since fP_v has this property.

Consider now a projection $r \in (A_1 \overline{\otimes} Z(B_2))b$. The inequality

$$\Phi \otimes \text{Tr}_2(P_v r) \leq \Phi \otimes \tau(r) \quad (4.29)$$

is valid by Lemma 4.5. Thus the hypotheses of Theorem 4.1 are satisfied with P replaced by bP_v . We conclude that there is an element $w \in \mathcal{GN}_{A_1 \overline{\otimes} M_2}^{(1)}(A_1 \overline{\otimes} B_2)$ so that $bP_v = w^*(1 \otimes e_{B_2})w$. Since b lies under $1-q$, this contradicts maximality of the q_i 's, proving that $P_v q = P_v$. Thus

$$P_v = \sum_{n=1}^{\infty} w_n^*(1 \otimes e_{B_2})w_n, \quad (4.30)$$

which we also write as $P_v = \sum_{n=1}^{\infty} W_n^*W_n$ where W_n is defined to be $(1 \otimes e_{B_2})w_n \in A_1 \overline{\otimes} \langle M_2, e_{B_2} \rangle$.

As in Lemma 4.5, we regard (A_1, Tr_1) as $L^\infty(\Omega)$ for a σ -finite measure space (Ω, μ) . We can then identify $A_1 \overline{\otimes} \langle M_2, e_{B_2} \rangle$ with $L^\infty(\Omega, \langle M_2, e_{B_2} \rangle)$, and we write elements of this tensor product as uniformly bounded measurable functions on Ω with values in $\langle M_2, e_{B_2} \rangle$. Then

$$\text{Tr}(P_v) = \int \sum_{n=1}^{\infty} \text{Tr}_2(W_n(\omega)^*W_n(\omega)) d\mu(\omega). \quad (4.31)$$

Since $\text{Tr}(P_v) < \infty$, we may neglect a countable number of null sets to conclude that

$$\text{Tr}_2(P_v(\omega)) = \text{Tr}_2 \left(\sum_{n=1}^{\infty} W_n(\omega)^*W_n(\omega) \right) < \infty, \quad \omega \in \Omega, \quad (4.32)$$

from which it follows that $P_v(\omega) \leq Q_2$ for $\omega \in \Omega$. Thus $P_v \leq 1 \otimes Q_2$ which gives $P_v \leq S_1 \otimes Q_2$, since the inequality $P_v \leq S_1 \otimes S_2$ has already been established. \square

We are now in a position to approximate an intertwiner in a tensor product.

Theorem 4.7. *Let $B_i \subseteq M_i$, $i = 1, 2$, be inclusions of finite von Neumann algebras satisfying $B'_i \cap M_i \subseteq B_i$. Given $v \in \mathcal{GN}_{M_1 \overline{\otimes} M_2}^{(1)}(B_1 \overline{\otimes} B_2)$ and $\varepsilon > 0$, there exist $x_1, \dots, x_k \in B_1 \overline{\otimes} B_2$, $w_{1,1}, \dots, w_{1,k} \in \mathcal{GN}_{M_1}^{(1)}(B_1)$ and $w_{2,1}, \dots, w_{2,k} \in \mathcal{GN}_{M_2}^{(1)}(B_2)$ such that:*

1. $\|x_j\| \leq 1$ for each j ;

2.

$$\left\| v - \sum_{j=1}^k x_j (w_{1,j} \otimes w_{2,j}) \right\|_{2,\tau} < \varepsilon. \quad (4.33)$$

Proof. Write $B \subseteq M$ for the inclusion $B_1 \overline{\otimes} B_2 \subseteq M_1 \overline{\otimes} M_2$ and fix $v \in \mathcal{GN}_M^{(1)}(B)$. Recall from Lemmas 3.6 and 4.6 that there is a projection $P_v \in Z(B' \cap \langle M, e_B \rangle)$ satisfying $P_v v^* v = v^* e_B v$, and $P_v \leq Q_1 \otimes Q_2$ where Q_i is the supremum of the set \mathcal{P}_i of projections in $Z(B'_i \cap \langle M_i, e_{B_i} \rangle)$ specified in Definition 4.4. Thus $Q_i = \sum_k R_{i,k}$ for some countable sum of orthogonal projections $R_{i,k} \in \mathcal{P}_i$. By Lemma 3.8 (iii), any subprojection of $R_{i,k}$ in $Z(B'_i \cap \langle M_i, e_{B_i} \rangle)$ is also in \mathcal{P}_i , so it follows that every subprojection of Q_i in $B' \cap \langle M_i, e_{B_i} \rangle$ is also of the form $\sum_k R'_{i,k}$ for some countable sum of orthogonal projections $R'_{i,k} \in \mathcal{P}_i$.

The restriction Φ_i of Tr_i to $Z(B'_i \cap \langle M_i, e_{B_i} \rangle) Q_i$ is a normal semifinite weight on this abelian von Neumann algebra A_i , and (A_i, Tr_i) can be identified with $L^\infty(\Omega_i)$ for a σ -finite measure space $(\Omega_i, \Sigma_i, \mu_i)$. Since $P_v \leq Q_1 \otimes Q_2$, this operator can be viewed as an element of $L^\infty(\Omega_1 \times \Omega_2, \mu_1 \times \mu_2)$, and it also lies in the corresponding L^2 -space since $\text{Tr}(P_v) \leq 1$. By Lemma 2.7 and the previous paragraph, P_v can be approximated in $\|\cdot\|_{2,\text{Tr}}$ -norm by finite sums of orthogonal projections of the form $R_1 \otimes R_2$, each lying in \mathcal{P}_i . These elementary tensors correspond to measurable rectangles in $\Omega_1 \times \Omega_2$. By the definition of \mathcal{P}_i , each R_i is close in $\|\cdot\|_{2,\text{Tr}_i}$ -norm to a finite sum $\sum_{j=1}^k w_{i,j}^* e_{B_i} w_{i,j}$, with $w_{i,j} \in \mathcal{GN}_{M_i}^{(1)}(B_i)$. This allows us to make the following approximation: given $\varepsilon > 0$, there exist finite sets $\{w_{i,j}\}_{j=1}^k \in \mathcal{GN}_{M_i}^{(1)}(B_i)$, $i = 1, 2$, such that

$$\left\| P_v - \sum_{j=1}^k (w_{1,j}^* e_{B_1} w_{1,j}) \otimes (w_{2,j}^* e_{B_2} w_{2,j}) \right\|_{2,\text{Tr}} < \varepsilon. \quad (4.34)$$

If we multiply on the right in (4.34) by $v^* e_B = v^* e_B v v^* e_B$, then the result is

$$\left\| v^* e_B - \left(\sum_{j=1}^k (w_{1,j}^* e_{B_1} w_{1,j}) \otimes (w_{2,j}^* e_{B_2} w_{2,j}) \right) (v^* e_B) \right\|_{2,\text{Tr}} < \varepsilon, \quad (4.35)$$

using the fact that $P_v v^* v = v^* e_B v$. A typical element of the sum in (4.35) is

$$(w_{1,j}^* \otimes w_{2,j}^*) (e_{B_1} \otimes e_{B_2}) (w_{1,j} \otimes w_{2,j}) v^* (e_{B_1} \otimes e_{B_2})$$

which has the form $(w_{1,j}^* \otimes w_{2,j}^*)x_j^*e_B$, where $x_j^* = \mathbb{E}_B((w_{1,j} \otimes w_{2,j})v^*) \in B$ has $\|x_j^*\| \leq \|(w_{1,j} \otimes w_{2,j})v^*\| \leq 1$. Thus (4.35) becomes

$$\left\| \left(v^* - \sum_{j=1}^k (w_{1,j} \otimes w_{2,j})^* x_j^* \right) e_B \right\|_{2, \text{Tr}} < \varepsilon. \quad (4.36)$$

For each $y \in M$,

$$\|ye_B\|_{2, \text{Tr}}^2 = \text{Tr}(e_B y^* y e_B) = \tau(y^* y) = \|y\|_{2, \tau}^2, \quad (4.37)$$

and so (4.36) implies that

$$\left\| v - \sum_{j=1}^k x_j (w_{1,j} \otimes w_{2,j}) \right\|_{2, \tau} < \varepsilon,$$

as required. \square

Corollary 4.8. *Let $B_i \subseteq M_i$, $i = 1, 2$, be inclusions of finite von Neumann algebras satisfying $B'_i \cap M_i \subseteq B_i$. Then*

$$W^*(\mathcal{GN}^{(1)}(B_1 \overline{\otimes} B_2)) = W^*(\mathcal{GN}^{(1)}(B_1)) \overline{\otimes} W^*(\mathcal{GN}^{(1)}(B_2)). \quad (4.38)$$

There are two extreme cases where the hypothesis $B' \cap M \subseteq B$ is satisfied. The first is when B is an irreducible subfactor where the result of Theorem 4.7 can be deduced from the stronger results of [20]. The second is when B is a masa in M where Theorem 4.7 is already known, [1]. The following example explains the preference given to intertwiners over unitary normalizers in intermediate cases, even in a simple setting.

Example 4.9. Let M be a II_1 factor, let $p \in M$ be projection whose trace lies in $(0, 1/2)$, and let $B = pMp + (1-p)M(1-p)$. This subalgebra has no nontrivial unitary normalizers, essentially because $\tau(p) \neq \tau(1-p)$. However, the tensor product $B \overline{\otimes} B \subseteq M \overline{\otimes} M$ does have such normalizers because the compressions by $p \otimes (1-p)$ and $(1-p) \otimes p$, which have equal traces, are conjugate by a unitary normalizer u which is certainly outside $B \overline{\otimes} B$. According to Theorem 4.7, u can be obtained as the limit of finite sums of elementary tensors from $W^*(\mathcal{GN}^{(1)}(B)) \overline{\otimes} W^*(\mathcal{GN}^{(1)}(B))$. \square

5 Groupoid normalizers of tensor products

In this section we return to the groupoid normalizers $\mathcal{GN}_M(B)$, namely those $v \in M$ such that $v, v^* \in \mathcal{GN}_M^{(1)}(B)$. Our objective in this section is to establish a corresponding version of Theorem 4.7 for $\mathcal{GN}(B)$, and consequently we will assume throughout that any inclusion $B \subseteq M$ satisfies the relative commutant condition $B' \cap M \subseteq B$.

We will need to draw a sharp distinction between those intertwiners v that are groupoid normalizers and those that are not, and so we introduce the following definition.

Definition 5.1. Say that $v \in \mathcal{GN}^{(1)}(B)$ is *strictly one-sided* if the only projection $p \in Z(Bv^*v) = Z(B)v^*v$ for which $vp \in \mathcal{GN}(B)$ is $p = 0$. When B is an irreducible subfactor of M then any unitary $u \in M$ satisfying $uBu^* \not\subseteq B$ is a strictly one-sided intertwiner (see [20, Example 5.4] for examples of such unitaries).

Given $v \in \mathcal{GN}(B)$, recall from Section 3 that there is a projection $P_v \in Z(B' \cap \langle M, e_B \rangle)$ such that $P_v v^* v = v^* e_B v$, and P_v has the form

$$P_v = \sum_{n \geq 0} v_n^* e_B v_n, \quad (5.1)$$

where there exist partial isometries $w_n \in B$ so that $v_n = v w_n^* \in \mathcal{GN}^{(1)}(B)$, and $w_m^* w_n = 0$ for $m \neq n$. Letting p_n denote the projection $w_n w_n^* \in B$, it also holds that $v_n^* v_n = p_n$. We will employ this notation below.

Lemma 5.2. *Let $v \in \mathcal{GN}(B)$, and let $u \in \mathcal{GN}^{(1)}(B)$ be strictly one-sided. Then $P_v u^* e_B u = 0$.*

Proof. By Lemmas 3.2 (i) and 3.6, $u^* e_B u \in Z(B' \cap \langle M, e_B \rangle) u^* u$ and $P_v \in Z(B' \cap \langle M, e_B \rangle)$, showing that P_v and $u^* e_B u$ are commuting projections. Let Q denote the projection $P_v u^* e_B u$ in $Z(B' \cap \langle M, e_B \rangle) u^* u$ which lies below both P_v and $u^* e_B u$. From Lemmas 3.4 and 3.8 we may find projections $z \in Z(B)$ and $f \in M \cap \{P_v\}'$ such that

$$Q = z u^* e_B u = f P_v. \quad (5.2)$$

From (5.1), write P_v as the strongly convergent sum

$$P_v = \sum_{n \geq 0} w_n v^* e_B v w_n^*, \quad (5.3)$$

so that (5.2) becomes

$$\sum_{n \geq 0} f w_n v^* e_B v w_n^* = z u^* e_B u. \quad (5.4)$$

If we multiply (5.4) on the right by $p_j = w_j w_j^*$ and on the left by $e_B u$, noting that $u z u^* \in B$, then the result is

$$e_B b_j v w_j^* = e_B u z u^* a p_j = e_B u u^* u z p_j = e_B u z p_j \quad (5.5)$$

for each $j \geq 0$, where $b_j = \mathbb{E}_B(u f w_j v^*) \in B$. Thus

$$b_j v w_j^* = u z p_j, \quad j \geq 0. \quad (5.6)$$

If we sum (5.6) over $j \geq 0$, then the right-hand side will converge strongly, implying strong convergence of $\sum_{j \geq 0} b_j v w_j^*$. If we return to (5.4) and multiply on the left by $e_B u$, then we obtain

$$u z = \sum_{n \geq 0} b_n v w_n^* \quad (5.7)$$

with strong convergence of this sum. It follows that, for each $b \in B$,

$$z u^* b u z = \lim_{k \rightarrow \infty} \sum_{m, n \leq k} w_n v^* b_n^* b b_m v w_m^* \quad (5.8)$$

strongly, and thus $z u^* b u z \in B$ since $v^* b_n^* b b_m v \in B$. Thus the projection $p = z u^* u \in Z(B u^* u)$ satisfies $p u^* B u p \subseteq B$. Since u is strictly one-sided, we conclude that $z u^* u = 0$, showing that

$$Q = z u^* e_B u = 0 \quad (5.9)$$

from (5.2). This proves the result. \square

We can use the preceding lemma to show that a strictly one-sided intertwiner $v \in \mathcal{GN}^{(1)}(B)$ has the property that the only projection $p \in Bv^*v$ for which $vp \in \mathcal{GN}(B)$ is $p = 0$. Indeed, take such a projection p for which $w = vp \in \mathcal{GN}(B)$. Then $P_v w^* w = w^* e_B w$ so that $P_v \geq P_w$ by Remark 3.7. Lemma 5.2 then gives $P_w v^* e_B v = 0$. Thus $P_w w^* e_B w = w^* e_B w = 0$ and so $w = 0$.

Lemma 5.3. *Let $v \in \mathcal{GN}^{(1)}(B)$. Then there exist orthogonal families of orthogonal projections $e_n, f_n \in v^*vBv^*v$ such that $\sum(e_n + f_n) = v^*v$, each ve_n is strictly one-sided, and each vf_n lies in $\mathcal{GN}(B)$.*

Proof. Let $\{e_n\}$ be a maximal orthogonal family of projections in $Z(B)v^*v$ such that ve_n is strictly one-sided, and set $e = \sum e_n$. Then choose a maximal family of orthogonal projections $\{f_n\} \in Z(B)(v^*v - e)$ such that $vf_n \in \mathcal{GN}(B)$. If $\sum e_n + \sum f_n = v^*v$ then the result is proved, so consider the projection $g = v^*v - \sum e_n - \sum f_n \in Z(B)v^*v$ and suppose that $g \neq 0$. Then vg cannot be strictly one-sided otherwise the maximality of $\{e_n\}$ would be contradicted. Thus there exists $z \in Z(B)$ such that vgz is a nonzero element of $\mathcal{GN}(B)$. But this contradicts maximality of $\{f_n\}$, proving the result. \square

We now return to considering two containments $B_i \subseteq M_i$ satisfying $B'_i \cap M_i \subseteq B_i$, and the tensor product containment $B = B_1 \overline{\otimes} B_2 \subseteq M_1 \overline{\otimes} M_2 = M$. The next lemma is the key step required to obtain a version of Theorem 4.7 for groupoid normalizers. We need a result from the perturbation theory of finite von Neumann algebras. For any containment $A \subseteq N$, where N has a specified trace τ , recall that $N \subset_{\delta, \tau} A$ means that

$$\sup\{\|x - \mathbb{E}_A(x)\|_2 : x \in N, \|x\| \leq 1\} \leq \delta$$

where $\|\cdot\|_2$ is defined using the given trace τ . If τ is scaled by a constant λ , then $\subset_{\delta, \lambda\tau}$ is the same as $\subset_{\delta/\sqrt{\lambda}, \tau}$. Then [17, Theorem 3.5] (see also [18, Theorem 10.3.5]), stated for normalized traces, has the following general interpretation: if $A \subseteq N$ and $N \subset_{\delta, \tau} A$ for some $\delta < (\tau(1)/23)^{1/2}$, then there exists a nonzero projection $p \in Z(A' \cap N)$ such that $Ap = pNp$.

Lemma 5.4. *Let $v \in \mathcal{GN}^{(1)}(B_1)$ and $w \in \mathcal{GN}^{(1)}(B_2)$. If v (or w) is strictly one-sided then $v \otimes w \in \mathcal{GN}^{(1)}(B)$ is strictly one-sided.*

Proof. Without loss of generality, suppose that v is strictly one-sided. Fix a nonzero projection $p \in Z(B)(v^*v \otimes w^*w)$, and let τ_i be the faithful normalized normal trace on $M_i \supset B_i$. Given $\varepsilon > 0$, we may choose projections $p_i \in Z(B_1)v^*v$ and $q_i \in Z(B_2)w^*w$, $1 \leq i \leq k$, with $\|p - \sum_{i=1}^k p_i \otimes q_i\|_{2, \tau} < \varepsilon$ and the p_i 's orthogonal since these projections lie in abelian von Neumann algebras. Here, $\|\cdot\|_{2, \tau}$ is with respect to the normalized trace $\tau = \tau_1 \otimes \tau_2$ on M . Now

$$p_i B_1 p_i = p_i v^* v B_1 v^* v p_i \subseteq p_i v^* B_1 v p_i \tag{5.10}$$

since $v B_1 v^* \subseteq B_1$. If it were true that $p_i v^* B_1 v p_i \subset_{\delta, \tau_1} p_i B_1 p_i$ for some $\delta < (\tau(p_i)/23)^{1/2}$, then it would follow from [17, Theorem 3.5] that there exists a nonzero projection $p'_i \in (p_i B_1 p_i)' \cap p_i v^* B_1 v p_i \subseteq Z(B_1)p_i$ such that $p'_i p_i B_1 p_i = p'_i p_i v^* B_1 v p_i p'_i$, contradicting the hypothesis that v is strictly one-sided. Thus there exists $b_i \in B_1$ satisfying

$$\|p_i v^* b_i v p_i\| \leq 1, \quad d_{2, \tau_1}(p_i v^* b_i v p_i, p_i B_1 p_i) \geq (\tau_1(p_i)/23)^{1/2}, \tag{5.11}$$

where $d_{2,\tau_1}(x, A) = \inf\{\|x - a\|_{2,\tau_1} : a \in A\}$ for any von Neumann algebra A .

Since each p_i lies under v^*v , the projections $vp_i v^*$ lie in B_1 and are orthogonal. We may then define an element $b \in B$ by

$$b = \sum_i vp_i v^* b_i vp_i v^* \otimes w q_i w^*, \quad (5.12)$$

and the orthogonality gives $\|b\| \leq 1$. Moreover,

$$(v \otimes w)^* b (v \otimes w) = \sum_i p_i v^* b_i v p_i \otimes q_i, \quad (5.13)$$

and

$$\begin{aligned} d_{2,\tau}((v \otimes w)^* b (v \otimes w), B)^2 &\geq \left(\sum_i \tau_1(p_i) \tau_2(q_i) \right) / 23 \\ &> (\tau(p) - \varepsilon) / 23. \end{aligned} \quad (5.14)$$

The right-hand side of (5.13) is unchanged by pre- and post-multiplication by the projection $\sum_i p_i \otimes q_i$, and this is close to p . This leads to the estimate

$$\|p(v \otimes w)^* b (v \otimes w) p - (v \otimes w)^* b (v \otimes w)\|_{2,\tau} \leq 2 \left\| p - \sum_i p_i \otimes q_i \right\|_{2,\tau} < 2\varepsilon. \quad (5.15)$$

From (5.14) and (5.15),

$$d_{2,\tau}(p(v \otimes w)^* b (v \otimes w) p, B) > ((\tau(p) - \varepsilon) / 23)^{1/2} - 2\varepsilon. \quad (5.16)$$

A sufficiently small choice of ε then shows that $p(v \otimes w)^* b (v \otimes w) p \notin B$, and thus $v \otimes w$ is strictly one-sided. \square

We can now give the two-sided counterpart of Theorem 4.7.

Theorem 5.5. *Let $B_i \subseteq M_i$, $i = 1, 2$, be inclusions of finite von Neumann algebras satisfying $B'_i \cap M_i \subseteq B_i$. Given $v \in \mathcal{GN}_{M_1 \overline{\otimes} M_2}(B_1 \overline{\otimes} B_2)$ and $\varepsilon > 0$, there exist $x_1, \dots, x_k \in B_1 \overline{\otimes} B_2$, $w_{1,1}, \dots, w_{1,k} \in \mathcal{GN}_{M_1}(B_1)$ and $w_{2,1}, \dots, w_{2,k} \in \mathcal{GN}_{M_2}(B_2)$ such that:*

1. $\|x_j\| \leq 1$ for each j ;

2.

$$\left\| v - \sum_{j=1}^k x_j (w_{1,j} \otimes w_{2,j}) \right\|_{2,\tau} < \varepsilon. \quad (5.17)$$

Proof. Consider $v \in \mathcal{GN}(B)$. Following the proof of Theorem 4.7, given $\varepsilon > 0$, there exist $v_{i,j} \in \mathcal{GN}^{(1)}(B_i)$, $1 \leq j \leq k$, so that

$$\left\| P_v - \sum_{j=1}^k (v_{1,j} \otimes v_{2,j})^* e_B(v_{1,j} \otimes v_{2,j}) \right\|_{2,\text{Tr}} < \varepsilon \quad (5.18)$$

as in (4.34). Using Lemma 5.3, we may replace this sum with one of the form

$$\sum (w_{1,j} \otimes w_{2,j})^* e_B(w_{1,j} \otimes w_{2,j}) + \sum (x_{1,j} \otimes x_{2,j})^* e_B(x_{1,j} \otimes x_{2,j})$$

where the $w_{i,j}$'s are two-sided and, for each j , at least one of $x_{1,j}, x_{2,j}$ is strictly one-sided. By Lemma 5.4, each $x_{1,j} \otimes x_{2,j}$ is strictly one-sided, so $(x_{1,j} \otimes x_{2,j})^* e_B(x_{1,j} \otimes x_{2,j}) P_v = 0$ by Lemma 5.2. If we multiply on the right by P_v , then

$$\left\| P_v - \sum_j (w_{1,j} \otimes w_{2,j})^* e_B(w_{1,j} \otimes w_{2,j}) P_v \right\|_{2,\tau} < \varepsilon. \quad (5.19)$$

Simple approximation allows us to obtain the same estimate for some finite subcollection of the $w_{1,j}$ and $w_{2,j}$, say $w_{1,1}, \dots, w_{1,k}$ and $w_{2,1}, \dots, w_{2,k}$. We now continue to follow the proof of Theorem 4.7 from (4.34) to obtain the required x_j . \square

Just as in Section 4, the next corollary follows immediately.

Corollary 5.6. *Let $B_i \subseteq M_i$, $i = 1, 2$, be inclusions of finite von Neumann algebras satisfying $B'_i \cap M_i \subseteq B_i$. Then*

$$\mathcal{GN}_{M_1}(B_1)'' \overline{\otimes} \mathcal{GN}_{M_2}(B_2)'' = \mathcal{GN}_{M_1 \overline{\otimes} M_2}(B_1 \overline{\otimes} B_2)''.$$

References

- [1] I. Chifan. On the normalizing algebra of a masa in a II_1 factor. arXiv:math.OA/0606225, 2006.
- [2] E. Christensen. Subalgebras of a finite algebra. *Math. Ann.*, 243:17–29, 1979.
- [3] J. Dixmier. Sous-anneaux abéliens maximaux dans les facteurs de type fini. *Ann. of Math. (2)*, 59:279–286, 1954.
- [4] H. A. Dye. On groups of measure preserving transformation. I. *Amer. J. Math.*, 81:119–159, 1959.
- [5] E. G. Effros and Z.-J. Ruan. On approximation properties for operator spaces. *Internat. J. Math.*, 1(2):163–187, 1990.
- [6] J. Fang. On completely singular von Neumann subalgebras. *Proc. Edinb. Math. Soc. (2)*, to appear. arXiv:math.OA/0606649.
- [7] J. Feldman and C. C. Moore. Ergodic equivalence relations, cohomology, and von Neumann algebras. I. *Trans. Amer. Math. Soc.*, 234(2):289–324, 1977.
- [8] J. Feldman and C. C. Moore. Ergodic equivalence relations, cohomology, and von Neumann algebras. II. *Trans. Amer. Math. Soc.*, 234(2):325–359, 1977.

- [9] V. Jones and V. S. Sunder. *Introduction to subfactors*, volume 234 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1997.
- [10] V. F. R. Jones. Index for subfactors. *Invent. Math.*, 72(1):1–25, 1983.
- [11] R. V. Kadison and J. R. Ringrose. *Fundamentals of the theory of operator algebras. Vol. I*, volume 15 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1997. Elementary theory, Reprint of the 1983 original.
- [12] R. V. Kadison and J. R. Ringrose. *Fundamentals of the theory of operator algebras. Vol. II*, volume 16 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1997. Advanced theory, Corrected reprint of the 1986 original.
- [13] M. Pimsner and S. Popa. Entropy and index for subfactors. *Ann. Sci. École Norm. Sup. (4)*, 19(1):57–106, 1986.
- [14] S. Popa. On a class of type II_1 factors with Betti numbers invariants. *Ann. of Math. (2)*, 163(3):809–899, 2006.
- [15] S. Popa. Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups, I. *Invent. Math.*, 165:369–408, 2006.
- [16] H. L. Royden. *Real analysis*, third edition. Macmillan Publishing Company, New York, 1988.
- [17] S. Popa, A. M. Sinclair, and R. R. Smith. Perturbations of subalgebras of type II_1 factors. *J. Funct. Anal.*, 213(2):346–379, 2004.
- [18] A. M. Sinclair and R. R. Smith. *Finite von Neumann algebras and masas*, volume 351 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2008.
- [19] A. M. Sinclair, R. R. Smith, S. A. White, and A. Wiggins. Strong singularity of singular masas in II_1 factors. *Illinois J. Math.*, 51(4):1077–1083, 2007.
- [20] R. R. Smith, S. A. White, and A. Wiggins. Normalizers of irreducible subfactors. arXiv:0705.0183, 2007.
- [21] M. Takesaki. *Theory of operator algebras. II*, volume 125 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2003. Operator Algebras and Non-commutative Geometry, 6.

Author Addresses

Junsheng Fang
Department of Mathematics
University of New Hampshire
Durham, NH 03824
USA
jfang@cisunix.unh.edu

Stuart White
Department of Mathematics
University of Glasgow
Glasgow G12 8QW
UK
s.white@maths.gla.ac.uk

Roger Smith
Department of Mathematics
Texas A&M University
College Station, TX 77843
USA
rsmith@math.tamu.edu

Alan Wiggins
Department of Mathematics
Vanderbilt University
Nashville, TN 37240
USA
alan.d.wiggins@vanderbilt.edu