

Section 1.1: Vectors

Quantities that we measure that have magnitude but not direction are called **scalars**.

DEFINITION 1. A **vector** is a quantity that has both magnitude and direction. A 2-dimensional vector is an ordered pair $\mathbf{a} = \langle a_1, a_2 \rangle$. The numbers a_1 and a_2 are called the **components** of the vector \mathbf{a} .

Typical notation to designate a vector is a boldfaced character or a character with an arrow on it (i.e. \mathbf{a} or \vec{a}).

DEFINITION 2. Given the points $A(a_1, a_2)$ and $B(b_1, b_2)$, the vector \mathbf{a} with representation \vec{AB} is

$$\vec{AB} = \langle b_1 - a_1, b_2 - a_2 \rangle.$$

The point A here is initial point and B is terminal one.

A vector with the initial point located at the origin is called the **position** vector (or we say that a vector is in standard position).

Vectors are equal if they have the same length and direction (same slope).

EXAMPLE 3. Graph the vector with initial point $A(1, -2)$ and terminal point $B(2, 1)$. Find the components of \vec{AB} and \vec{BA} .

Vector operations

- *Scalar Multiplication:* If c is a scalar and $\mathbf{a} = \langle a_1, a_2 \rangle$, then

$$c\mathbf{a} = c \langle a_1, a_2 \rangle = \langle ca_1, ca_2 \rangle.$$

DEFINITION 4. Two vectors \mathbf{a} and \mathbf{b} are called **parallel** if $\mathbf{b} = c\mathbf{a}$ with some scalar c .

If $c > 0$ then a and ca have the same direction, if $c < 0$ then a and ca have the opposite direction.

- *Vector addition:* If $\mathbf{a} = \langle a_1, a_2 \rangle$ and $\mathbf{b} = \langle b_1, b_2 \rangle$ then

$$\mathbf{a} + \mathbf{b} = \langle a_1 + b_1, a_2 + b_2 \rangle.$$

Triangle Law

Parallelogram Law

$\mathbf{a} + \mathbf{b}$ is called the **resultant vector**

EXAMPLE 5. Let $\mathbf{a} = \langle -1, 2 \rangle$ and $\mathbf{b} = \langle 2.1, -0.5 \rangle$. Then $3\mathbf{a} + 2\mathbf{b} =$

The **magnitude** or **length** of a vector $\mathbf{a} = \langle a_1, a_2 \rangle$ is denoted by $|\mathbf{a}|$,

$$|\mathbf{a}| =$$

EXAMPLE 6. Find: $|\langle 3, -8 \rangle|$, $\left| \left\langle \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right\rangle \right|$, $|\mathbf{0}|$

A **unit** vector is a vector with length one. Any vector can be made into a unit vector by dividing it by its length. So, a unit vector in the direction of \mathbf{a} is

$$\mathbf{u} = \frac{\mathbf{a}}{|\mathbf{a}|}.$$

Any vector \mathbf{a} can be fully represented by providing its length, $|\mathbf{a}|$ and a unit vector \mathbf{u} in its direction:

$$\mathbf{a} = |\mathbf{a}| \mathbf{u}$$

EXAMPLE 7. Given $\mathbf{a} = \langle 2, -1 \rangle$. Find

(a) a unit vector that has the same direction as \mathbf{a} ;

(b) a vector \mathbf{b} in the direction opposite to \mathbf{a} s.t $|\mathbf{b}| = 7$.

The **standard basis vectors** are given by the unit vectors $\mathbf{i} = \langle 1, 0 \rangle$ and $\mathbf{j} = \langle 0, 1 \rangle$ along the x and y directions, respectively. Using the basis vectors, one can represent any vector $\mathbf{a} = \langle a_1, a_2 \rangle$ as

$$\mathbf{a} = a_1 \mathbf{i} + a_2 \mathbf{j}.$$


EXAMPLE 8. Given $\mathbf{a} = 2\mathbf{i} - \mathbf{j}$, $\mathbf{b} = \langle 5, -2 \rangle$. Find a scalars s and t such that $s\mathbf{a} + t\mathbf{b} = -4\mathbf{j}$.

Applications: Quantities such as force, displacement or velocity that have direction as well as magnitude are represented by vectors.

EXAMPLE 9. *Ben walks due west on the deck of a ship at 5 mph. The ship is moving north at a speed of 25 mph. Find the direction and speed of Ben relative to the surface of the water.*

EXAMPLE 10. *Two forces F_1 and F_2 with magnitudes 14 pounds and 12 pounds act on an object at a point P as shown. Find the resultant force as well as its magnitude and direction.*

EXAMPLE 11. An 60 pound weight hangs from two wires as shown. Find the tensions (forces) in both wires and their magnitudes.



EXAMPLE 12. *An airplane, flying due east at an airspeed of 450mph, encounters a 50-mph wind acting in the direction of $E60^\circ N$ (60° North of East). The airplane holds its compass heading due east but, because of the wind, acquires a new ground speed (i.e. the magnitude of the resultant) and direction. What are they?*