

## Section 2.3: Calculating limits using the limits laws

LIMIT LAWS Suppose that  $c$  is a constant and the limits

$$\lim_{x \rightarrow a} f(x) \quad \text{and} \quad \lim_{x \rightarrow a} g(x)$$

exist. Then

1.  $\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x)$
2.  $\lim_{x \rightarrow a} [cf(x)] = c \lim_{x \rightarrow a} f(x)$
3.  $\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$
4.  $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$  if  $\lim_{x \rightarrow a} g(x) \neq 0$
5.  $\lim_{x \rightarrow a} c = c$
6.  $\lim_{x \rightarrow a} x = a$
7.  $\lim_{x \rightarrow a} [f(x)]^n = \left[ \lim_{x \rightarrow a} f(x) \right]^n$ , where  $n$  is a positive integer.
8.  $\lim_{x \rightarrow a} x^n = a^n$ , where  $n$  is a positive integer.
9.  $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}$  where  $n$  is a positive integer and if  $n$  is even, then we assume that  $\lim_{x \rightarrow a} f(x) > 0$ .
10.  $\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{\lim_{x \rightarrow a} x}$  where  $n$  is a positive integer and if  $n$  is even, then we assume that  $a > 0$ .

REMARK 1. Note that *all these properties also hold for the one-sided limits.*

REMARK 2. The analogues of the *laws 1-3 also hold when  $f$  and  $g$  are vector functions* (the product in Law 3 should be interpreted as a dot product).

EXAMPLE 3. *Compute the limit:*

$$\lim_{x \rightarrow -1} (7x^5 + 2x^3 - 8x^2 + 3) =$$

REMARK 4. If we had defined  $f(x) = 7x^5 + 2x^3 - 8x^2 + 3$  then Example 3 would have been,

$$\lim_{x \rightarrow -1} f(x) = \lim_{x \rightarrow -1} (7x^5 + 2x^3 - 8x^2 + 3) = 7(-1)^5 + 2(-1)^3 - 8(-1)^2 + 3 = -14 = f(-1)$$

EXAMPLE 5. Compute the limit:

$$\lim_{x \rightarrow -2} \frac{x^2 + x + 1}{x^3 - 10} =$$

REMARK 6. The function from Example 5 also satisfies "direct substitution property":

$$\lim_{x \rightarrow a} f(x) = f(a).$$

Later we will say that such functions are *continuous*. Note that in both examples it was important that  $a$  in the domain of  $f$ .

EXAMPLE 7. Compute the limit:

$$\lim_{x \rightarrow 3} \frac{x - 3}{x^2 - 9}$$

EXAMPLE 8. Compute the limit:

$$\lim_{x \rightarrow 1} \frac{x - 1}{x^2 - 4x + 3}$$

EXAMPLE 9. Given

$$g(x) = \begin{cases} x^2 + 4, & \text{if } x \leq -1 \\ 2 - 3x & \text{if } x > -1 \end{cases}$$

Compute the limits:

(a)  $\lim_{x \rightarrow 4} g(x)$

(b)  $\lim_{x \rightarrow -1} g(x)$

EXAMPLE 10. Evaluate these limits.

(a)  $\lim_{x \rightarrow 4} \frac{x^{-1} - 0.25}{x - 4}$

(b)  $\lim_{x \rightarrow 0} \frac{(x + 5)^2 - 25}{x}$

(c)  $\lim_{x \rightarrow -1} \frac{|x + 1|}{x + 1}$

(d)  $\lim_{x \rightarrow -1} \frac{x^2 + x}{|x + 1|}$

(e)  $\lim_{x \rightarrow 0^-} \left\{ \frac{1}{x} - \frac{1}{|x|} \right\}$

$$(f) \lim_{x \rightarrow 0} \frac{\sqrt{6-x} - \sqrt{6}}{x}$$

**Conclusion** from the above examples:

To calculate the limit of  $f(x)$  as  $x \rightarrow a$ :

PLUG IN  $x = a$  if  $a$  is in the domain of  $f$ .

Otherwise "FACTOR" or "MULTIPLY BY CONJUGATE" and then plug in.

Consider one sided limits if necessary.

**Squeeze Theorem.** Suppose that for all  $x$  in an interval containing  $a$  (except possibly at  $x = a$ )

$$g(x) \leq f(x) \leq h(x)$$

and  $\lim_{x \rightarrow a} g(x) = L = \lim_{x \rightarrow a} h(x)$ . Then

$$\lim_{x \rightarrow a} f(x) = L.$$

**Corollary.** Suppose that for all  $x$  in an interval containing  $a$  (except possibly at  $x = a$ )

$$|f(x)| \leq h(x) \quad (\text{equivalently, } -h(x) \leq f(x) \leq h(x))$$

and  $\lim_{x \rightarrow a} h(x) = 0$ . Then

$$\lim_{x \rightarrow a} f(x) = 0.$$

EXAMPLE 11. Given  $3x \leq f(x) \leq x^3 + 2$  for  $0 \leq x \leq 2$ . Find  $\lim_{x \rightarrow 1} f(x)$

EXAMPLE 12. *Evaluate:*

(a)  $\lim_{x \rightarrow 0} x \sin \frac{1}{x}$

(b)  $\lim_{t \rightarrow 0} (t^5) \cos^3\left(\frac{1}{t^2}\right)$

EXAMPLE 13. *Is there a number  $c$  such that*

$$\lim_{x \rightarrow -2} \frac{3x^2 + cx + c + 3}{x^2 + x - 2}$$

*exists? If so, find the value  $c$  and the value of the limit.*