

8.4: Integration Of Rational Functions By Partial Fractions

EXAMPLE 1. Evaluate the following integrals:

$$(a) \int \frac{2x - 5}{x^2 - 5x + 4} dx$$

$$(b) \int \frac{x - 6}{x^2 - 5x + 4} dx$$

We will use so called Partial Fraction Decomposition

Partial Fraction Decomposition Process

Rational function: $f(x) = \frac{P(x)}{Q(x)}$, where $P(x)$ and $Q(x)$ are polynomials.

Step 0. $\deg P(x) < \deg Q(x)$.

Important to remember: Partial fractions can only be done if the degree of the numerator is **strictly less** than the degree of denominator. (Otherwise, you must first do **long division**.)

Step 1. Factor the denominator as much as possible.

Step 2. For each **linear factor** in the denominator use the following table:

Factor in denominator	Term in partial fraction decomposition
$ax + b$	$\frac{A}{ax + b}$
$(ax + b)^2$	$\frac{A_1}{ax + b} + \frac{A_2}{(ax + b)^2}$
$(ax + b)^k$	$\frac{A_1}{ax + b} + \frac{A_2}{(ax + b)^2} + \cdots + \frac{A_k}{(ax + b)^k}$

For each **irreducible quadratic factor** in the denominator use the following table:

Factor in denominator	Term in partial fraction decomposition
$ax^2 + bx + c$	$\frac{Ax + B}{ax^2 + bx + c}$
$(ax^2 + bx + c)^2$	$\frac{A_1x + B_1}{ax^2 + bx + c} + \frac{A_2x + B_2}{(ax^2 + bx + c)^2}$
$(ax^2 + bx + c)^k$	$\frac{A_1x + B_1}{ax^2 + bx + c} + \frac{A_2x + B_2}{(ax^2 + bx + c)^2} + \cdots + \frac{A_kx + B_k}{(ax^2 + bx + c)^k}$

EXAMPLE 2. Evaluate $I = \int \frac{3x + 11}{x^2 - x - 6} dx$

EXAMPLE 3. Evaluate $I = \int \frac{x^2 + 1}{x^2 - x} dx$

EXAMPLE 4. Evaluate $I = \int \frac{x^2}{(x - 3)(x + 2)^2} dx$

EXAMPLE 5. Integrate $I = \int \frac{x^4 - x^3 - 12x^2 + 10}{x^3 - 4x^2} dx$

EXAMPLE 6. Write out the form of the partial fraction decomposition of the following function:

$$f(x) = \frac{x^3 + 5x^2 - 2012}{x(x+12)(x^2+2x-3)(x^2+x+1)(x^2+25)^3}$$