

**Fall 2012 Math 152**

Week in Review 4

courtesy: *Oksana Shatalov*

(covering Sections 8.1&amp;8.2 and Exam 1 Review )

**8.1: Integration By Parts****Key Points**

$$\int u \, dv = uv - \int v \, du$$

$$\int_a^b u \, dv = uv|_a^b - \int_a^b v \, du \quad \text{where} \quad uv|_a^b = u(b)v(b) - u(a)v(a).$$

To choose  $u$  use the LIATE rule, in order of preference for  $u$ :

Logarithms

Inverse Trigonometric Functions

Algebraic, powers of  $x$ 

Trigonometric functions

Exponential functions

**Examples**

1. Compute the following integrals

$$(a) \quad I = \int \frac{x}{3} \sin \frac{x-1}{3} \, dx$$

$$(b) \quad I = \int_0^1 \frac{x^2}{e^{10x}} \, dx.$$

$$(c) \quad I = \int \arctan \frac{1}{x} \, dx$$

**8.2: Trigonometric Integrals****Key Points**

- How to evaluate  $\int \sin^n x \cos^m x \, dx$

1. If  $n$  is odd use substitution  $u = \cos x$  ( Strip out one sine and convert the rest to cosine.)
2. If  $m$  is odd use substitution  $u = \sin x$  ( Strip out one cosine and convert the rest to sine.)
3. If both  $n$  and  $m$  are odd use 1 or 2.
4. If both  $n$  and  $m$  are even, use the half-angle identities:

$$\sin^2 x = \frac{1}{2}(1 - \cos 2x); \quad \cos^2 x = \frac{1}{2}(1 + \cos 2x); \quad \sin x \cos x = \frac{1}{2} \sin 2x$$

- <sup>1</sup> How to evaluate  $\int \sec^n x \tan^m x \, dx$

1. If  $n$  is even use formula  $\sec^2 x = 1 + \tan^2 x$  and substitution  $u = \tan x \Rightarrow du = \sec^2 x \, dx$ .
2. If  $m$  is odd use formula  $\tan^2 x = \sec^2 x - 1$  and substitution  $u = \sec x \Rightarrow du = \sec x \tan x \, dx$

- How to evaluate  $\int \sin(Ax) \cos(Bx) \, dx$ ,  $\int \sin(Ax) \sin(Bx) \, dx$ ,  $\int \cos(Ax) \cos(Bx) \, dx$

Use the following identities: (The identities below need not be memorized for the exam)

$$\begin{aligned}\sin A \cos B &= \frac{1}{2} (\sin(A - B) + \sin(A + B)) \\ \sin A \sin B &= \frac{1}{2} (\cos(A - B) - \cos(A + B)) \\ \cos A \cos B &= \frac{1}{2} (\cos(A - B) + \cos(A + B))\end{aligned}$$

## Examples

2. Compute the following integrals

(a)  $I = \int \frac{\sec^4 x}{\tan^7 x} \, dx$

(b)  $I = \int \sin^5 x \sqrt{\cos x} \, dx$

(c)  $I = \int \frac{\cos^5(\ln x)}{x} \, dx$

(d)  $\int \cos^2(3x) \sin^2(3x) \, dx$

## General Exam 1 Review

3. If  $F(x) = \int_0^{e^x} \cos(t^2) \, dt$  what is  $F'(x)$ ?

4. Find the area between the curves  $y = 3x^2$  and  $y = 6x$  from  $x = 0$  to  $x = 4$ .

5. Compute  $\int_0^1 \frac{2x}{\sqrt[4]{x^2 + 1}} \, dx$ .

6. After an appropriate substitution, the integral  $\int_{-1}^2 3x^5 \sqrt{x^3 + 1} \, dx$  is equivalent to which of the following?

(a)  $\int_0^3 (u^{3/2} - u^{1/2}) \, du$     (b)  $\int_{-1}^2 (u^{1/2} - u^{3/2}) \, du$     (c)  $\int_0^3 (u^{1/2} + u^{3/2}) \, du$

(d)  $\int_0^9 (u^{3/2} - u^{1/2}) \, du$     (e)  $\int_{-1}^2 x^3 u^{1/2} \, du$

7. Determine the area of the region bounded by the  $x$ -axis, the curve  $y = x^4$  and tangent line to this curve at the point  $(-1, 1)$ .
8. The height of a monument is 20 m. A horizontal cross-section at a distance  $x$  meters from the top is an isosceles triangle with base  $x/4$  meters and height  $x/3$  meters. Set up, but *do not evaluate*, an integral for the volume of the monument.

---

<sup>1</sup>Integral  $\int \csc^n x \cot^m x \, dx$  can be found by similar methods because of the identity  $1 + \cot^2 x = \csc^2 x$ .

9. Find the volume of the solid whose base is the area enclosed by  $y = e^x$  and  $y = e^{-x}$  from  $[0, 1]$  with cross-sections perpendicular to the  $x$ -axis that are equilateral triangles.
10. Find the volume of the solid formed by rotating the region bounded by  $x = 0$ ,  $y = \ln(x)$ ,  $y = 0$ , and  $y = 3$  about the  $y$ -axis.
11. Using **cylindrical shells** which of the following integrals gives the volume of the solid formed by rotating the region bounded by  $y = \sqrt{x}$  and  $y = x^2$  about the line  $x = -2$ ?
- (a)  $2\pi \int_0^1 (x-2)(\sqrt{x}-x^2) dx$       (b)  $2\pi \int_0^1 (x+2)(\sqrt{x}-x^2) dx$       (c)  $\pi \int_0^1 (x^2 - \sqrt{x})^2 dx$
- (d)  $\pi \int_0^1 [(y^2-2)^2 - (\sqrt{y}-2)^2] dy$       (e)  $\pi \int_0^1 (y^2 - \sqrt{y})(y+2) dy$
12. A 15-Newton weight is suspended vertically at the end of a 30 m long rope. The rope weighs 6 Newtons. How much work (in Newton-m) is required to pull the weight to the top?
13. If a 25J work is required to keep a spring 1m beyond its natural length, how much work is done in stretching the spring from 2m to 4m beyond its natural length?
14. A rectangular swimming pool 20 m long, 10 m wide and 3 m deep is full of water (density =  $\rho$  kg/m<sup>3</sup>). What is the work required to pump all the water out of the top of the pool? (Leave your answer in terms of density  $\rho$  and the gravitational constant  $g$ .)
15. A tank of water is a trough 10 feet long and has a vertical cross section in the shape of a semi circle with radius 4 feet, diameter at the top. The tank is filled with water. The water is pumped from a spout at the top of the tank that is 0.5ft high. Find the work needed to pump out the water until the water level is 1m from the bottom. **Just set up the integral** (assume that the density of water is 62.5lb/ft<sup>3</sup>.)
16. A tank has the shape of an upright circular cone with height 15m and radius 3m. In addition, there is a 1 meter high spout at the top of the cone from which the water exits the tank. If the tank is initially full to a water depth of 9m, find the work required to pump all of the water out of the spout. **Just set up the integral.** (Leave your answer in terms of density  $\rho$  and the gravitational constant  $g$ .)
17. Find the average value of  $f(x) = \sin^2 x$  over the interval  $\left[0, \frac{\pi}{4}\right]$ .
18. Compute the following integrals:
- (a)  $I = \int_0^1 (x^2 + 4x - 1)e^x dx$ .
- (b)  $\int \sqrt{x} \ln(x) dx$