

3.8: Higher Derivatives

The derivative of a differentiable function f is also a function and it may have a derivative of its own:

$$(f')' = f'' \quad \text{second derivative}$$

$$f''(x) = \frac{d}{dx}(f'(x)) = \frac{d}{dx}\left(\frac{d}{dx}f(x)\right)$$

Alternative Notation: If $y = f(x)$ then

$$y'' = f''(x) = \frac{d^2y}{dx^2} = D^2f(x).$$

Similarly, the **third derivative** $f''' = (f'')'$ or

$$y''' = f'''(x) = \frac{d}{dx}\left(\frac{d^2y}{dx^2}\right) = \frac{d^3y}{dx^3} = D^3f(x).$$

In general, the n^{th} derivative of $y = f(x)$ is denoted by $f^{(n)}(x)$:

$$y^{(n)} = f^{(n)}(x) = \frac{d}{dx}\left(\frac{d^{n-1}y}{dx^{n-1}}\right) = D^n f(x).$$

EXAMPLE 1. If $f(x) = x^5 + 3x + 1$ find $f^{(n)}(x)$

CONCLUSION: If $p(x)$ is a polynomial of degree n then, $p^{(k)}(x) = 0$ for $k \geq n + 1$.

EXAMPLE 2. Find $D^{2013} \sin x$.

Acceleration: If $s(t)$ is the position of an object then the acceleration of the object is the first derivative of the velocity (consequently, the acceleration is the second derivative of the position function.)

$$a(t) = v'(t) = s''(t).$$

EXAMPLE 3. If $s(t) = t^3 - \frac{9}{2}t^2 - 30t + 12$ is the position of a moving object at time t (where $s(t)$ is measured in feet and t is measured in seconds) find the acceleration at the times when the velocity is zero.

Implicit second derivatives:

EXAMPLE 4. Find $y''(x)$ if $x^6 + y^6 = 66$.