## 9.1 The Definition of Function (continued)

# **FUNCTIONS AND SETS**

### Image of a Set

DEFINITION 1. Let  $f: X \to Y$  be a function. If  $A \subseteq X$ , we define f(A), the **image** of A under f, by  $f(A) = \{ y \in Y | y = f(x) \text{ for some } x \in A \}.$ 

EXAMPLE 2. Let  $f: X \to Y$ . Complete:

(a) If 
$$A \subseteq X$$
 then  $f(A) \subseteq \underline{\hspace{1cm}} \subseteq \underline{\hspace{1cm}}$ 

**(b)** 
$$f(X) =$$
\_\_\_\_

(c) 
$$y \in f(A) \Leftrightarrow \underline{\hspace{1cm}}$$

(d) 
$$y \notin f(A) \Leftrightarrow$$

EXAMPLE 3. Let  $f: \mathbb{R} \to \mathbb{R}$  be defined by  $f(x) = \cos x$ . Find the following

(a) 
$$f([-\pi/2, \pi/2])$$

**(b)** 
$$f([-\pi/2, 0])$$

(c) 
$$f([-\pi/2, \pi/2017])$$

(d) 
$$f([-\pi/4, \pi/4])$$

(e) Does 
$$-\frac{1}{2017}$$
 belong to  $f([-\pi/2, \pi/2])$ ?

### **Inverse Image**

DEFINITION 4. Let  $f: X \to Y$  be a function and let B be a subset of its codomain (i.e.  $B \subseteq Y$ ). Then the inverse image of B (written  $f^{-1}(B)$ ) is the set

$$f^{-1}(B) = \{ x \in X | f(x) \in B \}.$$

The inverse image  $f^{-1}(B)$  is a subset of domain of f containing all preimages of points from B under f. EXAMPLE 5. Let  $f: X \to Y$  and  $B \subseteq Y$ . Complete:

- (a)  $f^{-1}(B) \subseteq$ \_\_\_\_\_
- **(b)**  $x \in f^{-1}(B) \Leftrightarrow \underline{\hspace{1cm}}$
- (c)  $x \notin f^{-1}(B) \Leftrightarrow \underline{\hspace{1cm}}$
- (d) If  $x \in X$  and  $y \in Y$ , then

$$f(x)$$
\_\_\_\_Y,  $f(\{x\})$ \_\_\_\_Y,  $f^{-1}(y)$ \_\_\_\_X,  $f^{-1}(\{y\})$ \_\_\_\_X,

EXAMPLE 6. Let  $f: \mathbb{R} \to \mathbb{R}$  be defined by  $f(x) = x^4$ . Find the following:

- (a) f([1,2]) =
- **(b)** f([-2,-1]) =
- (c)  $f^{-1}([1, 16]) =$
- (d)  $f^{-1}([-16, -1]) =$
- (e)  $f^{-1}([-1,1]) =$
- (f)  $f^{-1}(\mathbb{R}) =$
- (g)  $f^{-1}(\mathbb{R}^+) =$

EXAMPLE 7. Let  $A \subseteq \text{dom}(f)$  and  $B \subset \text{codom}(f)$ . Determine the truth or falsehood of the following (a)  $f^{-1}(f(A)) = A$ 

**(b)** 
$$f(f^{-1}(B)) = B$$

EXAMPLE 8. Let  $A_1, A_2 \subseteq \text{dom}(f)$ . Determine the truth or falsehood of the following  $f(A_1) \cap f(A_2) = f(A_1 \cap A_2)$ 

### Summary

Let  $f: X \to Y$  and  $A \subseteq X$  and  $B \subseteq Y$ . Then above definitions imply the following tautologies

- $\forall y \in Y, ((y \in \operatorname{ran}(f)) \Leftrightarrow (\exists x \in X \ni f(x) = y)).$
- $\forall y \in Y, ((y \in f(A)) \Leftrightarrow (\exists x \in A \ni f(x) = y))$ .
- $\forall x \in X, ((x \in f^{-1}(B)) \Leftrightarrow (f(x) \in B))$ .

Also note that

• If  $B \subseteq \operatorname{ran}(f)$  then  $(S = f^{-1}(B)) \Rightarrow (f(S) = B)$ .

EXAMPLE 9. Let  $f : \mathbb{R} \to \mathbb{R}$  be defined by f(x) = 3x + 4 and let  $B = \{x \in \mathbb{R} | x > 0\}$ . Find  $f^{-1}(B)$ . Justify your answer (give a formal proof).

EXAMPLE 10.  $f : \mathbb{R} \to \mathbb{R}$  is defined by f(x) = 5x - 4. Find f([0,1]). Justify your answer (give a formal proof).

PROPOSITION 11. Let  $f: X \to Y$ . If  $A_1 \subseteq A_2 \subseteq X$  then  $f(A_1) \subseteq f(A_2)$ .

Proof.

PROPOSITION 12. Let  $f: X \to Y$ . If  $B_1$  and  $B_2$  are subsets of Y then  $f^{-1}(B_1 \cup B_2) = f^{-1}(B_1) \cup f^{-1}(B_2).$ 

Proof.

EXAMPLE 13. (cf. Example 8.) Let  $f: X \to Y$  and  $A_1$  and  $A_2$  be subsets of X.

(a) Prove that  $f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$ .

(b) Prove that if, in addition, f is an injective function, then  $f(A_1 \cap A_2) = f(A_1) \cap f(A_2)$ .

EXAMPLE 14. (cf. Example 7.) Let  $f: X \to Y$  and B be a subset of Y.

(a) Prove that  $f(f^{-1}(B)) \subseteq B$ .

(b) Prove that if, in addition, f is an surjective function, then  $f(f^{-1}(B) = B)$ .