

1 Mathematical Reasoning

Prove: “For every odd integer n , the integer $3n+7$ is even.”

1.1 Statements

DEFINITION 1. *A statement is any declarative sentence that is either true or false.*

A statement cannot be neither true nor false and it cannot be both true and false.

1. The integer 5 is odd.
2. The integer 24 is prime.
3. $15 + 7 = 22$
4. Apple manufactures computers.
5. Apple manufactures the world’s best computers.
6. Did you buy IBM?
7. I am telling a lie.

An open sentence is any declarative sentence containing one or more variables that is not a statement but becomes a statement when the variables are assigned values.

8. $x + 5 = 7$
9. He is a student.
10. $x^2 + y^2 = 1$

Use quantifiers to transfer an open sentence to a statement.

Universal: $\forall x$ means for all/for every assigned value a of x .

Existential: $\exists x$ means that for some assigned vales a of x .

11. The area of a rectangle is its length times its width.

Quantifiers:

12. A triangle may be equilateral.

Quantifiers:

13. $15 - 5 = 10$

Quantifiers:

14. The sum of an even integer and an odd integer is even.

Quantifiers:

15. All positive real numbers have a square root.

Quantifiers:

16. A real-valued function that is continuous at 0 is not necessarily differentiable at 0.

Quantifiers:

NEGATIONS

DEFINITION 2. If P is a statement, then the **negation** of P , written $\neg P$ (read “not P ”), is the statement “ P is false”.

17. All continuous functions are differentiable.

18. P : Barack Obama won the Nobel Peace Prize.

19. P : $5^3 = 120$ $\neg P$: _____

20. $P(x)$: $x^2 + x + 1 = 0$ $\neg P(x)$: _____

21. $P(x, y)$: $x^4 + y^4 = 0$ $\neg P(x, y)$: _____

22. P : If n is an odd integer then $3n + 7$ is odd.

$\neg P$ _____

23. P : Every car on the parking lot #47 was with valid permit.

$\neg P$ _____

24. P : There exist real numbers a and b such that $(a + b)^2 = a^2 + b^2$.

$\neg P$ _____

Rules to negate statements with quantifiers:

25. P : For every even integer n there exists an integer m such that $n = 2m$.

$\neg P$ _____

26. P : There exists a positive integer n such that $m(n + 5) < 1$ for every integer m .

$\neg P$ _____

EXAMPLE 3. P : If n is an integer and n^2 is a multiple of 4 then n is a multiple of 4.

Question: Is the following “proof” valid?

Let $n = 6$. Then $n^2 = 6^2 = 36$ and 36 is a multiple of 4, but 6 is not a multiple of 4. Therefore, the statement P is FALSE.□

1.2 Compound Statements

1. P : Some math tests are long.
2. Q : Some math tests are difficult.

Logical connectivity	write	read	meaning
Conjunction	$P \wedge Q$	P and Q	Both P and Q are true
Disjunction	$P \vee Q$	P or Q	P is true or Q is true

TRUTH TABLES

P : Ben is a student.

Q : Ben is a teaching assistant.

P	Q	$P \wedge Q$	$Q \wedge P$	$P \vee Q$	$Q \vee P$

EXAMPLE 4. Rewrite the following open statements using disjunction or conjunction.

(a) P : $|x| \geq 10$.

(b) P : $|x| < 10$.

DEFINITION 5. We say that two compound statements are logically equivalent (write “ \equiv ”) if they have the same truth tables, which means they both are true or both are false.

EXAMPLE 6. Let P and Q be statement forms. Determine whether the compound statements $\neg P \wedge Q$ and $\neg P \vee Q$ are logically equivalent (i.e. both true or both false).

P	Q			

Some Fundamental Properties of Logical Equivalence

THEOREM 7. For the statement forms P , Q and R ,

- $\neg(\neg P) \equiv P$
- *Commutative Laws*
 $P \vee Q \equiv Q \vee P$
 $P \wedge Q \equiv Q \wedge P$
- *Associative Laws*
 $P \vee (Q \vee R) \equiv (P \vee Q) \vee R$
 $P \wedge (Q \wedge R) \equiv (P \wedge Q) \wedge R$
- *Distributive Laws*
 $P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)$
 $P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)$
- *De Morgan's Laws*
 $\neg(P \vee Q) \equiv (\neg P) \wedge (\neg Q)$
 $\neg(P \wedge Q) \equiv (\neg P) \vee (\neg Q)$

Proof. Each part of the theorem is verified by means of a truth table.

P	Q					

EXAMPLE 8. Rewrite the rules to negate statements with quantifiers in terms of logical equivalence:

$$\neg(\forall x, P(x)) \equiv$$

$$\neg(\exists x \ni P(x)) \equiv$$

$$\neg(\forall x, (P(x) \vee Q(x))) \equiv$$

$$\neg(\forall x, (P(x) \wedge Q(x))) \equiv$$

$$\neg(\exists x \ni (P(x) \vee Q(x))) \equiv$$

$$\neg(\exists x \ni (P(x) \wedge Q(x))) \equiv$$

EXAMPLE 9. *Negate:*

P : *There exists a prime number p which is greater than 7 and less than 10.*

$\neg P$ _____

Tautologies and Contradictions

Tautology: statement that is always true

Contradiction: statement that is always false

P	$\neg P$	$P \vee (\neg P)$	$P \wedge (\neg P)$
T			
F			

1.3 Implications

DEFINITION 10. *Let P and Q be statements. The **implication** $P \Rightarrow Q$ (read “ P implies Q ”) is the statement “If P is true, then Q is true.”*

EXAMPLE 11. *If n is odd, then $3n + 7$ is even.*

The truth table for implication:

P	Q	$P \Rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

EXAMPLE 12. P : *You earn an A on the final exam.*

Q : *You get an A for your final grade.*

$P \Rightarrow Q$

Different ways of expressing $P \Rightarrow Q$:

If P is true, then Q is true

Q is true if P is true

P implies Q

P is true only if Q is true

P is sufficient for Q

Q is necessary for P .

EXAMPLE 13. For a triangle T , let

$P(T)$: T is equilateral $Q(T)$: T is isosceles.

State $P(T) \Rightarrow Q(T)$ in a variety of ways:

EXAMPLE 14. P : The function $f(x) = \sin x$ is differentiable everywhere.

Q : The function $f(x) = \sin x$ is continuous everywhere.

$P \Rightarrow Q$

$Q \Rightarrow P$

Proving Statements Containing Implications.

Properties of Integers:

1. *The negative of every integer is an integer.*
2. *The sum (and difference) of every two integers is an integer.*
3. *The product of every two integers is an integer.*

• **DIRECT PROOF**

- Assume that P is true.
- Draw out consequences of P .
- Use these consequences to prove Q is true.

EXAMPLE 15. *If n is an even integer, then $5n^5$ is an even integer.*

EXAMPLE 16. Evaluate the proposed proof of the following result:

If a is an even integer and b is an odd integer, then $3a - 5b$ is odd.

Proof. Let a be an even integer and b be an odd integer. Then $a = 2n$ and $b = 2n + 1$ for some integer n . Therefore,

$$3a - 5b = 3(2n) - 5(2n + 1) = 6n - 10n - 5 = -4n - 5 = 2(-2n - 2) - 1.$$

Since $-2n - 2$ is an integer, $3a - 5b$ is odd. \square

EXAMPLE 17. *The sum of every two odd integers is even.*

EXAMPLE 18. *Let x be an integer. If $5x - 7$ is odd, then $9x + 2$ is even.*

- **PROOF BY CASES**

EXAMPLE 19. *If n is an integer, then $n^2 + 3n + 4$ is an even integer.*

Hint: Use the following fact: “*Every integer number is either even. or odd.*”

Negating an implication: **Counterexamples**

$$\neg(P \Rightarrow Q) \equiv P \wedge (\neg Q)$$

REMARK 20. The negation of an implication is not an implication!

EXAMPLE 21. *Negate the statement: “For all x , $P(x) \Rightarrow Q(x)$.”*

The value assigned to the variable x that makes $P(x)$ true and $Q(x)$ false is called a **counterexample** to the statement “For all x , $P(x) \Rightarrow Q(x)$.”

EXAMPLE 22. Disprove the following statement:

If a real-valued function is continuous at some point, then this function is differentiable there.

Necessary and Sufficient Conditions

$P \Rightarrow Q$ can be expressed as

P is sufficient for Q .

or

Q is necessary for P .

Equivalently,

In order for Q to be true it is sufficient that P be true.

or

Q must be true in order to P to be true.

EXAMPLE 23. Consider the following open sentences

$P(x)$: x is a multiple of 4. $Q(x)$: x is even. Complete:

- “ $\forall x, P(x) \Rightarrow Q(x)$ ” is _____.
- $P(x)$ is a _____ condition for Q to be true.
- $Q(x)$ is a _____ condition for $P(x)$ to be true.
- $Q(x)$ is not a _____ condition for $P(x)$ to be true.

EXAMPLE 24. Consider the following open sentences

$P(f)$: f is a differentiable function.

$Q(f)$: f is a continuous function.

Complete:

- “ $\forall f, P(f) \Rightarrow Q(f)$ ” is _____.
- “ $\forall f, Q(f) \Rightarrow P(f)$ ” is _____.
- $Q(f)$ is a _____ condition for f to be differentiable, but not a _____ condition.
- $P(f)$ is a _____ condition for f to be continuous.

REMARK 25. Note however, if $P \Rightarrow Q$ is true, then it is not necessary that P is true in order for Q to be true. Even if Q is true, P may be false.

1.4 Contrapositive and Converse

Contrapositive

DEFINITION 26. The statement $\neg Q \Rightarrow \neg P$ is called the **contrapositive** of the statement $P \Rightarrow Q$.

EXAMPLE 27. Let P and Q be statement forms. Prove that $\neg Q \Rightarrow \neg P$ is logically equivalent to $P \Rightarrow Q$.

P	Q	$P \Rightarrow Q$	$\neg Q$	$\neg P$	$\neg Q \Rightarrow \neg P$

Methods to prove an implication $P \Rightarrow Q$ (continued)

- **CONTRAPOSITIVE PROOF** (based on the equivalence $(P \Rightarrow Q) \equiv (\neg Q \Rightarrow \neg P)$)
 - Assume that $\neg Q$ is true.
 - Draw out consequences of $\neg Q$.
 - Use these consequences to prove $\neg P$ is true.
 - It follows that $P \Rightarrow Q$.

REMARK 28. If you use a contrapositive method, you must declare it in the beginning and then state **what is sufficient to prove**.

EXAMPLE 29. Let x be an integer. If $5x - 7$ is even, then x is odd.

Converse

DEFINITION 30. The statement $Q \Rightarrow P$ is called a **converse** of the statement $P \Rightarrow Q$.

Question: Are the statements $P \Rightarrow Q$ and $Q \Rightarrow P$ logically equivalent? _____

EXAMPLE 31. If m and n are odd integers then $m + n$ is even.

Biconditional “ \Leftrightarrow ”

DEFINITION 32. The statement $P \Leftrightarrow Q$ (or P iff Q) is the statement $(P \Rightarrow Q) \wedge (Q \Rightarrow P)$.

P	Q	$P \Rightarrow Q$	$Q \Rightarrow P$	$P \Leftrightarrow Q$
T	T			
T	F			
F	T			
F	F			

THEOREM 33. Let \vec{a} and \vec{b} be two non zero vectors. Then \vec{a} is orthogonal to \vec{b} iff $\vec{a} \cdot \vec{b} = 0$.

THEOREM 34. *Let n be an integer. Then n is even if and only if n^2 is even.*

Proof.

REMARK 35. $(P \Leftrightarrow Q) \equiv (\neg P \Leftrightarrow \neg Q)$

COROLLARY 36. *Let n be an integer. Then n is odd iff n^2 is odd.*

Methods to prove an implication $P \Rightarrow Q$ (continued)

EXAMPLE 37. Let S and C be statement forms. Prove that $\neg S \Rightarrow (C \wedge \neg C)$ is logically equivalent to S .

• PROOF BY CONTRADICTION

- Assume that P is true.
- To derive a contradiction, assume that $\neg Q$ is true.
- Prove a false statement C , using negation $\neg(P \Rightarrow Q) \equiv (P \wedge \neg Q)$.
- Prove $\neg C$. It follows that Q is true. (The statement $C \wedge \neg C$ must be false, i.e. a contradiction.)

PROPOSITION 38. If m and n are integers, then $m^2 - 4n \neq 2$.

Proof.

PROPOSITION 39. *Let a, b , and c be integers. If $a^2 + b^2 = c^2$ then a or b is an even integer.*

Proof.

DEFINITION 40. A real number x is **rational** if $x = \frac{m}{n}$ for some integer numbers m and n . Also, x is **irrational** if it is not rational, that is

PROPOSITION 41. The number $\sqrt{2}$ is irrational.