

## Chapter 5: The Integers

### 5.1: Axioms and Basic Properties

Operations on the set of integers,  $\mathbf{Z}$ : *addition* and *multiplication* with the following properties:

**A1.** Addition is **associative**:

**A2.** Addition is **commutative**:

**A3.**  $\mathbf{Z}$  has an **identity** element with respect to addition namely, the integer 0.

**A4.** Every integer  $x$  in  $\mathbf{Z}$  has an **inverse** w.r.t. addition, namely, its negative  $-x$  :

**A5.** Multiplication is **associative**:

**A6.** Multiplication is **commutative**:

**A7.**  $\mathbf{Z}$  has an **identity** element with respect to multiplication namely, the integer 1. (and  $1 \neq 0$ .)

**A8. Distributive Law:**

REMARK 1. We do not prove A1-A8. We take them as **axioms**: statements we *assume* to be true about the integers.

We use  $xy$  instead  $x \cdot y$  and  $x - y$  instead  $x + (-y)$ .

PROPOSITION 2. *Let  $a, b, c \in \mathbf{Z}$ .*

**P1.** *If  $a + b = a + c$  then  $b = c$ . (cancellation law for addition)*

**P2.**  $a \cdot 0 = 0 \cdot a = 0$ .

**P3.**  $(-a)b = a(-b) = -(ab)$

**P4.**  $-(-a) = a$

**P5.**  $(-a)(-b) = ab$

**P6.**  $a(b - c) = ab - ac$

**P7.**  $(-1)a = -a$

**P8.**  $(-1)(-1) = 1$ .

Proof

$\mathbf{Z}$  contains a subset  $\mathbf{Z}^+$ , called the **positive integers**, that has the following properties:

**A9. Closure property:**  $\mathbf{Z}^+$  is closed w.r.t. addition and multiplication:

**A10. Trichotomy Law:** for all  $x \in \mathbf{Z}$  exactly one is true:

PROPOSITION 3. *If  $x \in \mathbf{Z}$ ,  $x \neq 0$ , then  $x^2 \in \mathbf{Z}^+$ .*

*Proof.*

COROLLARY 4.  $\mathbf{Z}^+ = \{1, 2, 3, \dots, n, n + 1, \dots\}$

*Proof.*

### Inequalities (the order relation less than)

DEFINITION 5. For  $x, y \in \mathbf{Z}$ ,  $x < y$  if and only  $y - x \in \mathbf{Z}^+$ .

Note that  $\mathbf{Z}^+ = \{n \in \mathbf{Z} | n > 0\}$ .

PROPOSITION 6. For all  $a, b \in \mathbf{Z}$ :

**Q1.** Exactly one of the following holds:  $a < b$ ,  $b < a$ , or  $a = b$ .

**Q2.** If  $a > 0$  then  $-a < 0$ ; if  $a < 0$  then  $-a > 0$ .

**Q3.** If  $a > 0$  and  $b > 0$  then  $a + b > 0$  and  $ab > 0$ .

**Q4.** If  $a > 0$  and  $b < 0$  then  $ab < 0$ .

**Q5.** If  $a < 0$  and  $b < 0$  then  $ab > 0$ .

**Q6.** If  $a < b$  and  $b < c$  then  $a < c$ .

**Q7.** If  $a < b$  and  $a + c < b + c$ .

**Q8.** If  $a < b$  and  $c > 0$  then  $ac < bc$ .

**Q9.** If  $a < b$  and  $c < 0$  then  $ac > bc$ .

Proof.

**A11. The Well Ordering Principle** Every nonempty subset on  $\mathbf{Z}^+$  has a smallest element; that is, if  $S$  is a nonempty subset of  $\mathbf{Z}^+$ , then there exists  $a \in S$  such that  $a \leq x$  for all  $x \in S$ .

PROPOSITION 7. *There is no integer  $x$  such that  $0 < x < 1$ .*

*Proof.*

COROLLARY 8. *1 is the smallest element of  $\mathbf{Z}^+$ .*

COROLLARY 9. *The only integers having multiplicative inverses in  $\mathbf{Z}$  are  $\pm 1$ .*

## 5.2: Induction

*"Domino Effect"*

**Step 1.** The first domino falls.

**Step 2.** When any domino falls, the next domino falls.

**Conclusion.** All dominos will fall!

**THEOREM 10. (First Principle of Mathematical Induction)** *Let  $P(n)$  be a statement about the positive integer  $n$ . Suppose that  $P(1)$  is true. Whenever  $k$  is a positive integer for which  $P(k)$  is true, then  $P(k + 1)$  is true. Then  $P(n)$  is true*

*Proof.*

### Strategy

The proof by induction consists of the following steps:

**Basic Step:** Verify that  $P(1)$  is true.

**Induction hypothesis:** Assume that  $k$  is a positive integer for which  $P(k)$  is true .

**Inductive Step:** With the assumption made, prove that  $P(k + 1)$  is true.

**Conclusion:**  $P(n)$  is true for every positive integer  $n$ .

EXAMPLE 11. *Prove by induction the formula for the sum of the first  $n$  positive integers*

$$1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}. \quad (1)$$

EXAMPLE 12. *Find the sum of all odd numbers from 1 to  $2n+1$  ( $n \in \mathbf{Z}^+$ ).*

EXAMPLE 13. *Prove by induction the following formula*

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

*Paradox: All horses are of the same color.*

*Question: What's wrong in the following "proof" of G. Pólya?*

**Basic Step.** If there is only one horse, there is only one color.

**Inductive step.** Assume as induction hypothesis that within any set of  $k$  horses, there is only one color. Now look at any set of  $k+1$  horses. Number them:  $1, 2, 3, \dots, k, k+1$ . Consider the sets  $\{1, 2, 3, \dots, k\}$  and  $\{2, 3, 4, \dots, k+1\}$ . Each is a set of only  $k$  horses, therefore within each there is only one color. But the two sets overlap, so there must be only one color among all  $k+1$  horses.

### 5.3: The Division Algorithm And Greatest Common Divisor

THEOREM 14. (Division Algorithm) *Let  $a \in \mathbf{Z}$ ,  $b \in \mathbf{Z}^+$ . Then there exist unique integers  $q$  and  $r$  such that*

$$a = bq + r, \quad \text{where } 0 \leq r < b.$$

#### Divisors

DEFINITION 15. *Let  $a$  and  $b$  be integers. We say that  $b$  **divides**  $a$ , written  $b|a$ , if there is an integer  $c$  such that  $bc = a$ . We say that  $b$  and  $c$  are **factors** of  $a$ , or that  $a$  is **divisible** by  $b$  and  $c$ .*

For example,

EXAMPLE 16. *Prove that  $3|4^n - 1$ , where  $n \in \mathbf{Z}^+$ .*

PROPOSITION 17. *Let  $a, b, c \in \mathbf{Z}$ .*

- (a) *If  $a|1$ , then  $a = \pm 1$ .*
- (b) *If  $a|b$  and  $b|a$ , then  $a = \pm b$ .*
- (c) *If  $a|b$  and  $a|c$ , then  $a|(bx + cy)$  for any  $x, y \in \mathbf{Z}$ .*
- (d) *If  $a|b$  and  $b|c$ , then  $a|c$ .*



**Greatest common divisor (gcd)**

DEFINITION 18. Let  $a$  and  $b$  be integers, not both zero. The **greatest common divisor** of  $a$  and  $b$  (written  $\gcd(a, b)$ , or  $(a, b)$ ) is the largest positive integer  $d$  that divides both  $a$  and  $b$ .

EXAMPLE 19. Find  $\gcd(18, 24)$ .

**Euclidean Algorithm**

is based on the following two lemmas:

LEMMA 20. Let  $a$  and  $b$  be integers. If  $a|b$  then  $\gcd(a, b) = a$ .

LEMMA 21. Let  $a$  and  $b$  be two positive integers such that  $b \geq a$ . Then  $\gcd(a, b) = \gcd(a, b - a)$ .

*Proof.*

COROLLARY 22. Let  $a$  and  $b$  be integers, not both zero. Suppose that there exist integers  $q_1$  and  $r_1$  such that  $b = aq_1 + r_1$ ,  $0 \leq r_1 < a$ . Then  $\gcd(a, b) = \gcd(a, r_1)$ .

**Procedure for finding gcd of two integers (the Euclidean Algorithm)**

EXAMPLE 23. Find  $\gcd(1176, 3087)$ .

EXAMPLE 24. Find integers  $x$  and  $y$  such that  $147 = 1176x + 3087y$ .

COROLLARY 25. If  $d = \gcd(a, b)$  then there exist integers  $x$  and  $y$  such that  $ax + by = d$ . Moreover,  $d$  is the minimal natural number with such property.

### Relatively prime (or coprime) integers

DEFINITION 26. Two integers  $a$  and  $b$ , not both zero, are said to be **relatively prime (or coprime)**, if  $\gcd(a, b) = 1$ .

For example,

Combining the above definition and the proof of Corollary 25, we obtain

THEOREM 27.  $a$  and  $b$  are relatively prime integers if and only if there exist integers  $x$  and  $y$  such that  $ax + by = 1$ .

**THEOREM 28.** *Let  $a, b, c \in \mathbf{Z}$ . Suppose  $a|bc$  and  $\gcd(a, b) = 1$ . Then  $a|c$ .*

*Proof.*

## 5.4: Primes and Unique Factorization

**DEFINITION 29.** *An integer  $p$  greater than 1 is called a **prime** number if the only divisors of  $p$  are  $\pm 1$  and  $\pm p$ . If an integer greater than 1 is not prime, it is called **composite**.*

For example,

### Sieve of Eratosthenes.

The method to find all primes from 2 to  $n$ .

1. Write out all integers from 2 to  $n$ .
2. Select the smallest integer  $p$  that is not selected or crossed out.
3. Cross out all multiples of  $p$  (these will be  $2p, 3p, 4p, \dots$ ; the  $p$  itself should not be crossed out).
4. If not all numbers are selected or crossed out return to step 2. Otherwise, all selected numbers are prime.

**EXAMPLE 30.** *Find all two digit prime numbers.*

2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	

**REMARK 31.** It is sufficient to cross out the numbers in step 3 starting from  $p^2$ , as all the smaller multiples of  $p$  will have already been crossed out at that point. This means that the algorithm is allowed to terminate in step 4 when  $p^2$  is greater than  $n$ . In other words, *if the number  $p$  in step 2 is greater than  $\sqrt{n}$  then all numbers that are already selected or not crossed out are prime.*

**Prime Factorization** of a positive integer  $n$  greater than 1 is a decomposition of  $n$  into a product of primes.

EXAMPLE 32. Write 1224 in standard form (i.e. find its prime factorization).

LEMMA 33. Let  $a$  and  $b$  be integers. If  $p$  is prime and divides  $ab$ , then  $p$  divides either  $a$ , or  $b$ . (Note,  $p$  also may divide both  $a$  and  $b$ .)

*Proof.*

COROLLARY 34. Let  $a_1, a_2, \dots, a_n$  be integers. If  $p$  is prime and divides  $a_1 a_2 \cdot \dots \cdot a_n$ , then  $p$  divides at least one integer from  $a_1, a_2, \dots, a_n$ .

Note that Lemma 33 corresponds to  $n = 2$ . General proof of the above Corollary is by induction.

**THEOREM 35. (Second Principle of Mathematical Induction)** Let  $P(n)$  be a statement about the positive integer  $n$ . Suppose that  $P(1)$  is true. Whenever  $k$  is a positive integer for which  $P(i)$  is true for every positive integer  $i$  such that  $i \leq k$ , then  $P(k + 1)$  is true. Then  $P(n)$  is true for every positive integer  $n$ .

**THEOREM 36. Unique Prime Factorization Theorem.** *Let  $n \in \mathbf{Z}$ ,  $n > 1$ . Then  $n$  is a prime number or can be written as a product of prime numbers. Moreover, the product is unique, except for the order in which the factors appears.*

*Proof.*

**Existence:** Use the Second Principle of Mathematical Induction.

$P(n)$  :

Basic step:

Induction hypothesis:

Inductive step:

**Uniqueness** Use the Second Principle of Mathematical Induction.  $P(n)$  :

Basic step:

Induction hypothesis:

COROLLARY 37. *There are infinitely many prime numbers.*

*Proof.*

EXAMPLE 38. *Prove that if  $a$  is a positive integer of the form  $4n + 3$ , then at least one prime divisor of  $a$  is of the form  $4n + 3$ .*

*Proof.*