

## 14.5: The Chain Rule

Chain Rule for functions of a single variable: If  $y = f(x)$  and  $x = g(t)$  where  $f$  and  $g$  are differentiable functions, then  $y$  is indirectly a differentiable function of  $t$  and

$$\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt}.$$

EXAMPLE 1. Let  $z = x^y$ , where  $x = t^2$ ,  $y = \sin t$ . Compute  $z'(t)$ .

Assume that all functions below have continuous derivatives (ordinary or partial).

- CASE 1:  $z = f(x, y)$ , where  $x = x(t)$ ,  $y = y(t)$  and compute  $z'(t)$ .

*Chain Rule:*

$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}$$

SOLUTION OF EXAMPLE 1:

EXAMPLE 2. The radius of a right circular cone is increasing at a rate of 1.8 cm/s while its height is decreasing at a rate 2.5 cm/s. At what rate is the volume of the cone changing when the radius is 120 cm and the height is 140 cm.

- CASE 2:  $z = f(x, y)$ , where  $x = x(s, t)$ ,  $y = y(s, t)$  and compute  $z_s$  and  $z_t$ .

Chain Rule:

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s}$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial t}$$

Tree diagram:

EXAMPLE 3. Write out the Chain Rule for the case where  $w = f(x, y, z)$  and  $x = x(u, v)$ ,  $y = y(u, v)$  and  $z = z(u, v)$ .

EXAMPLE 4. If  $z = \sin x \cos y$ , where  $x = (s - t)^2$ ,  $y = s^2 - t^2$  find  $z_s + z_t$ .

EXAMPLE 5. If  $u = x^2y + y^3z^2$  where  $x = rse^t$ ,  $y = r + s^2e^{-t}$ ,  $z = rs \sin t$ , find  $u_s$  when  $(r, s, t) = (1, 2, 0)$

**Implicit differentiation:** Suppose that an equation

$$F(x, y) = 0$$

defines  $y$  implicitly as a differentiable function of  $x$ , i.e.  $y = y(x)$ , where  $F(x, y(x)) = 0$  for all  $x$  in the domain of  $y(x)$ . Find  $y'$ :

EXAMPLE 6. Find  $y'$  if  $x^4 + y^3 = 6e^{xy}$ .

Suppose that an equation

$$F(x, y, z) = 0$$

defines  $z$  implicitly as a differentiable function of  $x$  and  $y$ , i.e.  $z = z(x, y)$ , where

$$F(x, y, z(x, y)) = 0$$

for all  $(x, y)$  in the domain of  $z$ . Find the partial derivatives  $z_x$  and  $z_y$ :

EXAMPLE 7. If  $x^4 + y^3 + z^2 + xye^z = 10$  find

(a)  $z_x$  and  $z_y$

(b)  $x_y$  and  $x_z$