


14.7: Maximum and minimum values

Function $y = f(x)$	Function of two variables $z = f(x, y)$
<p>DEFINITION 1. A function $f(x)$ has a local maximum at $x = a$ if $f(a) \geq f(x)$ when x is near a (i.e. in a neighborhood of a). A function f has a local minimum at $x = a$ if $f(a) \leq f(x)$ when x is near a.</p> <p>.</p> <p>If the inequalities in this definition hold for ALL points x in the domain of f, then f has an absolute max (or absolute min) at a</p> <p>If the graph of f has a tangent line at a local extremum, then the tangent line is horizontal: $f'(a) = 0$.</p> 	<p>DEFINITION 2. A function $f(x, y)$ has a local maximum at $(x, y) = (a, b)$ if $f(a, b) \geq f(x, y)$ when (x, y) is near (a, b) (i.e. in a neighborhood of (a, b)). A function f has a local minimum at $(x, y) = (a, b)$ if $f(a, b) \leq f(x, y)$ when (x, y) is near (a, b).</p> <p>If the inequalities in this definition hold for ALL points (x, y) in the domain of f, then f has an absolute maximum (or absolute minimum) at (a, b).</p> <p>If the graph of f has a tangent plane at a local extremum, then the tangent PLANE is horizontal.</p>

THEOREM 3. If f has a local extremum (that is, a local maximum or minimum) at (a, b) and the first-order partial derivatives exist there, then


$$f_x(a, b) = f_y(a, b) = 0 \quad (\text{or, equivalently, } \nabla f(a, b) = 0.)$$

DEFINITION 4. A point (a, b) such that $f_x(a, b) = 0$ and $f_y(a, b) = 0$, or one of this partial derivatives does not exist, is called a **critical point** of f .

At a critical point, a function could have a local max or a local min, or neither.

We will be concerned with two important questions:

- Are there any local or absolute extrema?
- If so, where are they located?




<https://www.slideshare.net/abdulazizuinmlg/multivariate-calculus-mhsw-2>

in \mathbb{R}	in \mathbb{R}^2
close interval $[a, b]$	close set
open interval (a, b)	open set
end points of an interval	boundary points

DEFINITION 5. A **bounded set** in \mathbb{R}^2 is one that contained in some disk.

THE EXTREME VALUE THEOREM:

Function $y = f(x)$	Function of two variables $z = f(x, y)$
If f is continuous on a closed interval $[a, b]$, then f attains an absolute maximum value $f(x_1)$ and an absolute minimum value $f(x_2)$ at some points x_1 and x_2 in $[a, b]$.	If f is continuous on a closed bounded set \mathcal{D} in \mathbb{R}^2 , then f attains an absolute maximum value $f(x_1, y_1)$ and an absolute minimum value $f(x_2, y_2)$ at some points (x_1, y_1) and (x_2, y_2) in \mathcal{D} .



<https://www.slideshare.net/abdulazizuinmlg/multivariate-calculus-mhsw-2>

EXAMPLE 6. Find extreme values of $f(x, y) = x^2 + y^2$.

	Local	Absolute
Maximum		
Minimum		

Domain:

EXAMPLE 7. Find extreme values of $f(x, y) = 5 + \sqrt{1 - x^2 - y^2}$.

	Local	Absolute
Maximum		
Minimum		

Domain:

EXAMPLE 8. Find extreme values of $f(x, y) = x^2 - y^2$.

	Local	Absolute
Maximum		
Minimum		

Domain:

REMARK 9. Example 8 illustrates so called **saddle point** of f . Note that the graph of f crosses its tangent plane at (a, b) .

ABSOLUTE MAXIMUM AND MINIMUM VALUES on a closed bounded set.

THE EXTREME VALUE THEOREM:

To find the absolute maximum and minimum values of a continuous function f on a closed interval $[a, b]$:

1. Find the values of f at the critical points of f in (a, b) .
2. Find the values of f at the endpoints of the interval.
3. The largest of the values from steps 1&2 is the absolute max value; the smallest of the values from steps 1&2 is the absolute min value.

To find the absolute max and min values of a continuous function f on a closed bounded set D :

1. Find the values of f at the critical points of f in D .
2. Find the extreme values of f on the boundary of D . (This usually involves the Calculus I approach for this work.)
3. The largest of the values from steps 1&2 is the absolute maximum value; the smallest of the values from steps 1&2 is the absolute minimum value.

- The quantity to be maximized/minimized is expressed in terms of variables (as few as possible!)
- Any constraints that are presented in the problem are used to reduce the number of variables to the point they are independent,
- After computing partial derivatives and setting them equal to zero you get purely algebraic problem (but it may be hard.)
- Sort out extreme values to answer the original question.

EXAMPLE 10. A lamina occupies the region $D = \{(x, y) : 0 \leq x \leq 3, -2 \leq y \leq 4 - 2x\}$. The temperature at each point of the lamina is given by

$$T(x, y) = 4(x^2 + xy + 2y^2 - 3x + 2y) + 10.$$

Find the hottest and coldest points of the lamina.

Local/Relative Extrema

Second derivatives test:

Suppose f'' is continuous near a and $f'(c) = 0$ (i.e. a is a critical point).

- If $f''(c) > 0$ then $f(c)$ is a local minimum.
- If $f''(c) < 0$ then $f(c)$ is a local maximum.

NOTE:

- If $f''(c) = 0$, then the test gives no information.

Suppose that the second partial derivatives of f are continuous near (a, b) and $\nabla f(a, b) = \mathbf{0}$ (i.e. (a, b) is a critical point).

Let $\mathcal{D} = \mathcal{D}(a, b) = f_{xx}(a, b)f_{yy}(a, b) - [f_{xy}(a, b)]^2$

- If $\mathcal{D} > 0$ and $f_{xx}(a, b) > 0$ then $f(a, b)$ is a local minimum.
- If $\mathcal{D} > 0$ and $f_{xx}(a, b) < 0$ then $f(a, b)$ is a local maximum.
- If $\mathcal{D} < 0$ then $f(a, b)$ is not a local extremum (saddle point).

If $\mathcal{D} = 0$ or does not exist, then the test gives no information. fails.

To remember formula for \mathcal{D} :

$$\mathcal{D} = f_{xx}f_{yy} - [f_{xy}]^2 = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{vmatrix}$$

EXAMPLE 11. Use the Second Derivative Test to confirm that a local cold point of the lamina in the previous Example is $(2, -1)$.

EXAMPLE 12. *The surface $z = 4xy - x^4 - y^4$ has two peaks and one pass. Find them.*