

15.2: Double Integrals over General Regions

All functions below are continuous on their domains.

Let D be a bounded region enclosed in a rectangular region R . We define

$$F(x, y) = \begin{cases} f(x, y) & \text{if } (x, y) \text{ is in } D \\ 0 & \text{if } (x, y) \text{ is in } R \text{ but not in } D. \end{cases}$$



If F is integrable over R , then we say F is *integrable* over D and we define **the double integral of f over D** by

$$\iint_D f(x, y) \, dA = \iint_R F(x, y) \, dA$$

FACT: If $f(x, y) \geq 0$ and f is continuous on the region D then the volume V of the solid S that lies above D and under the graph of f , i.e.

$$S = \{(x, y, z) \in \mathbb{R}^3 \mid 0 \leq z \leq f(x, y), (x, y) \in D\},$$

is

$$V = \iint_D f(x, y) \, dA.$$

EXAMPLE 1. Evaluate the integral

$$\iint_D \sqrt{16 - x^2 - y^2} \, dA$$

where $D = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 16\}$ by identifying it as a volume of a solid.

Properties of double integrals:

- If $D = D_1 \cup D_2$, where D_1 and D_2 do not overlap except perhaps their boundaries then

$$\iint_D f(x, y) \, dA = \iint_{D_1} f(x, y) \, dA + \iint_{D_2} f(x, y) \, dA.$$

- If α and β are real numbers then

$$\iint_D (\alpha f(x, y) + \beta g(x, y)) \, dA = \alpha \iint_D f(x, y) \, dA + \beta \iint_D g(x, y) \, dA.$$

- If we integrate the constant function $f(x, y) = 1$ over D , we get **area** of D :




$$\iint_D 1 \, dA = A(D).$$

EXAMPLE 2. If $D = \{(x, y) \mid x^2 + y^2 \leq 25\}$ then

$$\iint_D dA =$$

Computation of double integral:A plain region of **TYPE I**:

$$D = \{(x, y) \mid a \leq x \leq b, g_1(x) \leq y \leq g_2(x)\}.$$






THEOREM 3. If D is a region of type I such that $D = \{(x, y) \mid a \leq x \leq b, g_1(x) \leq y \leq g_2(x)\}$ then

$$\iint_D f(x, y) \, dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) \, dy \, dx.$$

A plain region of **TYPE II**:

$$D = \{(x, y) \mid c \leq y \leq d, h_1(y) \leq x \leq h_2(y)\}.$$



THEOREM 4. If D is a region of type II s.t. $D = \{(x, y) \mid c \leq y \leq d, h_1(y) \leq x \leq h_2(y)\}$ then

$$\iint_D f(x, y) \, dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) \, dx \, dy.$$

EXAMPLE 5. Evaluate $I = \iint_D 30x^2y \, dA$, where D is the region bounded by the lines $x = 2$, $y = x$ and the hyperbola $xy = 1$ in two different ways (i.e. considering D as a type I and then as a type II region).

EXAMPLE 6. Find the volume of the solid bounded by the cylinder $x^2 + y^2 = 1$ and the planes $x = 0, y = z, z = 0$ in the first octant.

EXAMPLE 7. Evaluate the following iterated integral by reversing the order of integration:

$$I = \int_0^1 \int_{x^2}^1 x^3 \sin(y^3) \, dy \, dx$$