

15.7: Triple integrals in cylindrical coordinates

• Cylindrical coordinates:

$$P(x, y, z) \in \mathbb{R}^3$$

In the cylindrical coordinates P is represented by the ordered triple (r, θ, z) , where r, θ are the polar coordinates of P_{xy} and z is the directed distance from the xy -plane to P :


$$x = \quad \quad \quad y = \quad \quad \quad z =$$

where

$$r^2 = \quad \quad \quad \tan \theta = \quad \quad \quad z = z.$$

REMARK 1. The cylindrical coordinates

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \\ z &= z \\ r &\geq 0, \quad 0 \leq \theta \leq 2\pi \end{aligned}$$



are useful in problems that involve *symmetry about the z-axis*.

<https://i.stack.imgur.com/FgSBF.jpg>

EXAMPLE 2. Find an equation in cylindrical coordinates for the cone

$$z = \sqrt{x^2 + y^2}$$

THEOREM 3. Let $f(x, y, z)$ be a continuous function over a solid $E \subset \mathbb{R}^3$. Let E^* be its image in cylindrical coordinates. Then

$$\iiint_E f(x, y, z) \, dV = \iiint_{E^*} f(r \cos \theta, r \sin \theta, z) \, dV^*,$$

where

$$dV^* = r \, dr \, dz \, d\theta.$$

EXAMPLE 4. *The density at any point of the solid E ,*

$$E = \{(x, y, z) : x^2 + y^2 \leq 9, -1 \leq z \leq 4\},$$

equals to its distance from the axis of E . Find the mass of E .

REMARK 5. If E is a solid region of type I, i.e.

$$E = \{(x, y, z) | (x, y) \in D, \phi_1(x, y) \leq z \leq \phi_2(x, y)\},$$

where D is the projection of E onto the xy -plane then, as we know,

$$\iiint_E f(x, y, z) \, dV = \iint_D \left[\int_{\phi_1(x, y)}^{\phi_2(x, y)} f(x, y, z) \, dz \right] dA.$$

Passing to cylindrical coordinates here we actually have to replace D by its image D^* in polar coordinates and $dz \, dA$ by $r \, dz \, dr \, d\theta$.

EXAMPLE 6. Find the volume of the solid E bounded by the surfaces

$$y = x, \quad y = -x, \quad x^2 + y^2 = 5z, \quad z = 7$$

so that $y \geq 0$.