

## 7. Differences between Linear and Nonlinear Equations (section 2.4)

### 1. Existence and Uniqueness of Solutions

THEOREM 1. *Let the functions  $f$  and  $\partial f/\partial y$  be continuous in some rectangle*

$$R = \{(t, y) | \alpha < t < \beta, \quad \gamma < y < \delta\}$$

*containing the point  $(t_0, y_0)$ . Then in some interval  $t_0 - h < t < t_0 + h$  contained in  $I = \{t | \alpha < t < \beta\}$ , there is a **unique** solution  $y = y(t)$  of the initial value problem*

$$y' = f(t, y), \quad y(t_0) = y_0.$$

2. By this theorem we can guarantee the existence of solution only for values of  $t$  which are sufficiently closed to  $t_0$ , but not for all  $t$ .
3. Geometric consequence of the theorem is that two integral curves never intersect each other.
4. The condition “ $\partial f/\partial y$  be continuous in some rectangle...” is important for uniqueness.  
Illustration: Apply the existence and uniqueness theorem to the following IVP:

$$y' = y^{2/3}, \quad y(0) = 0.$$

### 5. Existence and Uniqueness of Solutions of Linear ODE

**THEOREM 2.** *If the functions  $p(t)$  and  $g(t)$  are continuous on the interval  $I = \{t \mid \alpha < t < \beta\}$ , then for any  $t = t_0$  on  $I$ , there is a **unique** solution  $y = y(t)$  of the initial value problem*

$$y' + p(t)y = g(t), \quad y(t_0) = y_0. \quad (1)$$

6. Note that the conditions of the Theorem 1 hold automatically for linear ODE.

7. Determine (without solving the problem) an interval in which the solution of the given IVP is certain to exist:

$$(t - 3)y' + (\ln |t|)y = 2t$$

(a)  $y(1) = 2$

(b)  $y(5) = 7$

8. Consider

$$ty' + 2y = 4t^2 \quad (2)$$

- (a) Determine (without solving the problem) an interval in which the solution (2) satisfying  $y(t_0) = y_0$  with  $t_0 > 0$  is certain to exist.

- (b) The solution of IVP from item (a) found by the method of integrating factor is given:

$$y(t) = t^2 + \frac{C}{t^2}, \quad C = t_0^2(y_0 - t_0^2).$$

Using that information discuss the domain of the solution and compare your conclusion with the answer of (a).