

10.1: Sequences

A sequence is a list of numbers written in a definite order.

General sequence terms are denoted as follows:


$$\begin{array}{llll}
 n=1 & a_1 & - & \text{first term} \\
 n=2 & a_2 & - & \text{second term} \\
 & \vdots & & \\
 n & a_n & - & n^{\text{th}} \text{ term} \\
 n+1 & a_{n+1} & - & (n+1)^{\text{th}} \text{ term} \\
 & \vdots & &
 \end{array}$$

Notice that, in general, $a_{n+1} \neq a_n + 1$.

A sequence can be defined as a function whose domain is the set of positive ^{whole} numbers:

$$\begin{array}{ccc}
 \mathbb{Z}^+ & \longrightarrow & \mathbb{R} \\
 \{1, 2, 3, \dots\} & \longrightarrow & \{a_1, a_2, a_3, \dots\}
 \end{array}$$

$$n \longrightarrow a_n = f(n)$$



NOTATION: $\{a_1, a_2, \dots, a_n, a_{n+1}, \dots\}$, $\{a_n\}$, $\{a_n\}_{n=1}^{\infty}$.

EXAMPLE 1. Write down the first few terms of the following sequences:

$$(a) \left\{ \frac{n+1}{n^2} \right\}_{n=1}^{\infty} = \left\{ \underset{n=1}{2}, \underset{n=2}{\frac{3}{4}}, \underset{n=3}{\frac{4}{9}}, \underset{n=4}{\frac{5}{16}}, \dots \right\}$$

$$(b) \left\{ \frac{(-1)^{n+1}}{2^n} \right\}_{n=0}^{\infty} = \left\{ -1, \frac{1}{2}, -\frac{1}{4}, \frac{1}{8}, \dots \right\}$$

(c) The Fibonacci sequence $\{f_n\}$ is defined recursively:

$$f_1 = 1, \quad f_2 = 1, \quad f_n = f_{n-1} + f_{n-2}, \quad n \geq 3.$$

$$f_1 = 1$$

$$f_2 = 1$$

$$f_3 = f_1 + f_2 = 1 + 1 = 2$$

$$f_4 = f_2 + f_3 = 1 + 2 = 3$$

$$f_5 = f_4 + f_3 = 3 + 2 = 5$$

\vdots

1, 1, 2, 3, 5, 8, 13, 21, ...



EXAMPLE 2. Find a general formula for the sequence:

$$(a) \left\{ \frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \frac{1}{9}, \frac{1}{11}, \dots \right\} = \left\{ \frac{1}{2n+1} \right\}_{n=1}^{\infty}$$

$$(b) \left\{ \begin{array}{cccc} \frac{1}{4} & \frac{1}{9} & -\frac{1}{16} & \frac{1}{25} & \dots \\ 2^2 & 3^2 & 4^2 & 5^2 & \end{array} \right\} = \left\{ \frac{(-1)^{n+1}}{n^2} \right\}_{n=2}^{\infty}$$

DEFINITION 3. If $\lim_{n \rightarrow \infty} a_n$ exists ^{and finite} then we say that the sequence $\{a_n\}$ converges (or is convergent.) Otherwise, we say the sequence $\{a_n\}$ diverges (or is divergent.)

Graphs of two sequences with $\lim_{n \rightarrow \infty} a_n = L$.



EXAMPLE 4. Determine if $\{a_n\}_{n=1}^{\infty}$ converges or diverges. If converges, find its limit.

(a) $a_n = \frac{n+1}{2n+3}$

$$\lim_{h \rightarrow \infty} \frac{h+1}{2h+3} = \lim_{h \rightarrow \infty} \frac{\frac{h+1}{h}}{\frac{2h+3}{h}} = \lim_{h \rightarrow \infty} \frac{1 + \frac{1}{h}}{2 + \frac{3}{h}} = \frac{1}{2}$$

convergent

(b) $a_n = \frac{3n^2 - 1}{10n + 5n^2}$

$$\lim_{h \rightarrow \infty} \frac{3h^2 - 1}{10h + 5h^2} = \frac{3}{5} \quad \left(\begin{array}{l} \text{AS limit of} \\ \text{rational function} \\ \text{at } \infty \end{array} \right)$$

convergent

(c) $a_n = \arctan(2n)$

$$\lim_{n \rightarrow \infty} \arctan(2n) = \lim_{m \rightarrow \infty} \arctan(m) = \frac{\pi}{2} \quad \text{convergent}$$

(d) $a_n = \ln(2n+4) - \ln n = \ln \frac{2n+4}{n} = \ln \left(2 + \frac{4}{n} \right)$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \ln \left(2 + \frac{4}{n} \right) = \ln \left(\lim_{n \rightarrow \infty} \left(2 + \frac{4}{n} \right) \right)$$

$$= \boxed{\ln 2} \quad \text{convergent}$$

$$(e) a_n = \cos \frac{\pi n}{2}, n \geq 1$$

$$a_1 = \cos \frac{\pi}{2} = 0$$

$$a_2 = \cos \pi = -1$$


$$a_3 = \cos \frac{3\pi}{2} = 0$$

$$a_4 = \cos 2\pi = 1$$

$$a_5 = \cos \frac{5\pi}{2} = 0$$

⋮

$$\{a_n\} = \{0, -1, 0, 1, 0, -1, 0, 1, \dots\}$$



Terms of the sequence oscillate between -1 and 1. Thus, the sequence doesn't approach any number. It is divergent.

Note: $b_n = \cos 2\pi n \Rightarrow b_n = 1$ for all n ,

i.e. $\{b_n\}$ is constant sequence


$\lim_{n \rightarrow \infty} b_n = 1 \Rightarrow \{b_n\}$ is convergent

$$(f) a_n = \frac{3 + (-1)^n}{n^2} = \begin{cases} \frac{4}{n^2} & \text{for even } n \\ \frac{2}{n^2} & \text{for odd } n \end{cases}$$

Squeeze Theorem

$$0 < a_n = \frac{3 + (-1)^n}{n^2} < \frac{4}{n^2} \rightarrow 0 \quad n \rightarrow \infty$$

$$\lim_{n \rightarrow \infty} a_n = 0$$






DEFINITION 5. A sequence $\{a_n\}$ is **bounded above** if there is a number M s.t.

$$a_n \leq M \quad \text{for all } n.$$

A sequence $\{a_n\}$ is **bounded below** if there is a number m s.t.

$$m \leq a_n \quad \text{for all } n.$$

If its bounded above and below, then a_n is a **bounded sequence**.




DEFINITION 6. A sequence $\{a_n\}$ is increasing if

$$a_n < a_{n+1} \quad \text{for all } n.$$

A sequence $\{a_n\}$ is decreasing if

$$a_n > a_{n+1} \quad \text{for all } n.$$

MONOTONIC SEQUENCE THEOREM. *Every bounded, monotonic sequence is convergent.*



EXAMPLE 7. Determine whether a_n is increasing, decreasing or not monotonic.

(a) $a_n = -n^2, n \geq 1$

Way 1

$$n < n+1$$

$$n^2 < (n+1)^2$$

$$-n^2 > -(n+1)^2$$

$\Rightarrow a_n > a_{n+1} \Rightarrow \{a_n\}$ is decreasing

(b) $\left\{ \frac{2}{n^2} \right\}_{n=5}^{\infty}$

$$f(x) = \frac{2}{x^2}, x \geq 5$$

$$f(n) = \frac{2}{n^2}$$

$$f'(x) = \left(\frac{2}{x^2} \right)' = -\frac{4}{x^3} < 0 \Rightarrow f \downarrow \Rightarrow \left\{ \frac{2}{n^2} \right\}_{n=5}^{\infty} \text{ is decreasing.}$$

Way 2

$$a_n = -n^2 = f(n), n \geq 1$$

$$f(x) = -x^2, x \geq 1$$

$$f'(x) = -2x < 0 \text{ for } x \geq 1$$

$f \downarrow \Rightarrow \{a_n\}$ is decreasing


(c) $\{(-1)^{n+1}\}_{n=1}^{\infty} = \{1, -1, 1, -1, 1, -1, \dots\}$ oscillates
not monotonic

(d) $a_n = \frac{\sqrt{n+1}}{5n+3}, n = 0, 1, 2, \dots$

Define $f(x) = \frac{\sqrt{x+1}}{5x+3}$ for all $x \geq 0$ so that $f(n) = a_n$

$$f'(x) = \frac{\frac{1}{2\sqrt{x+1}}(5x+3) - 5\sqrt{x+1}}{(5x+3)^2} = \text{common denom.}$$

$$f'(x) = \frac{5x+3 - 10(x+1)}{2\sqrt{x+1}(5x+3)^2} = - \frac{5x+7}{2\sqrt{x+1}(5x+3)^2} > 0$$



$f'(x) < 0$ for all $x \geq 0$

$\Rightarrow f(n) \downarrow$ as $n \rightarrow \infty$

$\{a_n\}_{n=0}^{\infty}$ is decreasing

Example 8.
Consider the sequence defined by $a_1 = 1$, $a_{n+1} = \frac{1}{3-a_n}$

Find the first 5 terms of this sequence. Find the limit of the sequence.

$$a_1 = 1$$

$$a_2 = \frac{1}{3-a_1} = \frac{1}{2}$$

$$a_3 = \frac{1}{3-\frac{1}{2}} = \frac{2}{5}$$

$$a_4 = \frac{1}{3-\frac{2}{5}} = \frac{5}{13}$$

$$a_5 = \frac{1}{3-\frac{5}{13}} = \frac{13}{34}$$

$$1, \frac{1}{2}, \frac{2}{5}, \frac{5}{13}, \frac{13}{34}, \dots$$

$\{a_n\}$ is bounded

$$0 < a_n < 1$$

decreasing (monotonic)

\Rightarrow convergent

There exist a real number L
Such that

$$L = \lim_{n \rightarrow \infty} a_n$$

Note here
that $0 < L < 1$

Find L .

$$L = \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} a_{n+1} = \lim_{n \rightarrow \infty} \frac{1}{3-a_n} = \frac{1}{3-L}$$

because $a_n \neq 3$

$$L = \frac{1}{3-L} \Rightarrow L(3-L) = 1$$

$$3L - L^2 = 1$$

$$L^2 - 3L + 1 = 0$$

$$L_{1,2} = \frac{3 \pm \sqrt{3^2 - 4 \cdot 1 \cdot 1}}{2}$$

$$L_{1,2} = \frac{3 \pm \sqrt{5}}{2} \quad \text{b/c } 0 < L < 1$$

$$L = \boxed{\frac{3 - \sqrt{5}}{2}}$$