

New Conjectured Reality: Welschinger signs and the Wronski map

Nonlinear Algebra in Applications

SIAM TX-LA Meeting


University of Louisiana, Lafayette, 4 November 2023



Frank Sottile

Texas A&M University
sottile@tamu.edu


With Tom Brazelton
and Stephen McKean



Nodes and Signs of Rational Plane Curves

An irreducible degree d plane curve C has arithmetic genus $\binom{d-1}{2}$.

\Rightarrow when C is rational ($g = 0$), it necessarily has singularities.

If C is also general, it has $\binom{d-1}{2}$ ordinary double points .

Real curves have three types of ordinary double points:



node




solitary
point

Definition: (Welschinger) The sign of a real rational normal curve C is $w(C) := (-1)^{\#\text{solitary points}}$.

Nodes and Signs of Rational Plane Curves

An irreducible degree d plane curve C has arithmetic genus $\binom{d-1}{2}$.

\Rightarrow when C is rational ($g = 0$), it necessarily has singularities.

If C is also general, it has $\binom{d-1}{2}$ ordinary double points 


Real curves have three types of ordinary double points:



node




solitary
point



conjugate pair of nodes

Definition: (Welschinger) The sign of a real rational normal curve C is $w(C) := (-1)^{\#\text{solitary points}}$.

Welschinger's Lower Bound



c. 1990, Kontsevich gave a formula for the number N_d of rational curves through $3d-1$ general points in \mathbb{P}^2 .

Welschinger c. 2002: *If each of the $3d-1$ points are real, then*

$$\sum_{C \text{ real}} w(C)$$

is independent of the choice of $3d-1$ general real points.

IKS: This sum, W_d is at least $\frac{d!}{3}$ and $\lim_{d \rightarrow \infty} \frac{\log W_d}{\log N_d} = 1$.

d	1	2	3	4	5
N_d	1	1	12	620	87304
W_d	1	1	8	240	18264

Parametrized Rational Curves From Grassmannians

The rational normal curve is

$$\gamma: \mathbb{P}^1 \rightarrow \mathbb{P}^d = \mathbb{P}(\mathbb{C}_d[s, t]) = \mathbb{P}(V).$$

A codimension $k+1$ plane H in \mathbb{P}^d ($H \in \mathbb{G}$)

is the centre of a linear projection

$$\pi_H: \mathbb{P}^d - H \rightarrow \mathbb{P}(V/H) =: \mathbb{P}^k.$$


The induced map $\gamma_H: \mathbb{P}^1 \rightarrow \mathbb{P}^k$ is a rational curve of degree d .

Singularities of γ_H correspond to the interaction of H with γ .

E.G. a *flex* (first k derivatives dependent) at $\gamma_H(s)$ corresponds to H meeting the k -plane $F_k(s)$ osculating γ at $s \in \mathbb{P}^1$.

Cusps and other *ramification* of γ_H correspond to Schubert conditions that H satisfies w.r.t osculating flags $F_\bullet(s)$.

(From the 19th c. and used by Eisenbud-Harris in the 1980's.)



Two curves
with $d = 4$

The Wronski Map

Given $H \in \mathbb{G}$, we get the rational curve $\gamma_H = (f_0(s, t), \dots, f_k(s, t))$ (f_i homogeneous of degree d). The *Wronskian* of H is

$$\text{Wr}(H) := \det \left(\frac{\partial^a}{\partial s^a} \frac{\partial^b}{\partial t^b} f_i(s, t) \right)_{\substack{i=0, \dots, k \\ a+b=k}} \in \mathbb{P}(\mathbb{C}_N[s, t]).$$

Here, $N := (k+1)(d-k) = \dim \mathbb{G}$.

Zeros of $\text{Wr}(H) \longleftrightarrow$ flexes of γ_H .

This *Wronski map* $\mathbb{G} \ni H \mapsto \text{Wr}(H)$ is the restriction to \mathbb{G} of a linear projection

$$\text{Wr} : \mathbb{P}(\wedge^{k+1} \mathbb{C}_d[s, t]) \longrightarrow \mathbb{P}(\mathbb{C}_N[s, t]) =: \mathbb{P}^N.$$

Easy fact: This is a finite map $\mathbb{G} \rightarrow \mathbb{P}^N$ of degree


$$\deg \mathbb{G} = \frac{1!2! \cdots (d-k-1)! \cdot N!}{k!(k+1)! \cdots (d-1)!}.$$

Maximally Inflected Curves

Theorem. (MTV) If $f \in \mathbb{P}^N$ has all roots real, then $Wr^{-1}(f) \subset \mathbb{G}_{\mathbb{R}}$.
If f also has distinct roots, it is a regular value of Wr .

(If $Wr(H)$ has all roots real, then H is necessarily real.)

Definition. (Kharlamov-S.) If $Wr(H)$ has all roots real, then γ_H is *maximally inflected* in that all flexes occur at real points.




Maximally inflected



Not maximally inflected

These curves are beautiful; here are a few quintics.



New Conjectured Reality

Restricting Wr to the big cell of $\mathbb{G}_{\mathbb{R}}$ gives a proper map $Wr: \mathbb{R}^N \rightarrow \mathbb{R}^N$ (=monic real polynomials of degree N).

Eremenko and Gabrielov computed its degree for all k and d (formula omitted).

Curious Conjecture: (Brazelton-S.) Fix $k = 2$. When γ_H is maximally inflected with only flexes (e.g. $Wr(H)$ has (all) N real roots), then $\deg_H Wr = (-1)^d w(\gamma_H)$:

Sign of the Wronski map at $H =$ Welschinger sign of curve γ_H .

This is easily proven when $d = 4$, and there is significant evidence for $d = 5, 6$. Computations, even for $d = 6$ are challenging.

Obvious generalizations do not appear to hold.

Even More Reality, Experimentally

When $k = 2$, $d = 5$, and f has all $N = 9$ roots real, then $\#Wr^{-1}(f) = 42$.

Define $S(j)_f := \#\{H \in Wr^{-1}(f) \mid \gamma_H \text{ has } j \text{ solitary points}\}$.

In each of $\gtrsim 10^6$ examples, we find that

$$S_5 = (S(j)_f \mid j = 0 \dots 6) = (0, 0, 0, 12, 18, 9, 3).$$

When $d = 4$, we have $\#Wr^{-1}(f) = 5$, and it is a result of Kharlamov-S. that $S_4 = (0, 0, 3, 2)$ for any f .

$d = 6$, we have $\#Wr^{-1}(f) = 462$, and in each of about 200 challenging examples, we find that

$$S_6 = (0, 0, 0, 0, 55, 132, 132, 88, 39, 12, 4).$$


Another Reality Conjecture

Other ramifications may be imposed on plane curves: E.g. in a local parameter, the curve is $s \mapsto (s^{1+b}, s^{2+a})$ with $a \geq b \geq 0$. (A simple flex is $(a, b) = (1, 0)$.)







This ramification has order $a + b$, and the sum of all local ramifications of a curve is $N = 3(d - 2)$.

Assigning ramifications to points of $\mathbb{RP}^1 \simeq S^1$ gives a necklace, e.g. $(\text{cusp}, \text{flex}, \text{cusp}, \text{flex}, \text{cusp}, \text{flex}) \neq (\text{flex}, \text{flex}, \text{flex}, \text{cusp}, \text{cusp}, \text{cusp})$.

For example, there are three necklaces with three cusps (κ) and three flexes (ι):



Some Pictorial Data

		
		
<p>(1,3,2,0)</p>	<p>(2,3,1,0)</p>	<p>(5,0,0,1)</p>

Another Reality Conjecture

Conjecture. For a given necklace ν of ramification, the vector
(#curves with given ramification and i solitary points | i)
is independent of the placement of the points of ramification.

This has been tested thousands of times for all ramification when $d = 5$ and many times for $d = 6$.