

## Math 409 midterm 2 practice #2

Name: \_\_\_\_\_

This exam has 4 questions, for a total of 100 points.

Please answer each question in the space provided. No aids are permitted.

### Question 1. (40 pts)

In each of the following eight cases, indicate whether the given statement is true or false. No justification is necessary.

- (a) Every subsequence of a Cauchy sequence is also a Cauchy sequence.

**Solution:** True.

- (b) If  $f$  is a continuous function on a closed and bounded interval  $I$ , then there exists  $x_0 \in I$  such that  $f(x_0)$  is the maximum of  $f$  on  $I$ .

**Solution:** True.

- (c)  $\lim_{x \rightarrow 0} \frac{x^3 \sin(1/x) + x}{x \cos x}$  exists.

**Solution:** True.

- (d) A bounded sequence  $\{x_n\}$  in  $\mathbb{R}$  can have two subsequences converging to two different numbers.

**Solution:** True.

- (e) If  $g(x) \leq -1$  for all  $x \in \mathbb{R}$  and  $\lim_{x \rightarrow a} f(x) = 0$ , then  $\lim_{x \rightarrow a} \frac{g(x)}{f(x)} = -\infty$ .

**Solution:** False.

- (f) A bounded increasing sequence converges to a finite number.

**Solution:** True.

- (g) If  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a uniformly continuous function, then  $f$  is bounded on  $\mathbb{R}$ .

**Solution:** False.

- (h) Given a sequence  $\{x_n\}$  with  $x_n > 0$  for all  $n$ , if  $\{x_n\}$  has no converging subsequences, then  $x_n \rightarrow \infty$ , as  $n \rightarrow \infty$ .

**Solution:** True.

**Question 2. (20 pts)**

- (a) State the definition of Cauchy sequences.

**Solution:** Omitted. You can find it in the textbook.

- (b) Let  $\{x_n\}$  be a real sequence such that

$$x_{n+1} = x_n + \left(\frac{1}{3}\right)^n.$$

Prove that  $\{x_n\}$  is a Cauchy sequence.

**Solution:** For any  $n > m$ , we have

$$\begin{aligned} |x_n - x_m| &= |x_n - x_{n-1} + x_{n-1} - x_{n-2} + \cdots - x_{m+1} + x_{m+1} - x_m| \\ &\leq |x_n - x_{n-1}| + |x_{n-1} - x_{n-2}| + \cdots + |x_{m+1} - x_m| \\ &= \left(\frac{1}{3}\right)^{n-1} + \left(\frac{1}{3}\right)^{n-2} + \cdots + \left(\frac{1}{3}\right)^m \\ &= \frac{(1/3)^m(1 - (1/3)^{n-m})}{1 - (1/3)} \leq (3/2) \cdot (1/3)^m \end{aligned}$$

So for any  $\varepsilon > 0$ , choose  $N \in \mathbb{N}$  such that  $(3/2) \cdot (1/3)^N < \varepsilon$ . Then for any  $n, m > N$ , we have

$$|x_n - x_m| < (3/2) \cdot (1/3)^N < \varepsilon.$$

So  $\{x_n\}$  is Cauchy.

**Question 3. (20 pts)**

- (a) State the Intermediate Value Theorem.

**Solution:** Omitted. You can find it in the textbook.

- (b) Prove that there exists an  $x \in \mathbb{R}$  such that  $4^x = x^3 + \sin x + x^2 + 2$ .

**Solution:** Consider the function  $g(x) = 4^x - (x^3 + \sin x + x^2 + 2)$ . On the interval  $[0, 3]$ , we have

$$g(0) = -1 \text{ and } g(3) = 64 - (27 + \sin 3 + 9 + 2) > 0.$$

So we have  $g(0) \leq 0 \leq g(3)$ . It follows from the intermediate value theorem that there exists  $x_0 \in [0, 3]$  such that  $g(x_0) = 0$ .

**Question 4. (20 pts)**

- (a) State the definition of uniform continuity.

**Solution:** Omitted. You can find it in the textbook.

- (b) Let  $f$  and  $g$  be uniformly continuous functions on  $\mathbb{R}$ . Prove that  $f + g$  is uniformly continuous on  $\mathbb{R}$ .

**Solution:** Since  $f$  is uniformly continuous on  $\mathbb{R}$ , for any  $\varepsilon > 0$ , there exists  $\delta_1$  such that

$$|f(x) - f(y)| < \varepsilon/2$$

for all  $x, y \in \mathbb{R}$  with  $|x - y| < \delta_1$ . Similarly, there exists  $\delta_2$  such that

$$|g(x) - g(y)| < \varepsilon/2$$

for all  $x, y \in \mathbb{R}$  with  $|x - y| < \delta_2$ . Let  $\delta = \min\{\delta_1, \delta_2\}$ . Then we have

$$|(f + g)(x) - (f + g)(y)| \leq |f(x) - f(y)| + |g(x) - g(y)| < \varepsilon$$

for all  $x, y \in \mathbb{R}$  with  $|x - y| < \delta$ .

- (c) Let  $f$  and  $g$  be uniformly continuous functions on  $\mathbb{R}$ . If both  $f$  and  $g$  are bounded on  $\mathbb{R}$ , then  $fg$  is also uniformly continuous on  $\mathbb{R}$ .

**Solution:** Since  $f$  and  $g$  are bounded over  $\mathbb{R}$ , there exists  $M > 0$  such that

$$|f(x)| < M \text{ and } |g(x)| < M$$

for all  $x \in \mathbb{R}$ .

Since  $f$  is uniformly continuous on  $\mathbb{R}$ , for any  $\varepsilon > 0$ , there exists  $\delta_1$  such that

$$|f(x) - f(y)| < \varepsilon/2M$$

for all  $x, y \in \mathbb{R}$  with  $|x - y| < \delta_1$ . Similarly, there exists  $\delta_2$  such that

$$|g(x) - g(y)| < \varepsilon/2M$$

for all  $x, y \in \mathbb{R}$  with  $|x - y| < \delta_2$ .

Let  $\delta = \min\{\delta_1, \delta_2\}$ . Then for all  $x, y \in \mathbb{R}$  with  $|x - y| < \delta$ , we have

$$\begin{aligned} |f(x)g(x) - f(y)g(y)| &= |f(x)g(x) - f(x)g(y) + f(x)g(y) - f(y)g(y)| \\ &\leq |f(x)g(x) - f(x)g(y)| + |f(x)g(y) - f(y)g(y)| \\ &= |f(x)||g(x) - g(y)| + |g(y)||f(x) - f(y)| < \varepsilon. \end{aligned}$$