

Name _____

MATH 221 Final Exam Version A Fall 2019

Section 504 P. Yasskin

Multiple Choice: (5 points each. No part credit.)

1-11	/55	13	/10
12	/15	14	/25
		Total	/105

1. A triangle has vertices $A = (3,2,1)$, $B = (3,3,2)$ and $C = (4,4,2)$. Find the angle at A .

- a. 90°
- b. 60°
- c. 45°
- d. 30°
- e. 0°

2. Find the area of the triangle with vertices $A = (3,2,1)$, $B = (3,3,2)$ and $C = (4,4,2)$.

- a. $\frac{\sqrt{3}}{2}$
- b. $\sqrt{3}$
- c. $\frac{\sqrt{6}}{2}$
- d. $\sqrt{6}$
- e. $2\sqrt{6}$

3. Find the arc length of the curve $\vec{r}(t) = (t^2, 2t, \ln t)$ between $(1, 2, 0)$ and $(e^2, 2e, 1)$.

- a. $e^2 + 1$
- b. $e^2 - 1$
- c. e^2
- d. $\sqrt{e^2 + 1}$
- e. $\sqrt{e^2 + 1} - 1$

4. Find the average value of the function $f(x, y, z) = x$ along the curve $\vec{r}(t) = (t^2, 2t, \ln t)$ between $(1, 2, 0)$ and $(e^2, 2e, 1)$.

- a. $\frac{e^2}{2} + \frac{1}{2} - \frac{1}{e^2}$
- b. $\frac{e^4}{2} + \frac{e^2}{2} - 1$
- c. $\frac{e^2}{2} - \frac{1}{2} - \frac{1}{e^2}$
- d. $\frac{e^4}{2} - \frac{e^2}{2} - 1$
- e. $\frac{e^2}{2} - \frac{1}{e^2}$

5. Find the plane tangent to $x \sin y + 2z \cos y = 2$ at $(x, y, z) = \left(\sqrt{2}, \frac{\pi}{4}, \frac{1}{\sqrt{2}}\right)$. The z -intercept is:

- a. 0
- b. $\frac{1}{2\sqrt{2}}$
- c. $\frac{1}{\sqrt{2}}$
- d. $\sqrt{2}$
- e. 2

6. The velocity field of the water in a sink is $\vec{V} = \langle -x^2y, xy^2 \rangle$. Find the circulation of the water, $Circ = \oint \vec{V} \cdot d\vec{s}$, counterclockwise around the circle $x^2 + y^2 = 4$.

HINT: Use Green's Theorem.

- a. 64π
- b. 32π
- c. 8π
- d. $\frac{16}{3}\pi$
- e. 2π

7. Compute $\int \vec{F} \cdot d\vec{s}$ for $\vec{F} = \langle 4x^3, 3y^2, 2z \rangle$ along the curve $\vec{r}(t) = \left(t \sin\left(\frac{\pi}{2}t\right), 2^t \cos\left(\frac{\pi}{2}t\right), 2^t \sin\left(\frac{\pi}{2}t\right) \right)$ from $t = 0$ to $t = 1$.

HINT: Find a scalar potential.

- a. 4
- b. 3
- c. 2
- d. 1
- e. 0

8. Compute $\iint_S \vec{\nabla} \times \vec{G} \cdot d\vec{S}$ for $\vec{G} = \langle y, -x, z \rangle$ over the surface $\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, 16 - r^4)$ for $0 \leq r \leq 2$ and $0 \leq \theta \leq 2\pi$ oriented up and out.

HINT: Use a theorem.

- a. 4π
- b. 2π
- c. -2π
- d. -4π
- e. -8π

9. Find the centroid of the solid hemisphere $0 \leq z \leq \sqrt{4 - x^2 - y^2}$.

- a. $(0, 0, 2\pi)$
- b. $(0, 0, 4\pi)$
- c. $(0, 0, \frac{3}{4})$
- d. $(0, 0, \frac{4}{3})$
- e. $(0, 0, \frac{1}{2})$

10. Find the area of the surface $\vec{R}(u, v) = (u, v, u + v)$ for $0 \leq u \leq 2$ and $0 \leq v \leq 3$.

- a. $3\sqrt{2}$
- b. $6\sqrt{3}$
- c. $6\sqrt{2}$
- d. $12\sqrt{3}$
- e. $12\sqrt{2}$

11. Find the flux of $\vec{F} = \langle y, -z, x \rangle$ upward thru the surface $\vec{R}(u, v) = (u, v, u + v)$ for $0 \leq u \leq 2$ and $0 \leq v \leq 3$.

- a. 0
- b. 12
- c. 18
- d. 24
- e. 30

Work Out: (Points indicated. Part credit possible. Show all work.)

12. (15 points) The Ideal Gas Law says the Pressure, P , Density, δ , and Temperature, T , are related by $P = k\delta T$ for some constant k . We will assume the atmosphere is an ideal gas with $k = 2$. A weather balloon measures the Density and Temperature to be

$$\delta = 0.1 \frac{\text{gm}}{\text{m}^3} \quad T = 270^\circ\text{K}$$

and their gradients to be

$$\vec{\nabla}\delta = \langle 0.03, 0.02, 0.01 \rangle \quad \vec{\nabla}T = \langle -1, 1, 3 \rangle$$


Find the gradient of the Pressure.

13. (10 points) Find the largest value of $f = xy^2z^3$ on the plane $6x + 3y + 2z = 72$.

14. (25 points) Verify Gauss' Theorem $\iiint_V \vec{\nabla} \cdot \vec{F} dV = \iint_{\partial V} \vec{F} \cdot d\vec{S}$

for the vector field $\vec{F} = (xz, yz, 3z\sqrt{x^2 + y^2})$ and the solid cone $\sqrt{x^2 + y^2} \leq z \leq 2$.

Be careful with orientations. Use the following steps:



First the Left Hand Side:

a. Compute the divergence:

$$\vec{\nabla} \cdot \vec{F} =$$

b. Express the divergence and the volume element in the appropriate coordinate system:

$$\vec{\nabla} \cdot \vec{F} = \qquad dV =$$

c. Compute the left hand side:

$$\iiint_V \vec{\nabla} \cdot \vec{F} dV =$$

Second the Right Hand Side:

The boundary surface consists of a hemisphere H and a disk D with appropriate orientations.

The disk D may be parametrized as: $\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, 2)$

d. Compute the tangent vectors:

$$\vec{e}_r =$$

$$\vec{e}_\theta =$$

e. Compute the normal vector:

$$\vec{N} =$$

f. Evaluate $\vec{F} = (xz, yz, 3z\sqrt{x^2 + y^2})$ on the disk:

$$\vec{F}|_{\vec{R}(r,\theta)} =$$

g. Compute the dot product:

$$\vec{F} \cdot \vec{N} =$$

h. Compute the flux through D :

$$\iint_D \vec{F} \cdot d\vec{S} =$$

The cone C may be parametrized as: $\vec{R}(r, \theta) = (r \cos \theta, r \sin \theta, r)$

i. Compute the tangent vectors:

$$\vec{e}_r =$$

$$\vec{e}_\theta =$$

j. Compute the normal vector:

$$\vec{N} =$$

k. Evaluate $\vec{F} = (xz, yz, 3z\sqrt{x^2 + y^2})$ on the cone:

$$\vec{F}|_{\vec{R}(r,\theta)} =$$

l. Compute the dot product:

$$\vec{F} \cdot \vec{N} =$$

m. Compute the flux through C :

$$\iint_C \vec{F} \cdot d\vec{S} =$$

n. Compute the **TOTAL** right hand side: