

Name _____

MATH 251 Exam 1 Version A Fall 2018
Sections 504/505 Solutions P. Yasskin

1-9	/54	11	/16
10	/36	Total	/106

Multiple Choice: (6 points each. No part credit.)

1. The Galactic Federation is trying to keep a stasis pod stationary in intergalactic space where there is no gravity. They already have 2 tractor beams pulling on the pod with the forces

$$\vec{F}_1 = \langle 4, 1, -3 \rangle \quad \text{and} \quad \vec{F}_2 = \langle -2, 2, 1 \rangle$$

If they apply a 3rd tractor beam on the pod, what should its force \vec{F}_3 be to keep the pod stationary?

- a. $\vec{F}_3 = \langle -2, -3, 2 \rangle$ Correct Choice
- b. $\vec{F}_3 = \langle -2, 3, 2 \rangle$
- c. $\vec{F}_3 = \langle 2, 3, -2 \rangle$
- d. $\vec{F}_3 = \langle 2, -3, -2 \rangle$
- e. $\vec{F}_3 = \langle 2, 3, 2 \rangle$

Solution: To balance forces, $\vec{F}_1 + \vec{F}_2 + \vec{F}_3 = \vec{0}$. So

$$\vec{F}_3 = -\vec{F}_1 - \vec{F}_2 = -\langle 4, 1, -3 \rangle - \langle -2, 2, 1 \rangle = \langle -2, -3, 2 \rangle$$

2. The Galactic Federation moves a stasis pod from $(2, 3, 4)$ to $(6, 9, 0)$ by applying the 2 forces:

$$\vec{F}_1 = \langle 4, 1, -3 \rangle \quad \text{and} \quad \vec{F}_2 = \langle -2, 2, 1 \rangle$$

How much work is done by the force \vec{F}_1 only?

- a. $W = 0$
- b. $W = 10$
- c. $W = 22$
- d. $W = 33$
- e. $W = 34$ Correct Choice

Solution: The displacement is $\vec{D} = (6, 9, 0) - (2, 3, 4) = \langle 4, 6, -4 \rangle$. So the work is

$$W = \vec{F}_1 \cdot \vec{D} = 16 + 6 + 12 = 34$$

3. If \vec{u} points Up and \vec{v} points SouthEast, where does $\vec{u} \times \vec{v}$ point?

- a. Down
- b. NorthEast Correct Choice
- c. SouthWest
- d. NorthWest
- e. South

Solution: Hold your right hand with the fingers pointing Up and the palm facing SouthEast. Then the thumb points NorthEast.

4. Convert the polar equation $r = 4 \cos \theta$ to rectangular coordinates and identify the shape of the curve.
- Circle of radius 4 centered at a point on the x -axis.
 - Circle of radius 4 centered at a point on the y -axis.
 - Circle of radius 2 centered at a point on the x -axis. **Correct Choice**
 - Circle of radius 2 centered at a point on the y -axis.
 - Parabola opening to the right.

Solution: $r = 4 \cos \theta = 4 \frac{x}{r} \Rightarrow r^2 = 4x \Rightarrow x^2 + y^2 - 4x = 0 \Rightarrow (x - 2)^2 + y^2 = 4$
 This is a circle of radius 2 centered at (2,0).

5. Find the angle between the direction of the line $(x, y, z) = (3 + 3t, 3 - 3t, 4)$ and the normal to the plane $2x + 2z = 15$.
- 0°
 - 30°
 - 45°
 - 60° **Correct Choice**
 - 90°

Solution: The tangent to the line is $\vec{v} = \langle 3, -3, 0 \rangle$. The normal to the plane is $\vec{N} = \langle 2, 0, 2 \rangle$.

So $\cos \theta = \frac{\vec{v} \cdot \vec{N}}{|\vec{v}| |\vec{N}|} = \frac{6}{\sqrt{18} \sqrt{8}} = \frac{6}{3\sqrt{2} \cdot 2\sqrt{2}} = \frac{1}{2}$ and $\theta = 60^\circ$

6. Find the point where the line $(x, y, z) = \vec{r}(t) = (t + 2, t - 2, 2t - 1)$ intersects the plane $3x - y + 2z = 12$. At this point $x + y + z =$
- 3
 - 1
 - 0
 - 1
 - 3 **Correct Choice**

Solution: Plug the line into the plane and solve for t :

$$3x - y + 2z = 3(t + 2) - (t - 2) + 2(2t - 1) = 6t + 6 = 12 \Rightarrow t = 1$$

So the point is $(x, y, z) = \vec{r}(1) = (3, -1, 1)$ and so $x + y + z = 3$.

7. The graph of the equation $x^2 + y^2 - z = -1$ is a
- Hyperboloid of 1 sheet
 - Hyperboloid of 2 sheets
 - Cone
 - Elliptic Paraboloid opening up **Correct Choice**
 - Elliptic Paraboloid opening down

Solution: We write it as $z = 1 + x^2 + y^2$.

It is an Elliptic Paraboloid because z is to the first power and x^2 and y^2 have the same sign. It opens up because $z \geq 1$.

8. Find the equation of the plane thru the point $P = \langle 1, 2, 3 \rangle$ tangent to the vectors $\vec{a} = \langle 1, 2, 3 \rangle$ and $\vec{b} = \langle 3, 2, 1 \rangle$.
- $-4x + 8y - 4z = 0$ **Correct Choice**
 - $-4x - 8y - 4z = 0$
 - $-4x + 8y - 4z = 32$
 - $-4x - 8y - 4z = 32$
 - $-4x - 8y - 4z = 16$

Solution: $\vec{N} = \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 3 \\ 3 & 2 & 1 \end{vmatrix} = \hat{i}(2 - 6) - \hat{j}(1 - 9) + \hat{k}(2 - 6) = (-4, 8, -4)$

$$\vec{N} \cdot X = \vec{N} \cdot P \quad -4x + 8y - 4z = -4(1) + 8(2) - 4(3) = 0$$

9. Find the area of the triangle with adjacent edges $\vec{a} = \langle 1, 2, 3 \rangle$ and $\vec{b} = \langle 3, 2, 1 \rangle$.
- $\frac{1}{2}\sqrt{6}$
 - $\sqrt{6}$
 - $2\sqrt{6}$ **Correct Choice**
 - $4\sqrt{6}$
 - $2\sqrt{2}$

Solution: $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 3 \\ 3 & 2 & 1 \end{vmatrix} = \hat{i}(2 - 6) - \hat{j}(1 - 9) + \hat{k}(2 - 6) = (-4, 8, -4)$

$$A = \frac{1}{2} |\vec{a} \times \vec{b}| = \frac{1}{2} \sqrt{16 + 64 + 16} = \frac{1}{2} \sqrt{96} = 2\sqrt{6}$$

Work Out: (Points indicated. Part credit possible. Show all work.)

10. (36 points) For the twisted cubic $\vec{r}(t) = \left(\frac{t^3}{3}, 2t, t^2\right)$ compute each of the following:

a. (6 pts) The velocity \vec{v}

Solution:

$$\vec{v} = \underline{\langle t^2, 2, 2t \rangle}$$

b. (6 pts) The speed $\frac{ds}{dt}$ (Simplify!)

Solution: $\frac{ds}{dt} = |\vec{v}| = \sqrt{t^4 + 4 + 4t^2} = \sqrt{(t^2 + 2)^2} = t^2 + 2$

$$\frac{ds}{dt} = \underline{t^2 + 2}$$

c. (6 pts) The tangential acceleration a_T

Solution: $a_T = \frac{d|\vec{v}|}{dt} = \frac{d}{dt}(t^2 + 2) = 2t$

$$a_T = \underline{2t}$$

d. (6 pts) The mass of a wire in the shape of this twisted cubic between $(0,0,0)$ and $(9,6,9)$ if the linear mass density is $\delta = yz$.

Solution: $|\vec{v}| = t^2 + 2$ $\delta = yz = 2t \cdot t^2 = 2t^3$ $(0,0,0) = \vec{r}(0)$ $(9,6,9) = \vec{r}(3)$

$$M = \int_{(0,0,0)}^{(9,6,9)} \delta ds = \int_0^3 yz |\vec{v}| dt = \int_0^3 2t^3 (t^2 + 2) dt = \left[\frac{t^6}{3} + t^4 \right]_0^3 = 3^5 + 3^4 = 243 + 81 = 324$$

$$M = \underline{324}$$

e. (6 pts) The z -component of the center of mass of the wire between $(0,0,0)$ and $(9,6,9)$ if the linear mass density is $\delta = yz$.

Solution: $M_z = \int_{(0,0,0)}^{(9,6,9)} z \delta ds = \int_0^3 yz |\vec{v}| dt = \int_0^3 t^2 2t^3 (t^2 + 2) dt = \left[\frac{t^8}{4} + \frac{4t^6}{6} \right]_0^3$

$$= \frac{3^8}{4} + 2 \cdot 3^5 = \frac{3^5}{4} (3^3 + 8) = \frac{3^5 \cdot 35}{4}$$

$$\bar{z} = \frac{M_z}{M} = \frac{3^5 \cdot 35}{4 \cdot 324} = \frac{105}{16}$$

$$\bar{z} = \underline{\frac{105}{16}}$$

f. (6 pts) The work done to move a bead along of a wire in the shape of this twisted cubic between $(0,0,0)$ and $(9,6,9)$ by the force $\vec{F} = (2y, 3x, z)$.

Solution: $\vec{F}(\vec{r}(t)) = (2y, 3x, z) = (4t, t^3, t^2)$ $\vec{v} = (t^2, 2, 2t)$

$$\vec{F} \cdot \vec{v} = 4t^3 + 2t^3 + 2t^3 = 8t^3$$

$$W = \int_{(0,0,0)}^{(9,6,9)} \vec{F} \cdot d\vec{S} = \int_0^3 \vec{F} \cdot \vec{v} dt = \int_0^3 8t^3 dt = \left[2t^4 \right]_0^3 = 2 \cdot 81 = 162$$

$$W = \underline{162}$$

11. (15 points) Write the vector $\vec{a} = \langle 6, 2, 2 \rangle$ as the sum of two vectors \vec{b} and \vec{c} with \vec{b} parallel to $\vec{d} = \langle 2, 1, -1 \rangle$ and \vec{c} perpendicular to \vec{d} .
Check \vec{c} is perpendicular to \vec{d} .

Solution: $\vec{b} = \text{proj}_{\vec{d}} \vec{a} = \frac{\vec{a} \cdot \vec{d}}{|\vec{d}|^2} \vec{d} = \frac{12 + 2 - 2}{4 + 1 + 1} \langle 2, 1, -1 \rangle = 2 \langle 2, 1, -1 \rangle = \langle 4, 2, -2 \rangle$

$$\vec{c} = \vec{a} - \vec{b} = \langle 6, 2, 2 \rangle - \langle 4, 2, -2 \rangle = \langle 2, 0, 4 \rangle$$

Check: $\vec{d} \cdot \vec{c} = \langle 2, 1, -1 \rangle \cdot \langle 2, 0, 4 \rangle = 4 - 4 = 0$

$$\vec{a} = \frac{\langle 4, 2, -2 \rangle}{\vec{b}} + \frac{\langle 2, 0, 4 \rangle}{\vec{c}}$$