

3. Compute $\iint_S \vec{\nabla} \times \vec{F} \cdot d\vec{S}$ over the quartic surface $z = x^4 + 2x^2y^2 + y^4$ with $z \leq 16$ oriented **down** and **out**, for $\vec{F} = \langle -yz, xz, z^2 \rangle$.

HINT: Use a theorem.

- a. 128π
- b. 256π
- c. 512π
- d. 1024π
- e. 2048π



4. The two legs of a right triangle are \vec{a} and \vec{b} and the hypotenuse is \vec{c} . So $\vec{a} \perp \vec{b}$ and $\vec{c} = \vec{a} + \vec{b}$. Given that $\vec{c} = \langle 9, 9, -9 \rangle$ and the direction of \vec{a} is $\hat{a} = \langle \frac{2}{3}, \frac{-1}{3}, \frac{-2}{3} \rangle$, find the magnitude $|\vec{b}|$.

- a. $|\vec{b}| = 36$
- b. $|\vec{b}| = 18\sqrt{2}$
- c. $|\vec{b}| = 18$
- d. $|\vec{b}| = 9\sqrt{2}$
- e. $|\vec{b}| = 9$

5. An ant is walking across a frying pan where the temperature is $T = \frac{1}{12}x^3y^2$. If the ant is currently at $P = (2, 3)$, in what unit vector direction should the ant walk to reduce the temperature as fast as possible?

- a. $\langle \frac{9}{13}, \frac{4}{13} \rangle$
- b. $\langle \frac{-9}{5}, \frac{-4}{5} \rangle$
- c. $\langle \frac{9}{5}, \frac{4}{5} \rangle$
- d. $\langle \frac{9}{\sqrt{97}}, \frac{4}{\sqrt{97}} \rangle$
- e. $\langle \frac{-9}{\sqrt{97}}, \frac{-4}{\sqrt{97}} \rangle$

6. The point $(1,2)$ is a critical point of the function $f(x,y) = 16x^4 + y^4 - 32xy$.
Classify the point $(1,2)$ using the Second Derivative Test.
- Local Minimum
 - Local Maximum
 - Saddle Point
 - Inflection Point
 - Test Fails

7. Find the mass of the piece of the solid paraboloid $z = x^2 + y^2$
for $2 \leq z \leq 4$ if the density is $\delta = z$.



- 64π
 - 60π
 - $\frac{112}{3}\pi$
 - $\frac{56}{3}\pi$
 - 20π
8. Find the center of mass of the piece of the solid paraboloid $z = x^2 + y^2$
for $2 \leq z \leq 4$ if the density is $\delta = z$.
- $\frac{14}{75}$
 - $\frac{14}{15}$
 - $\frac{45}{14}$
 - $\frac{14}{45}$
 - $\frac{15}{14}$

9. Find the equation of the plane tangent to the hyperboloid $xyz = 6$ at the point $(3, 2, 1)$.

a. $(x, y, z) = (3 + 2t, 2 + 3t, 1 + 6t)$

b. $(x, y, z) = (2 + 3t, 3 + 2t, 6 + t)$

c. $3x + 2y + z = 14$

d. $3x + 2y + z = 18$

e. $2x + 3y + 6z = 18$

10. Find the volume under the surface $z = 2xy$ above the region bounded by $y = x$ and $y = 2\sqrt{x}$.

The base is shown at the right.


a. $\frac{128}{3}$

b. $\frac{128}{5}$

c. $\frac{64}{3}$

d. $\frac{64}{5}$

e. $\frac{64}{7}$



11. Compute
$$\begin{vmatrix} 0 & 0 & 0 & 2 & 0 & 0 \\ 3 & 0 & 0 & 0 & 5 & 0 \\ 0 & 0 & 3 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 4 \\ 0 & 5 & 0 & 0 & 0 & 0 \end{vmatrix}$$


- a. $2!$
- b. $3!$
- c. $4!$
- d. $5!$
- e. $6!$

12. In \mathbb{R}^2 , compute $\oint_C \vec{F} \cdot d\vec{n}$, the flux of $\vec{F} = \langle x^3, y^3 \rangle$ outward thru the circle $x^2 + y^2 = 4$.
HINT: Use a Theorem.




- a. 12π
- b. 16π
- c. 24π
- d. 36π
- e. 48π

Work Out: (Points indicated. Part credit possible. Show all work.)

13. (5 points) At the right is the contour plot of a function $f(x,y)$. The contours are labeled by the function values. If you start at the cross at $(-6,4)$ and move so that your velocity is always in the direction of $\vec{\nabla}f$, the gradient of f , roughly sketch your path on the plot.



14. (20 points) A 118 cm wire is cut into 3 pieces. As shown in the plots, one piece is bent into a square of side a . Another piece is bent into a rectangle with sides b and $2b$. The third piece is bent into a rectangle with sides c and $4c$. Note: a , b and c may not be the same length, even if they look that way in the plots. Find a , b and c which minimize the total area enclosed in the three shapes. Note: the constraint is the sum of the perimeters. You do NOT need to check it is a minimum rather than a maximum.



15. (12 points) Determine whether or not each of these limits exists. If it exists, find its value.

a. $\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2y^2}{x^6 + 3y^3}$


b. $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^2}$

16. (25 points) Verify Gauss' Theorem $\iiint_V \vec{\nabla} \cdot \vec{F} dV = \iint_{\partial V} \vec{F} \cdot d\vec{S}$

for the vector field $\vec{F} = \langle -xz^2, -yz^2, z^3 \rangle$ and the solid above the cone $z = \sqrt{x^2 + y^2}$ below the plane $z = 2$.

Be careful with orientations. Use the following steps:

First the Left Hand Side:



a. Compute the divergence and give the volume element in the appropriate coordinate system:

$\vec{\nabla} \cdot \vec{F} =$ $dV =$

b. Compute the left hand side:

$\iiint_V \vec{\nabla} \cdot \vec{F} dV =$

Second the Right Hand Side:

The boundary surface consists of the cone C and a disk D with appropriate orientations.

c. Parametrize the disk D :

$\vec{R}(r, \theta) =$

d. Compute the tangent vectors:

$\vec{e}_r =$

$\vec{e}_\theta =$

e. Compute the normal vector:

$\vec{N} =$

f. Evaluate $\vec{F} = \langle -xz^2, -yz^2, z^3 \rangle$ on the disk:

$\vec{F}|_{\vec{R}(r, \theta)} =$

g. Compute the dot product:

$$\vec{F} \cdot \vec{N} =$$

h. Compute the flux through D :

$$\iint_D \vec{F} \cdot d\vec{S} =$$

.....
Parametrize the cone C as $\vec{R}(r, \theta) = \langle r \cos \theta, r \sin \theta, r \rangle$

i. Compute the tangent vectors:

$$\vec{e}_r =$$

$$\vec{e}_\theta =$$

j. Compute the normal vector:

$$\vec{N} =$$

k. Evaluate $\vec{F} = \langle -xz^2, -yz^2, z^3 \rangle$ on the cone:

$$\vec{F}|_{\vec{R}(r, \theta)} =$$

l. Compute the dot product

$$\vec{F} \cdot \vec{N} =$$

m. Compute the flux through C :

$$\iint_C \vec{F} \cdot d\vec{S} =$$

.....
n. Compute the **TOTAL** right hand side:

$$\iint_{\partial V} \vec{F} \cdot d\vec{S} =$$