

Name \_\_\_\_\_

MATH 251      Exam 3 Version B      Fall 2020

Sections 519      P. Yasskin

Multiple Choice: (6 points each. No part credit.)

1-8	/48	10	/20
9	/20	11	/20
		Total	/108

1. Compute  $\int_0^2 \int_{x^2}^{2x} 2xy \, dy \, dx$ .

a.  $-\frac{16}{5}$

b.  $\frac{32}{5}$

c.  $\frac{16}{5}$

d.  $\frac{32}{3}$

e.  $\frac{16}{3}$

2. Find the area of the heart shaped region inside the polar curve  $r = |\theta|$ .  
HINT: Double the upper half.


a.  $\frac{\pi^3}{6}$

b.  $\frac{\pi^3}{3}$

c.  $\frac{4\pi^3}{3}$


d.  $\frac{8\pi^3}{3}$

e.  $\frac{16\pi^3}{3}$



3. Find the mass of a triangular plate with vertices  $(0,0)$ ,  $(3,6)$  and  $(3,-6)$  whose surface mass density is  $\delta = x$ .

- a. 12
- b. 24
- c. 36
- d. 48
- e. 60



4. Find the center of mass of a triangular plate with vertices  $(0,0)$ ,  $(3,6)$  and  $(3,-6)$  whose surface mass density is  $\delta = x$ .

- a.  $(\frac{9}{4}, 0)$
- b.  $(0, \frac{9}{4})$
- c.  $(\frac{4}{9}, 0)$
- d.  $(0, \frac{4}{9})$
- e.  $(81, 0)$

5. Compute  $\iiint x^2 + y^2 dV$  over the region between the cones  $z = \sqrt{x^2 + y^2}$  and  $z = 4 - \sqrt{x^2 + y^2}$ .

- a.  $\frac{8\pi}{3}$
- b.  $\frac{16\pi}{3}$
- c.  $\frac{32\pi}{3}$
- d.  $\frac{16\pi}{5}$
- e.  $\frac{32\pi}{5}$

6. The surface of an apple is given in spherical coordinates by

$$\rho = 2 - 2 \cos \varphi$$

Its volume is given by the integral:

- a.  $V = \int_0^{2\pi} \int_0^{\pi/2} \int_0^{1-\cos \varphi} 1 \, d\rho \, d\varphi \, d\theta$
- b.  $V = \int_0^{2\pi} \int_0^{\pi} \int_0^{2-2\cos \varphi} 1 \, d\rho \, d\varphi \, d\theta$
- c.  $V = \int_0^{2\pi} \int_0^{\pi/2} \int_0^{1-\cos \varphi} \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta$
- d.  $V = \int_0^{2\pi} \int_0^{\pi} \int_0^{2-2\cos \varphi} \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta$
- e.  $V = \int_0^{2\pi} \int_0^{\pi} \int_0^1 (2 - 2 \cos \varphi) \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta$



7. If  $f = x^2 + y^2 - 2z^2$  and  $\vec{F} = (xz, yz, -z^2)$ , which of the following is FALSE?

- a.  $\vec{\nabla} \times \vec{\nabla} f = \vec{0}$
- b.  $\vec{\nabla} \cdot \vec{\nabla} f = 0$
- c.  $\vec{\nabla} \cdot \vec{\nabla} \times \vec{F} = 0$
- d.  $\vec{\nabla}(\vec{\nabla} \cdot \vec{F}) = \vec{0}$
- e. None of the above. They are all true.

8. Let  $f$  be a scalar potential for  $\vec{F} = (y, x, z)$ . Find  $f(2, 2, 2) - f(0, 0, 0)$

- a. 2
- b. 4
- c. 6
- d. 8
- e. 10


---

Work Out: (Points indicated. Part credit possible. Show all work.)

9. (20 points) Draw the region of integration and compute  $\int_0^4 \int_{\sqrt{x}}^2 \sqrt{y^3 + 1} dy dx$  by reversing the order of integration.

10. (20 points) Consider the vector field  $\vec{F} = (x^3, y^3, x^2z + y^2z)$ . First compute  $\vec{\nabla} \cdot \vec{F}$  in rectangular coordinates. Then convert  $\vec{\nabla} \cdot \vec{F}$  into cylindrical coordinates. Finally, compute  $\iiint \vec{\nabla} \cdot \vec{F} dV$  over the solid region below the cone  $z = 4 - \sqrt{x^2 + y^2}$  and above the  $xy$ -plane.

11. (20 points) Compute  $\iint_R x^2 dx dy$  over the diamond shaped region  $R$  bounded by  $xy = 1$ ,  $xy = 9$ ,  $y = x$ ,  $y = 4x$
- HINT: Use the curvilinear coordinates  $(u, v)$  where  $x = uv$  and  $y = \frac{u}{v}$ .



- a. (4 pts) What are the boundaries in terms of  $u$  and  $v$ ?
- b. (6 pts) Find the Jacobian factor  $J = \left| \frac{\partial(x,y)}{\partial(u,v)} \right|$ .
- c. (3 pts) Express the integrand,  $x^2$ , in terms of  $u$  and  $v$ .
- d. (7 pts) Compute the integral.