

MATH 311

Topics in Applied Mathematics I

Lecture 8:

Transpose of a matrix.

Determinants.

Transpose of a matrix

Definition. Given a matrix A , the **transpose** of A , denoted A^T , is the matrix whose rows are columns of A (and whose columns are rows of A). That is, if $A = (a_{ij})$ then $A^T = (b_{ij})$, where $b_{ij} = a_{ji}$.

Examples.
$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}^T = \begin{pmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{pmatrix},$$

$$\begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix}^T = (7, 8, 9), \quad \begin{pmatrix} 4 & 7 \\ 7 & 0 \end{pmatrix}^T = \begin{pmatrix} 4 & 7 \\ 7 & 0 \end{pmatrix}.$$

Properties of transposes:

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(rA)^T = rA^T$
- $(AB)^T = B^T A^T$
- $(A_1 A_2 \dots A_k)^T = A_k^T \dots A_2^T A_1^T$
- $(A^{-1})^T = (A^T)^{-1}$

Definition. A square matrix A is said to be **symmetric** if $A^T = A$.

For example, any diagonal matrix is symmetric.

Proposition For any square matrix A the matrices $B = AA^T$ and $C = A + A^T$ are symmetric.

Proof:

$$B^T = (AA^T)^T = (A^T)^T A^T = AA^T = B,$$

$$C^T = (A + A^T)^T = A^T + (A^T)^T = A^T + A = C.$$

Determinants

Determinant is a scalar assigned to each square matrix.

Notation. The determinant of a matrix

$A = (a_{ij})_{1 \leq i, j \leq n}$ is denoted $\det A$ or

$$\begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix}.$$

Principal property: $\det A \neq 0$ if and only if a system of linear equations with the coefficient matrix A has a unique solution. Equivalently, $\det A \neq 0$ if and only if the matrix A is invertible.

Definition in low dimensions

Definition. $\det(a) = a$, $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$,

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \\ - a_{13}a_{22}a_{31} - a_{12}a_{21}a_{33} - a_{11}a_{23}a_{32}.$$

$$+ : \begin{pmatrix} \boxed{*} & * & * \\ * & \boxed{*} & * \\ * & * & \boxed{*} \end{pmatrix}, \begin{pmatrix} * & \boxed{*} & * \\ * & * & \boxed{*} \\ \boxed{*} & * & * \end{pmatrix}, \begin{pmatrix} * & * & \boxed{*} \\ \boxed{*} & * & * \\ * & \boxed{*} & * \end{pmatrix}.$$

$$- : \begin{pmatrix} * & * & \boxed{*} \\ * & \boxed{*} & * \\ \boxed{*} & * & * \end{pmatrix}, \begin{pmatrix} * & \boxed{*} & * \\ \boxed{*} & * & * \\ * & * & \boxed{*} \end{pmatrix}, \begin{pmatrix} \boxed{*} & * & * \\ * & * & \boxed{*} \\ * & \boxed{*} & * \end{pmatrix}.$$

Examples: 2×2 matrices

$$\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1, \quad \begin{vmatrix} 3 & 0 \\ 0 & -4 \end{vmatrix} = -12,$$

$$\begin{vmatrix} -2 & 5 \\ 0 & 3 \end{vmatrix} = -6, \quad \begin{vmatrix} 7 & 0 \\ 5 & 2 \end{vmatrix} = 14,$$

$$\begin{vmatrix} 0 & -1 \\ 1 & 0 \end{vmatrix} = 1, \quad \begin{vmatrix} 0 & 0 \\ 4 & 1 \end{vmatrix} = 0,$$

$$\begin{vmatrix} -1 & 3 \\ -1 & 3 \end{vmatrix} = 0, \quad \begin{vmatrix} 2 & 1 \\ 8 & 4 \end{vmatrix} = 0.$$

Examples: 3×3 matrices

$$\begin{vmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{vmatrix} = 3 \cdot 0 \cdot 0 + (-2) \cdot 1 \cdot (-2) + 0 \cdot 1 \cdot 3 - \\ - 0 \cdot 0 \cdot (-2) - (-2) \cdot 1 \cdot 0 - 3 \cdot 1 \cdot 3 = 4 - 9 = -5,$$

$$\begin{vmatrix} 1 & 4 & 6 \\ 0 & 2 & 5 \\ 0 & 0 & 3 \end{vmatrix} = 1 \cdot 2 \cdot 3 + 4 \cdot 5 \cdot 0 + 6 \cdot 0 \cdot 0 - \\ - 6 \cdot 2 \cdot 0 - 4 \cdot 0 \cdot 3 - 1 \cdot 5 \cdot 0 = 1 \cdot 2 \cdot 3 = 6.$$

General definition

The general definition of the determinant is quite complicated as there is no simple explicit formula.

There are several approaches to defining determinants.

Approach 1 (original): an explicit (but very complicated) formula.

Approach 2 (axiomatic): we formulate properties that the determinant should have.

Approach 3 (inductive): the determinant of an $n \times n$ matrix is defined in terms of determinants of certain $(n - 1) \times (n - 1)$ matrices.

Axiomatic definition

$\mathcal{M}_{n,n}(\mathbb{R})$: the set of $n \times n$ matrices with real entries.

Theorem There exists a unique function $\det : \mathcal{M}_{n,n}(\mathbb{R}) \rightarrow \mathbb{R}$ (called the determinant) with the following properties:

(D1) if a row of a matrix is multiplied by a scalar r , the determinant is also multiplied by r ;

(D2) if we add a row of a matrix multiplied by a scalar to another row, the determinant remains the same;

(D3) if we interchange two rows of a matrix, the determinant changes its sign;

(D4) $\det I = 1$.

Corollary 1 Suppose A is a square matrix and B is obtained from A applying elementary row operations. Then $\det A = 0$ if and only if $\det B = 0$.

Corollary 2 $\det B = 0$ whenever the matrix B has a zero row.

Hint: Multiply the zero row by the zero scalar.

Corollary 3 $\det A = 0$ if and only if the matrix A is not invertible.

Idea of the proof: Let B be the reduced row echelon form of A . If A is invertible then $B = I$; otherwise B has a zero row.

Remark. The same argument proves that properties (D1)–(D4) are enough to evaluate any determinant.

Row echelon form of a square matrix A :

A 7x7 matrix in row echelon form. The main diagonal elements are represented by squares, and all other elements are asterisks. A blue line connects the bottom-left corners of the squares, forming a staircase pattern that descends from the top-left to the bottom-right.

$$\begin{pmatrix} \square & * & * & * & * & * & * \\ & \square & * & * & * & * & * \\ & & \square & * & * & * & * \\ & & & \square & * & * & * \\ & & & & \square & * & * \\ & & & & & \square & * \\ & & & & & & \square \end{pmatrix}$$

$$\det A \neq 0$$

A 7x7 matrix in row echelon form, similar to the first one, but with two asterisks circled in red. The first red circle is at the element in the third row and fourth column, and the second red circle is at the element in the sixth row and seventh column. A blue line connects the bottom-left corners of the squares, forming a staircase pattern that descends from the top-left to the bottom-right.

$$\begin{pmatrix} \square & * & * & * & * & * & * \\ & \square & * & * & * & * & * \\ & & \square & * & * & * & * \\ & & & \square & * & * & * \\ & & & & \square & * & * \\ & & & & & \square & * \\ & & & & & & \square \end{pmatrix}$$

$$\det A = 0$$

Example. $A = \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}$, $\det A = ?$

Earlier we have transformed the matrix A into the identity matrix using elementary row operations:

- interchange the 1st row with the 2nd row,
- add -3 times the 1st row to the 2nd row,
- add 2 times the 1st row to the 3rd row,
- multiply the 2nd row by -0.5 ,
- add -3 times the 2nd row to the 3rd row,
- multiply the 3rd row by -0.4 ,
- add -1.5 times the 3rd row to the 2nd row,
- add -1 times the 3rd row to the 1st row.

Example. $A = \begin{pmatrix} 3 & -2 & 0 \\ 1 & 0 & 1 \\ -2 & 3 & 0 \end{pmatrix}$, $\det A = ?$

Earlier we have transformed the matrix A into the identity matrix using elementary row operations.

These included two row multiplications, by -0.5 and by -0.4 , and one row exchange.

It follows that

$$\det I = -(-0.5)(-0.4) \det A = (-0.2) \det A.$$

Hence $\det A = -5 \det I = -5.$