

MATH 311

Topics in Applied Mathematics I

**Lecture 33:**

**Area and volume.**

**Multiple integrals.**

Let  $\mathcal{P}$  be the smallest collection of subsets of  $\mathbb{R}^2$  such that it contains all polygons and if  $X, Y \in \mathcal{P}$ , then  $X \cup Y, X \cap Y, X \setminus Y \in \mathcal{P}$ .

**Theorem** There exists a unique function  $\mu : \mathcal{P} \rightarrow \mathbb{R}$  (called the **area function**) that satisfies the following conditions:

- **(positivity)**  $\mu(X) \geq 0$  for all  $X \in \mathcal{P}$ ;
- **(additivity)**  $\mu(X \cup Y) = \mu(X) + \mu(Y)$  if  $X \cap Y = \emptyset$ ;
- **(translation invariance)**  $\mu(X + \mathbf{v}) = \mu(X)$  for all  $X \in \mathcal{P}$  and  $\mathbf{v} \in \mathbb{R}^2$ ;
- $\mu(Q) = 1$ , where  $Q = [0, 1] \times [0, 1]$  is the unit square.

The area function satisfies an extra condition:

- **(monotonicity)**  $\mu(X) \leq \mu(Y)$  whenever  $X \subset Y$ .

Now for any bounded set  $X \subset \mathbb{R}^2$  we let  $\bar{\mu}(X) = \inf_{X \subset Y} \mu(Y)$  and  $\underline{\mu}(X) = \sup_{Z \subset X} \mu(Z)$ . Note that  $\underline{\mu}(X) \leq \bar{\mu}(X)$ . In the case of equality, the set  $X$  is called **Jordan measurable** and we let  $\text{area}(X) = \bar{\mu}(X)$ .

## Area, volume, and determinants

- $2 \times 2$  determinants and plane geometry


Let  $P$  be a parallelogram in the plane  $\mathbb{R}^2$ . Suppose that vectors  $\mathbf{v}_1, \mathbf{v}_2 \in \mathbb{R}^2$  are represented by adjacent sides of  $P$ . Then  $\text{area}(P) = |\det A|$ , where  $A = (\mathbf{v}_1, \mathbf{v}_2)$ , a matrix whose columns are  $\mathbf{v}_1$  and  $\mathbf{v}_2$ .

Consider a linear operator  $L_A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  given by  $L_A(\mathbf{v}) = A\mathbf{v}$  for any column vector  $\mathbf{v}$ . Then  $\text{area}(L_A(D)) = |\det A| \text{area}(D)$  for any bounded domain  $D$ .

- $3 \times 3$  determinants and space geometry

Let  $\Pi$  be a parallelepiped in space  $\mathbb{R}^3$ . Suppose that vectors  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3 \in \mathbb{R}^3$  are represented by adjacent edges of  $\Pi$ . Then  $\text{volume}(\Pi) = |\det B|$ , where  $B = (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ , a matrix whose columns are  $\mathbf{v}_1, \mathbf{v}_2$ , and  $\mathbf{v}_3$ .

Similarly,  $\text{volume}(L_B(D)) = |\det B| \text{volume}(D)$  for any bounded domain  $D \subset \mathbb{R}^3$ .



$\text{volume}(\Pi) = |\det B|$ , where  $B = (\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ . Note that the parallelepiped  $\Pi$  is the image under  $L_B$  of a unit cube whose adjacent edges are  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ .

The triple  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$  obeys the right-hand rule. We say that  $L_B$  **preserves orientation** if it preserves the hand rule for any basis. This is the case if and only if  $\det B > 0$ .

## Riemann sums in two dimensions

Consider a closed coordinate rectangle

$$R = [a, b] \times [c, d] \subset \mathbb{R}^2.$$

*Definition.* A **Riemann sum** of a function  $f : R \rightarrow \mathbb{R}$  with respect to a partition  $P = \{D_1, D_2, \dots, D_n\}$  of  $R$  generated by samples  $t_j \in D_j$  is a sum

$$\mathcal{S}(f, P, t_j) = \sum_{j=1}^n f(t_j) \text{ area}(D_j).$$

The norm of the partition  $P$  is  $\|P\| = \max_{1 \leq j \leq n} \text{diam}(D_j)$ .

*Definition.* The Riemann sums  $\mathcal{S}(f, P, t_j)$  **converge** to a limit  $I(f)$  as the norm  $\|P\| \rightarrow 0$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that  $\|P\| < \delta$  implies  $|\mathcal{S}(f, P, t_j) - I(f)| < \varepsilon$  for any partition  $P$  and choice of samples  $t_j$ .

If this is the case, then the function  $f$  is called **integrable** on  $R$  and the limit  $I(f)$  is called the **integral** of  $f$  over  $R$ .

## Double integral

Closed coordinate rectangle  $R = [a, b] \times [c, d]$   
 $= \{(x, y) \in \mathbb{R}^2 \mid a \leq x \leq b, c \leq y \leq d\}$ .

Notation:  $\iint_R f \, dA$  or  $\iint_R f(x, y) \, dx \, dy$ .

**Theorem 1** If  $f$  is continuous on the closed rectangle  $R$ , then  $f$  is integrable.

**Theorem 2** A function  $f : R \rightarrow \mathbb{R}$  is Riemann integrable on the rectangle  $R$  if and only if  $f$  is bounded on  $R$  and continuous almost everywhere on  $R$  (that is, the set of discontinuities of  $f$  has zero area).

## Fubini's Theorem

Fubini's Theorem allows us to reduce a multiple integral to a repeated one-dimensional integral.

**Theorem** If a function  $f$  is integrable on  $R = [a, b] \times [c, d]$ , then

$$\iint_R f \, dA = \int_a^b \left( \int_c^d f(x, y) \, dy \right) dx = \int_c^d \left( \int_a^b f(x, y) \, dx \right) dy.$$

In particular, this implies that we can change the order of integration in a repeated integral.

**Corollary** If a function  $g$  is integrable on  $[a, b]$  and a function  $h$  is integrable on  $[c, d]$ , then the function  $f(x, y) = g(x)h(y)$  is integrable on  $R = [a, b] \times [c, d]$  and

$$\iint_R g(x)h(y) \, dx \, dy = \int_a^b g(x) \, dx \cdot \int_c^d h(y) \, dy.$$

## Integrals over general domains

Suppose  $f : D \rightarrow \mathbb{R}$  is a function defined on a (Jordan) measurable set  $D \subset \mathbb{R}^2$ . Since  $D$  is bounded, it is contained in a rectangle  $R$ . To define the integral of  $f$  over  $D$ , we extend the function  $f$  to a function on  $R$ :

$$f^{\text{ext}}(x, y) = \begin{cases} f(x, y) & \text{if } (x, y) \in D, \\ 0 & \text{if } (x, y) \notin D. \end{cases}$$

*Definition.*  $\iint_D f \, dA$  is defined to be  $\iint_R f^{\text{ext}} \, dA$ .

In particular,  $\text{area}(D) = \iint_D 1 \, dA$ .



## Integration as a linear operation

**Theorem 1** If functions  $f, g$  are integrable on a set  $D \subset \mathbb{R}^2$ , then the sum  $f + g$  is also integrable on  $D$  and

$$\iint_D (f + g) dA = \iint_D f dA + \iint_D g dA.$$

**Theorem 2** If a function  $f$  is integrable on a set  $D \subset \mathbb{R}^2$ , then for each  $\alpha \in \mathbb{R}$  the scalar multiple  $\alpha f$  is also integrable on  $D$  and

$$\iint_D \alpha f dA = \alpha \iint_D f dA.$$

## More properties of integrals

**Theorem 3** If functions  $f, g$  are integrable on a set  $D \subset \mathbb{R}^2$ , and  $f(x, y) \leq g(x, y)$  for all  $(x, y) \in D$ , then

$$\iint_D f \, dA \leq \iint_D g \, dA.$$

**Theorem 4** If a function  $f$  is integrable on sets  $D_1, D_2 \subset \mathbb{R}^2$ , then it is integrable on their union  $D_1 \cup D_2$ . Moreover, if the sets  $D_1$  and  $D_2$  are disjoint up to a set of zero area, then

$$\iint_{D_1 \cup D_2} f \, dA = \iint_{D_1} f \, dA + \iint_{D_2} f \, dA.$$

## Change of variables in a double integral

**Theorem** Let  $D \subset \mathbb{R}^2$  be a measurable domain and  $f$  be an integrable function on  $D$ . If  $\mathbf{T} = (u, v)$  is a smooth coordinate mapping such that  $\mathbf{T}^{-1}$  is defined on  $D$ , then

$$\begin{aligned} & \iint_D f(u, v) \, du \, dv \\ &= \iint_{\mathbf{T}^{-1}(D)} f(u(x, y), v(x, y)) \left| \det \frac{\partial(u, v)}{\partial(x, y)} \right| \, dx \, dy. \end{aligned}$$

In particular, the integral in the right-hand side is well defined.

**Problem** Evaluate a double integral

$$\iint_P (2x + 3y - \cos(\pi x + 2\pi y)) \, dx \, dy$$

over a parallelogram  $P$  with vertices  $(-1, -1)$ ,  $(1, 0)$ ,  $(2, 2)$ , and  $(0, 1)$ .

Adjacent edges of the parallelogram  $P$  are represented by vectors  $\mathbf{v}_1 = (1, 0) - (-1, -1) = (2, 1)$  and  $\mathbf{v}_2 = (0, 1) - (-1, -1) = (1, 2)$ .

Consider a transformation  $L$  of the plane  $\mathbb{R}^2$  given by

$$L \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} -1 \\ -1 \end{pmatrix} = \begin{pmatrix} 2u + v - 1 \\ u + 2v - 1 \end{pmatrix}$$

(columns of the matrix are vectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$ ). By construction,  $L$  maps the unit square  $[0, 1] \times [0, 1]$  onto the parallelogram  $P$ . The Jacobian matrix  $J$  of  $L$  is the same at

any point:  $J = \frac{\partial(x, y)}{\partial(u, v)} = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ .

Changing coordinates in the integral from  $(x, y)$  to  $(u, v)$  so that  $(x, y) = L(u, v) = (2u + v - 1, u + 2v - 1)$ , we obtain

$$\begin{aligned} & \iint_P (2x + 3y - \cos(\pi x + 2\pi y)) \, dx \, dy \\ &= \iint_{L^{-1}(P)} (7u + 8v - 5 - \cos(4\pi u + 5\pi v - 3\pi)) |\det J| \, du \, dv \\ &= \int_0^1 \int_0^1 3(7u + 8v - 5 + \cos(4\pi u + 5\pi v)) \, du \, dv \\ &= \frac{21}{2} + 12 - 15 + \int_0^1 \int_0^1 3 \cos(4\pi u + 5\pi v) \, du \, dv. \end{aligned}$$

$$\begin{aligned} \text{Further, } & \int_0^1 3 \cos(4\pi u + 5\pi v) \, du = \frac{3}{4\pi} \sin(4\pi u + 5\pi v) \Big|_{u=0}^1 \\ &= \frac{3}{4\pi} (\sin(4\pi + 5\pi v) - \sin(5\pi v)) = 0 \text{ for all } v. \end{aligned}$$

$$\text{It follows that } \iint_P (2x + 3y - \cos(\pi x + 2\pi y)) \, dx \, dy = \frac{15}{2}.$$

## Triple integral

To integrate in  $\mathbb{R}^3$ , volumes are used instead of areas in  $\mathbb{R}^2$ . Instead of coordinate rectangles, basic sets are coordinate boxes (or bricks)  $B = [a_1, b_1] \times [a_2, b_2] \times [a_3, b_3] \subset \mathbb{R}^3$ . Then we can define an integral of a function  $f$  over a measurable set  $D \subset \mathbb{R}^3$ .

Notation:  $\iiint_D f \, dV$  or  $\iiint_D f(x, y, z) \, dx \, dy \, dz$ .

The properties of triple integrals are completely analogous to those of double integrals. In particular, Fubini's Theorem is formulated as follows.

**Theorem** If a function  $f$  is integrable on a brick  $B = [a_1, b_1] \times [a_2, b_2] \times [a_3, b_3] \subset \mathbb{R}^3$ , then

$$\iiint_B f \, dV = \int_{a_1}^{b_1} \left( \int_{a_2}^{b_2} \left( \int_{a_3}^{b_3} f(x, y, z) \, dz \right) dy \right) dx.$$